

ON FIRST ORDER CONGRUENCES OF LINES IN \mathbb{P}^4 WITH GENERICALLY NON-REDUCED FUNDAMENTAL SURFACE*

PIETRO DE POI†

Abstract. In this article we obtain a complete description of the congruences of lines in \mathbb{P}^4 of order one provided that the fundamental surface F is non-reduced (and possibly reducible) at one of its generic points, and their classification under the hypothesis that $(F)_{\text{red}}$ is smooth.

Key words. Algebraic Geometry, focal loci, Grassmannians, congruences

AMS subject classifications. Primary 14M15, 14N15; Secondary 51N35

Introduction. A congruence of d -dimensional linear subspaces in \mathbb{P}^n is an irreducible subvariety (“family”) B' of dimension $n - d$ of the projective Grassmannian $\mathbb{G}(d, n)$. By taking a resolution B of singularities of B' , one can pull back the universal subbundle on $\mathbb{G}(d, n)$, obtaining a smooth \mathbb{P}^d -bundle Λ on B , which has dimension n . Since $\Lambda \subset B \times \mathbb{P}^n$, the second projection induces a morphism $p: \Lambda \rightarrow \mathbb{P}^n$ between manifolds of the same dimension, and one defines the *order* of the family as the degree of p . Geometrically, this is the number of d -planes of the family passing through a general point in \mathbb{P}^n .

The case where B is a surface ($n - d = 2$), and the order is one, was classified by Ziv Ran ([Ran86]), extending the classical work of Kummer ([Kum66]) for $n = 3$ and order at most two.

The case where $n = 4$, $d = 1$ and the order is one was considered by G. Marletta ([Mar09b], [Mar09a]) and, since [De 99], we are trying to complete Marletta’s incomplete classification and to bring it up to modern standards. In fact, in [De 99] we have started the classification of first order congruences of lines in \mathbb{P}^4 , and the first two steps of the classification are in in [De 05] and [De 01].

In this article, we study one of the most difficult and exciting cases, namely when the fundamental surface (see below) of our congruence is non-reduced at one of its generic points. The study of these cases is not only interesting in its own and for its applications to projective geometry, but it will also be useful—hopefully—for mathematical physics, in the theory of systems of conservation laws, see [AF1] and [AF2]. In particular, the results contained here can be used to classify the three dimensional Temple systems of conservation laws which are hyperbolic but *not* strictly hyperbolic, and so, among other things, can complete the results contained in [AF3].

From now on, we will consider always, for simplicity, $d = 1$. The set up for studying congruences of lines is the following: let us denote by R the ramification divisor of p , and by Φ its schematic image, *i.e.* the branch locus, which will be called the *focal locus*.

We can observe that, if the order is at least two, by the purity of the branch locus, Φ contains a hypersurface, whereas, if the order is one, the morphism p is birational. Thus, by Zariski’s main theorem, each component of the focal locus has codimension two, since it is precisely the locus where the fibre dimension is positive; moreover,

*Received March 19, 2005; accepted for publication April 26, 2007.

†Dipartimento di Matematica e Informatica, Università degli studi di Trieste, Via Valerio, 12/1, 34127 Trieste, Italy (depoi@dmi.units.it). This research was partially supported by the DFG Forschungsschwerpunkt “Globalen Methoden in der Komplexen Geometrie”, and the EU (under the EAGER network).

set-theoretically, Φ coincides with the *fundamental locus*, *i.e.* the set of points in \mathbb{P}^n through which there pass infinitely many lines of the congruence.

The *pure fundamental locus* $(\emptyset \neq)F \subset \Phi$ is instead the image under p of the components of R which surjectively project onto B ; this means geometrically that every line of B intersects each component of F .

If the order is one and $n = 4$, it can be proven that either $\dim(\Phi) = 0$, in which case B is a star of lines (*i.e.* the set of lines passing through the point $(\Phi)_{\text{red}}$) or $\dim(\Phi) = 2$ (see Theorem 2.2 in [De 04]). In what follows, we shall only consider congruences of lines in \mathbb{P}^4 such that Φ has pure dimension two. In this situation B' is a three-dimensional subvariety of the trisecant lines to F , by a result of C. Segre (see Proposition 1.1 below). The cases in which F contains a component of dimension one is treated in [De 01], while the case in which the pure fundamental locus F is a (generically) reduced and irreducible surface is in [De 05].

Here we are interested in the cases in which F has some component of dimension two such that F is nonreduced at the corresponding generic point. F will also be called *fundamental surface* and it is either irreducible or reducible. For both these possibilities we give a precise description.

Classically, congruences of order one in \mathbb{P}^4 were treated by G. Marletta in [Mar09b] for the generically reduced case and in [Mar09a] for the rest. In this paper Marletta's classifications are reproved and completed. In particular we deduce the complete list of these congruences provided that all the reduced components of F are smooth.

This article is structured as follows: after giving, in Section 1, the basic definitions, we give examples of all the possible congruences with a non-reduced component of the fundamental locus (finding also a case which is missing in Marletta's list) in Section 2, and in Sections 3 and 4 we prove that no other congruence with this characteristic property can exist. A partial list of these congruences has also been obtained by A. Oblomkov ([O]). If we suppose that all the components of the reduced locus of the pure fundamental locus are smooth, we obtain the complete list in Theorem 0.1.

In this theorem and throughout this article we will use the following notation and conventions: F will be the pure fundamental locus, and moreover we set $D := (F_1)_{\text{red}}$, where F_1 is a two-dimensional component of F such that F is nonreduced at the generic point of F_1 . If $b \in B$, then $\Lambda(b) \subset \mathbb{P}^4$ is the corresponding line of the congruence.

THEOREM 0.1. *If the fundamental surface F of a first order congruence of lines in \mathbb{P}^4 is such that $F_1 \neq \emptyset$ and all the components of $(F)_{\text{red}}$ are smooth, then we have the following possibilities:*

1. $F(= F_1)$ is irreducible, and we have the following cases:
 - (a) $\text{length}(\Lambda(b) \cap D) = 1$, in which case D is a plane and the congruence is as in Example 1; or
 - (b) $\text{length}(\Lambda(b) \cap D) = 2$, D is a rational normal cubic scroll, and the congruence is as in Example 2; or
2. F has two irreducible components, F_1 non-reduced and F_2 reduced, and we have the following cases:
 - (a) $D = (F_1)_{\text{red}}$ is a plane and F_2 is a rational normal cubic scroll, and we have the following cases:
 - i. $D \cap F_2$ is either a line or a conic and the congruence is as in Example 4; or
 - ii. $D \cap F_2$ is a (smooth) conic and the congruence is as in Example 3; or

(b) F_2 is a plane, and $D = (F_1)_{\text{red}}$ is a rational normal cubic scroll; $D \cap F_2$ is a smooth conic, unisecant to D , and the congruence is as in Example 5.

Vice versa a family of lines of dimension three constructed as in each of the cases of the theorem is a first order congruence in \mathbb{P}^4 .

Acknowledgements. This article has been highly improved by many discussions with and suggestions of F. Catanese. I want to thank him, E. Mezzetti and E. Arrondo for their help in the elaboration of this paper.

I would also like to thank A. A. Oblomkov for the useful discussions on the subject.

1. Notation, definitions and preliminary results. We will work with schemes and varieties over the complex field \mathbb{C} , with standard notation and definitions as in [Har77]. In this article, a *variety* will always be projective and irreducible. We refer to [De 01] and [De 03] for general results and references about families of lines, focal diagrams and congruences, and to [GH78] for notations about Schubert cycles. So, we denote by σ_{a_0, a_1} the Schubert cycle of the lines in \mathbb{P}^4 contained in a fixed $(4 - a_1)$ -dimensional subspace $H \subset \mathbb{P}^4$ and which meet a fixed $(3 - a_0)$ -dimensional subspace $\Pi \subset H$. We recall that a *congruence of lines* in \mathbb{P}^4 is a (flat) family (Λ, B, p) of lines in \mathbb{P}^4 obtained as the pull-back of the universal family under the desingularization of a subvariety B' of dimension three of the Grassmannian $\mathbb{G}(1, 4)$. So, $\Lambda \subset B \times \mathbb{P}^4$ and p is the restriction of the projection $p_1: B \times \mathbb{P}^4 \rightarrow B$ to Λ , while we will denote the restriction of $p_2: B \times \mathbb{P}^4 \rightarrow \mathbb{P}^4$ by f . $\Lambda_b := p^{-1}(b)$, ($b \in B$) is a line of the family and $f(\Lambda_b) =: \Lambda(b)$ is a line in \mathbb{P}^4 . Λ is smooth of dimension three: therefore we can define the *focal divisor* $R \subset \Lambda$ as the ramification divisor of f . The *focal locus* $\Phi = f(R) \subset \mathbb{P}^4$, is the scheme theoretic image of R .

In this article, we study the *first order congruences* of lines, *i.e.* congruences B for which through a general point in \mathbb{P}^4 there passes only one line of B ; or, equivalently, we can write

$$(1) \quad [B'] = \sigma_3 + a\sigma_{2,1},$$

where $[B']$ is the rational equivalence class of B' —as a linear combination of Schubert cycles of the Grassmannian; so, the *class* a is the degree of the ruled surface generated by the lines of the congruence which belong to a general \mathbb{P}^3 (*i.e.* the intersection number $[B'] \cdot \sigma_{2,1}$). Given a first order congruence B , through a focal point there will pass infinitely many lines in B , *i.e.* a focal point is a *fundamental point* and the focal locus coincides set-theoretically with the *fundamental locus*.

An important result, due to Corrado Segre, is the following (see [De 01], Proposition 1 for a proof):

PROPOSITION 1.1 (C. Segre, [Seg88]). *On every line $\Lambda(b) \subset \mathbb{P}^4$ of the family, the focal locus Φ either coincides with the whole $\Lambda(b)$ —in which case $\Lambda(b)$ is called focal line—or is a zero dimensional scheme of $\Lambda(b)$ of length three.*

Let us observe that, if the order is one, the morphism p is birational, thus, by Zariski main theorem (finite plus birational cover of a normal variety is an isomorphism), the focal locus is not a hypersurface, so $\dim(\Phi) \leq 2$. Actually, we have proven in [De 04] that $\dim(\Phi) = 2$ if B is not a star of lines.

Here we are interested in first order congruences of lines in \mathbb{P}^4 for which the focal locus Φ has pure dimension two, so the “important” component of Φ is the *fundamental 2-locus* (see [De 01], [De 03] and [De 04]), which is characterized by the fact that it is formed by the components of Φ of pure dimension two such that the general line of the

congruence meets it in a zero dimensional scheme (see [De 01], Proposition 2). The fundamental 2-locus will be called *pure fundamental locus* or, in what follows, simply *fundamental surface* and it will be indicated with F (For the other “non-important” components of Φ , *i.e.* the *parasitic planes*, see [De 05]).

In the rest of the paper, we need more notation. To a first order congruence B we associate:

- the hypersurface V_Π in \mathbb{P}^4 given by the lines of B which intersect a general plane $\Pi \subset \mathbb{P}^4$. So, $\deg(V_\Pi) = 1 + a$, where a is the class of B (see (1));
- the surface Σ_H —of degree a —of the lines of B contained in a general hyperplane H ; we can think of it as the image of the \mathbb{P}^1 -bundle over the curve $\Gamma_H \subset B$, obtained by pulling back the intersection of B' with the Schubert variety of the lines contained in H .
- we set also $C_H := (F)_{\text{red}} \cap H (\subset \Sigma_H)$.

REMARK. First of all we note that $\text{Sing}(\Sigma_H) \subset C_H$, since through a singular point P of the surface there pass more than a line of the congruence (possibly infinitely near). Besides, for the generality of H and dimensional reasons, Σ_H cannot be contained in F .

LEMMA 1.2. *Let D be a surface in \mathbb{P}^4 and P a smooth point in D such that, for a general point Q in D , the tangent planes $\mathbb{T}_{P,D}$ and $\mathbb{T}_{Q,D}$ do not span \mathbb{P}^4 . Then the surface D is degenerate.*

Proof. Take projective coordinates such that the local parametrization of D at P is $F := (1 : x : y : f(x, y) : g(x, y))$, where f, g vanish of order ≥ 2 for $x = y = 0$. Our statement is that the 6 vectors e_0, e_1, e_2 and F, F_x, F_y have rank (at most) 4. This simply means that the three vectors $(f, g), (f_x, g_x), (f_y, g_y)$ have rank (at most) 1, and implies that the map $(x : y) \mapsto (f : g)$ has constant image in \mathbb{P}^1 . Without loss of generality we may then assume $g \equiv 0$, hence D is degenerate. \square

PROPOSITION 1.3. *The congruence B of the tangent lines to a surface $D \subset \mathbb{P}^4$ cannot have order one.*

Proof. First we note that if D is a cone, B has order zero. Moreover, D is contained in the focal locus F of B . If $P \in \mathbb{T}_{P_1,D} \cap \mathbb{T}_{P_2,D}$, then $P \in F$, since two lines pass through it.

We cannot have $\dim(\mathbb{T}_{P_1,D} \cap \mathbb{T}_{P_2,D}) = 1$ by the preceding lemma, so we suppose that $\mathbb{T}_{P_1,D} \cap \mathbb{T}_{P_2,D} =: Q$ is a point and that B is a first order congruence. Now, if $\dim(\mathbb{T}_{P,D} \cap F) = 0$ for the general $P \in D$, then $Q \in (\mathbb{T}_{P,D} \cap F)$, since $\mathbb{T}_{P,D} \cap \mathbb{T}_{P',D} \in \mathbb{T}_{P,D} \cap F$ as P' varies in D . Therefore Q is fixed; but then the dual variety D^* would be degenerate, and D would be a cone (this argument holds only on a field of characteristic zero).

Instead $\dim(\mathbb{T}_{P,D} \cap F) = 1$ means that there is a family of dimension two of planes curves contained in F . But the surfaces with this property in \mathbb{P}^4 are classified in C. Segre’s Theorem (see [MP97], Theorem 4 for a modern proof) and they are the projected Veronese surface, the rational normal smooth cubic scroll and the cones. If D is this surface, we conclude by observing that couples of tangent planes to either the Veronese surface or the cubic ruled surface do not meet in a point in D .

Otherwise, there is another nondegenerate (if it is degenerate, also D is degenerate) surface $D' \subset F$ which is in the list of C. Segre’s Theorem, and the congruence B is given by the lines which are tangent to B and these lines moreover meet D' . In particular, on every line r of B , the three foci of r are a double point in D and

a single point in D' . D' cannot be a cone, since again D^* would be degenerate; in fact all the tangent planes would pass through the vertex of D' . Then D' is either the projected Veronese surface or the cubic scroll, and in both these cases the plane curves are irreducible conics $C \subset \mathbb{T}_{P,D}$. We recall that the lines of B contained in $\mathbb{T}_{P,D}$ form the pencil of lines through the point $P \in D$, \mathbb{P}_P^1 . But if $C \subset \mathbb{T}_{P,D}$ is the conic, and $\ell_P \in \mathbb{P}_P^1$, then either the two points $\ell_P \cap C$ are focal, but then $\ell_P \subset F$ and therefore every line in B is focal, or only one of the two points is focal for ℓ_P . But then, we would obtain a regular map from $\mathbb{P}_P^1 \cong \mathbb{P}^1$ to C which is injective not but surjective. \square

COROLLARY 1.4. Σ_H does not contain a component S_H which is the surface of the tangents to a curve C .

Proof. If it were so, $C \subset C_H$ and then B would be given by the family of tangent lines to $D = (F_1)_{\text{red}}$. \square

LEMMA 1.5. Σ_H does not contain a component S_H which is a cone.

Proof. If every S_H would be a cone, since this cone spans the hyperplane H , we would have ∞^4 such cones, whence ∞^5 pairs (S_H, L) , where L is a line of the cone S_H . Since such a line L belongs to ∞^2 hyperplanes, it follows that we get in this way ∞^3 lines, hence the whole congruence of lines.

Consider the irreducible variety S image of the rational map $H \rightarrow V_H$. We know that $\dim(S) \leq 2$, hence a general $V = V_H$ is a vertex of at least ∞^2 cones, and *a fortiori* there are at least ∞^2 lines of the congruence which pass through V . Hence we obtain that S is a 1-dimensional component of the focal locus, which is excluded by our assumption. \square

LEMMA 1.6. Σ_H does not contain a component S_H which is a quadric.

Proof. If S_H is a quadric, it must be smooth, by the preceding lemma.

If S_H is a smooth quadric, only one of the two families of lines contained in it can be contained in B , otherwise $S_H \subset F$, and this is absurd by varying H .

Let ℓ be a line of the congruence contained in S_H . If ℓ' is another general line of the congruence not contained in H , the hyperplane $H' := \overline{\ell'\ell}$ determines another quadric $S_{H'}$. By construction, $S_H \cap S_{H'} = \ell \cup \ell'$, and $\ell' \cap \ell'' \neq \emptyset$; therefore ℓ' intersects S_H , which is absurd since every line of B should intersect S_H . This again implies $S_H \subset F$, the same contradiction. \square

2. The examples. Since the congruences we are studying are rather complicated to construct, we first give some examples describing them, and afterwards we will show that these examples exhaust all the possible congruences.

REMARK. We observe first that if a component of the focal locus is contained set-theoretically in a hyperplane H , then the lines of the congruence contained in H form a first order congruence. Therefore a way of constructing congruences in \mathbb{P}^4 (or in general in \mathbb{P}^n) is to reverse this by considering a one-dimensional linear system of hyperplanes in \mathbb{P}^4 such that in each of them there is a first order congruence of lines in \mathbb{P}^3 , and such that these congruences vary algebraically.

Then, we start recalling the classification of first order congruences in \mathbb{P}^3 : we follow the notation used in [De 04]: let ℓ be a fixed line in \mathbb{P}^3 , then \mathbb{P}_ℓ^1 is the set of the planes containing ℓ . Let ϕ be a general nonconstant morphism from \mathbb{P}_ℓ^1 to ℓ and let Π be a general element in \mathbb{P}_ℓ^1 . We define $\mathbb{P}_{\phi(\Pi),\Pi}^1$ as the pencil of lines passing through the point $\phi(\Pi)$ and contained in Π .

If \mathcal{B} is a first order congruence of lines in \mathbb{P}^3 with focal locus \mathcal{F} , then

1. $P := \mathcal{F}_{\text{red}}$ is a point and \mathcal{B} is the star of lines passing through P ; or
2. \mathcal{F} is a rational normal curve in \mathbb{P}^3 and \mathcal{B} is the family of secant lines of \mathcal{F} ; or
3. $\ell := \mathcal{F}_{\text{red}}$ is a line, and the congruence is $\cup_{\Pi \in \mathbb{P}_k^1} \mathbb{P}_{\phi(\Pi), \Pi}^1$; or
4. $\mathcal{F} = \mathcal{F}_1 \cup \mathcal{F}_2$ where \mathcal{F}_1 is a line and \mathcal{F}_2 is a rational curve, such that $\text{length}(\mathcal{F}_1 \cap \mathcal{F}_2) = \text{deg}(\mathcal{F}_2) - 1$ and \mathcal{B} is the family of lines meeting \mathcal{F}_1 and \mathcal{F}_2 .

In order to state the examples of congruences in \mathbb{P}^4 , we need some more notations:

- if $H \subset \mathbb{P}^4$ is a linear space, H^* denotes its dual, *i.e.* the linear space of the hyperplanes in H ; if $K \subset H$ is another subspace, we set $K_H^\vee := \{\Pi \in H^* \mid \Pi \supset K\} \subset H^*$ (or simply K^\vee if $H = \mathbb{P}^4$) *i.e.* K_H^\vee is the projective dual subspace of K in H ; k will denote an element in K_H^\vee ;
- $|C|$ denotes the complete linear system associated to the divisor C ;
- a general (rational) map from X to Y will be denoted by $\phi_{X,Y}$;
- if $X, Y \subset \mathbb{P}^4$, $J(X, Y) \subset \mathbb{G}(1, 4)$ is $J(X, Y) := \{\ell \in \mathbb{G}(1, 4) \mid \ell \cap X \neq \emptyset \neq \ell \cap Y\}$; \overline{X} and $\langle X \rangle$ are, respectively, the (Zariski) closure and the span of X .
- We recall that we set $D := (F_1)_{\text{red}}$, where F_1 is the component of F which is non-reduced.
- Finally, in the case in which $F = F_1 \cap F_2$ is also reducible, we denote $C_{H,1} := (F_1)_{\text{red}} \cap H$ and $C_{H,2} := F_2 \cap H$, so $C_H = C_{H,1} \cup C_{H,2}$. k_i is the algebraic multiplicity of $C_{H,i}$ in Σ_H .

EXAMPLE 1. Let us see how to construct a congruence of lines in \mathbb{P}^4 from the case (3) in \mathbb{P}^3 , with set-theoretically linear focal locus, *i.e.* $D := F_{\text{red}}$ is a plane. We want to follow the remark above, so we need a pencil of hyperplanes, and therefore we have to fix a plane $D \subset \mathbb{P}^4$, so that the pencil is D^\vee . Now, for every element $d \in D^\vee$, we have to find a line $\ell_d \subset D$, in order to construct the congruence of lines in $d \cong \mathbb{P}^3$. Since we do not want (embedded) components of dimension one, we suppose that the line ℓ_d is not the same for all the elements in D^\vee . In brief, we have fixed a morphism $\phi := \phi_{D^\vee, D^*}$.

Now, we have to construct a congruence in every $d \in D^\vee$, as in case (3), and therefore we have to fix a morphism $\psi_d := \phi_{\phi(d)_d^\vee, \phi(d)}$; moreover, we are in the algebraic category, and so we assume that we have an algebraic family of morphisms $\{\psi_d\}_{d \in D^\vee}$.

Now it is clear that our congruence in \mathbb{P}^4 is formed by the lines of the pencils $(\psi_d(h))_h^\vee$ as h varies in $\phi(d)_d^\vee$ and d in D^\vee , *i.e.*

$$B = \overline{\cup_{d \in D^\vee} \cup_{h \in \phi(d)_d^\vee} (\psi_d(h))_h^\vee};$$

an easy calculation (or one can see Theorem 8, case 3 of [De 01] where this congruence is also introduced) shows that the bidegree of Λ is $(1, d_1 d_2 m + 1)$, where $d_1 := \text{deg}(\phi)$, $d_2 := \text{deg}(\psi_d)$ and $m := \text{deg}(\phi(D^\vee))$ and if r is the general line in D , then $\text{deg}(f^{-1}(r)) = d_1 d_2 m$.

We note that we could construct more first order congruences if we associate to d a general line in \mathbb{P}^4 (and not contained in D). Actually, in this case the reduced focal locus $D := F_{\text{red}}$ is a ruled surface and it is easy to show that B is given by a family of secant lines to D , and in fact this case is contemplated in the example which follows.

EXAMPLE 2. We will see now how to construct a first order congruence of lines B such that it is a subfamily of the secant lines to the reduced locus $D := (F)_{\text{red}}$. Let us suppose that D is a rational normal cubic scroll in \mathbb{P}^4 , *i.e.* either a non-degenerate cone or a rational normal (smooth) scroll of type (1, 2). In order to do so, let us

consider a curve C_0 which is either the zero section (see Proposition V.2.8 of [Har77]) if the scroll is smooth, *i.e.* an irreducible conic, or a pair of lines if D is a cubic cone, and then let us fix a map $\phi_{|C_0|,F}$ which associates to the curve $C \in |C_0|$ a point $P_C := \phi_{|C_0|,F}(C) \in C$; then we define a congruence as

$$B := \overline{\cup_{C \in |C_0|} (P_C)_{(C)}^\vee};$$

it is clearly a first order congruence of lines and moreover, if $k = \deg(\phi_{|C_0|,F})$ then the class of B is $2k$.

If we want to obtain congruences of lines in \mathbb{P}^4 from case (4), we have in fact many possibilities, since we may suppose either that the line or the rational curve is in the non-reduced locus. In particular, we will see that in Example 3 we have the case of the curve (a conic, indeed), and in Example 4 the line; in both cases the non-reduced locus is a plane.

EXAMPLE 3. So, let us first suppose that $D := (F_1)_{\text{red}}$ is a plane, and that the rational curves of case (4) are conics contained in D . Then, we also suppose that F_2 is a non-degenerate rational scroll of type $(1, d)$ and by what we are supposing, the ruling lines must be the lines of case (4); therefore, $D \cap F_2$ is a unisecant curve (of degree d) in F_2 . Now, let $|H_1| \cong \mathbb{P}^1$ be the linear system of the lines of the ruling in F_2 and $|C_2|$ the linear system of conics in D ; then the congruence is defined as follows: fix a morphism $\phi := \phi_{|H_1|,|(C_2)}$ such that $\phi_{|H_1|,|(C_2)}(\ell) \cap \ell \neq \emptyset$; then we set

$$B := \overline{\cup_{\ell \in |H_1|} \cup_{r \in J(\ell, \phi_{|H_1|,|(C_2)}(\ell))} r};$$

we also have that $k_1 = (d + 1) \deg(\phi) \deg(\text{Im}(\phi))$, $k_2 = 2$, and so $a = 2 + \deg(F_2) \deg(\phi) \deg(\text{Im}(\phi))$.

EXAMPLE 4. Again, let us first suppose that $D := (F_1)_{\text{red}}$ is a plane, but now we suppose that in it there are contained the lines of case (4). Now, we suppose also that F_2 is a non-degenerate rational surface with sectional genus zero (*i.e.* either a rational scroll or the projected Veronese surface); let $d \in D^\vee$ and let $|H_1|$ be the corresponding linear system of the curves $H_1 := \overline{(F_2 \cap d) \setminus D}$; then the congruence is given in this way: fix a morphism $\phi := \phi_{|H_1|,D^*}$ such that $\text{length}(\phi_{|H_1|,D^*}(C_2) \cap C_2) = \deg(C_2) - 1$; then we set

$$B := \overline{\cup_{C_2 \in |H_1|} \cup_{r \in J(C_2, \phi_{|H_1|,D^*}(C_2))} r};$$

we have also that $k_1 = \deg(F_2) \deg(\phi) \deg(\text{Im}(\phi))$ and $k_2 = 1$, and so $a = 1 + \deg(F_2) \deg(\phi) \deg(\text{Im}(\phi))$.

In the last example the focal locus is reducible, and the non-reduced component will be a rational ruled surface. Also this, as Example 1, is obtained from case (3) in \mathbb{P}^3 :

EXAMPLE 5. Now F_2 is a plane, suppose that $D = (F_1)_{\text{red}}$ is a non-degenerate rational scroll of type $(1, c)$ and that $D \cap F_2$ is a unisecant curve in D of degree c . Let $|H_1| \cong \mathbb{P}^1$ be the linear system of the lines of the ruling in D ; the congruence is obtained in the following way: $\forall L \in |H_1|$, fix a morphism $\psi := \phi_{L_{(L, F_2)}, L}$; then we set

$$B := \overline{\cup_{L \in |H_1|} \cup_{l \in L_{(L, F_2)}^\vee} \psi(l)_l^\vee};$$

we have also that $k_1 = \deg(\psi)$ and $k_2 = c + 1$, and so $a = \deg(D) + \deg(\psi)$.

3. The case of an irreducible fundamental surface. We start considering the case in which F is irreducible; since we are interested in the cases in which F is non-reduced at a generic point, we have two possibilities: either the general line $\Lambda(b)$ intersects F in a fat point of length three, or it intersects F in two points, one of which is a fat point of length two and the other is a simple point.

3.1. The case in which $\text{length}(\Lambda(b) \cap (F)_{\text{red}}) = 1$. If the general line of the congruence intersects F in only a fat point, we can prove the following:

THEOREM 3.1. *With notation as in the introduction, if $\text{length}(\Lambda(b) \cap D) = 1$, then D is a plane and the congruence is given in as in Example 1.*

Vice versa a family of lines constructed in this way is a first order congruence.

Proof. Let us consider the surface Σ_H of degree a associated to a general hyperplane H ; it contains the curve C_H with some algebraic multiplicity, say k .

If $L \subset \Sigma_H$ is a (general) line, and $l \in L_H^\vee$ a general plane, then $l \cap \Sigma_H = L \cup C$, where C is a curve of degree $a - 1$, with a point of multiplicity $k - 1$ in $L \cap F$. C intersects L in only another point, *i.e.* the sole point in which l is tangent to the irreducible component S_L of Σ_H which contains L . We recall in fact that a general tangent plane to a surface S is tangent at only one point in S but in the case in which S is a surface of tangents to a curve or a cone (see [Zak93]), and S_L can be neither a surface of tangents to a curve nor a cone, by Corollary 1.4, and Lemmas 1.5 and 1.6; therefore we have that $a = k + 1$.

I claim now that C_H is a plane curve. Indeed, observe first that any (bi)secant line to C_H through a general point in L meets Σ_H in at least $k + k + 1$ points counting multiplicities. Since Σ_H has degree $a = k + 1 < 2k + 1$, it follows that such a line is contained in Σ_H . If C_H is not a plane curve, it follows that through a general point of it there pass at least two lines of it (the line L and one of the bisecant lines mentioned above). But it is well known that a ruled surface with such a property is necessarily a quadric surface, which contradicts Lemma 1.6.

Therefore F is a degenerate surface; actually, it must be a plane: in fact, if P is a general point in $\langle C_H \rangle$, then through it there passes only one line ℓ_P of the congruence, and its focal point $\ell_P \cap D$ is contained in C_H .

By Theorem 8 of [De 01], in which we classified all first order congruences in \mathbb{P}^4 whose fundamental locus is set-theoretically linear, we finish the proof. \square

3.2. The case in which $\text{length}(\Lambda(b) \cap (F)_{\text{red}}) = 2$. Of these congruences we give a complete classification in the following

THEOREM 3.2. *If B is a first order congruence of lines given by a subfamily of the secant lines to the reduced locus $D := (F)_{\text{red}}$ of its pure fundamental locus F , then D is a non-degenerate cubic scroll in \mathbb{P}^4 and the congruence is given as in Example 2.*

Vice versa a B constructed in this way is a first order congruence such that $\text{length}(\Lambda(b) \cap D) = 2$.

Proof. Let us consider the surface Σ_H of degree a associated to a general hyperplane H ; it contains the curve C_H with some (algebraic) multiplicity, say k' .

As in the proof of the preceding theorem, we fix $L \subset \Sigma_H$, and $l \in L_H^\vee$. Then $l \cap \Sigma_H = L \cup C$, where C is a curve of degree $a - 1$, with two points of multiplicity $k' - 1$ in $L \cap F$. As before, C intersects L in only another point, by Corollary 1.4, and Lemmas 1.5 and 1.6 and $a = 2k'$.

In this case, C_H must be a twisted cubic: in fact, we recall that the twisted cubic is the sole irreducible curve with only one apparent double point, and so if C_H is not

the twisted cubic, the secant lines to C_H passing through a point $P \in L$ which are distinct from L are contained in Σ_H , by degree reasons, and so Σ_H must be a quadric, contradicting Lemma 1.6. Therefore we have only a secant line to C_H through P , L , and C_H can only be a twisted cubic and $(F)_{\text{red}}$ a rational normal scroll of degree three in \mathbb{P}^4 . We recall that the planes of the conics in the cubic scrolls cover \mathbb{P}^4 : therefore the lines of the congruence contained in one of these planes can be a congruence of order one and we are in the case of the assertion of the theorem, or it is a finite set. But the last case cannot occur: in fact, every secant line determines in a unique way a conic in the scroll, so we would have that $\dim(B) = 2$.

$k' = \deg(\phi|_{C_0, F}) := k$ since k' is the degree of the cone of the lines in B through a general point in the scroll. So, by what we said above, $k = k'$ and $a = 2k$.

Vice versa, it is easy to see that a congruence constructed in such a way, has order one. \square

REMARK. We note that the case of the cone in the preceding theorem was not considered by G. Marletta in [Mar09a].

Moreover, the only possible smooth surface in Theorem 3.2 is the rational normal cubic scroll in \mathbb{P}^4 , so we have done case (1b) of Theorem 0.1.

4. The case of a reducible and non-reduced fundamental surface. Now we consider the case in which F is reducible but not (generically) reduced; we have only one possibility: the general line $\Lambda(b)$ intersects a component F_1 of F in a fat point of length two, and so F_1 is non-reduced, and the other component F_2 (which is reduced) of F in a simple point.

We recall we have denoted $C_{H,1} := (F_1)_{\text{red}} \cap H$ and $C_{H,2} := F_2 \cap H$, so $C_H = C_{H,1} \cup C_{H,2}$ and k_i is the algebraic multiplicity of $C_{H,i}$ in Σ_H .

We start with

LEMMA 4.1. *Either $D := (F_1)_{\text{red}}$ or F_2 is a plane. Moreover, the class of the congruence is $a = k_1 + k_2$.*

Proof. As in the proof of the preceding two theorems, we fix $L \subset \Sigma_H$, and $l \in L_H^\vee$. Then $l \cap \Sigma_H = L \cup C$, where C is a curve of degree $a - 1$, with a point of multiplicity $k_1 - 1$ and a point of multiplicity $k_2 - 1$ in $L \cap F$. C intersects L in only another point, by Corollary 1.4, and Lemmas 1.5 and 1.6, and $a = k_1 + k_2$. Reasoning as before, through a general point in L there will not pass another joining line $C_{H,1}$ and $C_{H,2}$, and so these lines generate a first order congruence in H . Then, since all the first order congruences of lines in \mathbb{P}^3 are classified (see, for example, [De 04] Theorem 0.1), we have that one of these curves is a line. \square

PROPOSITION 4.2. *Let B be a first order congruence with non-reduced and reducible fundamental locus F , whose non-reduced component is F_1 and the other is F_2 . Then we have the following possibilities:*

1. $D := (F_1)_{\text{red}}$ is a plane, and we have the following cases:
 - (a) F_2 is a non-degenerate rational scroll of type $(1, d)$ and $D \cap F_2$ is a unisecant line in F_2 and the congruence is as in Example 3; or
 - (b) F_2 is a non-degenerate rational surface with sectional genus zero (i.e. either a rational scroll or the projected Veronese surface) and the congruence is as in Example 4; or
2. F_2 is a plane, and $D = (F_1)_{\text{red}}$ is a non-degenerate rational scroll of type $(1, c)$, $D \cap F_2$ is a unisecant curve in D of degree c and the congruence is as in Example 5.

Proof. As we observed in the remark in Section 2, if we have a linear component in the pure fundamental locus, then the lines of B contained in a general hyperplane containing this component give a first order congruence in \mathbb{P}^3 . So, if F_2 is this plane, the congruence induced in an $f_2 \in F_2^Y$ has a non-reduced line as a fundamental locus, given by $F_1 \cap f_2$ (see for example Theorem 0.1, case (1b) of [De 04]), from which we obtain case (2).

If instead $D = (F_1)_{\text{red}}$ is a plane, then the congruence induced in a $d \in D^V$ is a congruence with two fundamental curves, L and C , L is a line and if $\deg(C) = c$, then $\text{length}(C \cap L) = c - 1$ (Theorem 0.1, case (2) of [De 04]). If $L \subset F_2$, we have case (1a), while if $L \subset (F_1)_{\text{red}}$ we have case (1b).

The calculations of k_1 and k_2 are immediate once one remembers that these numbers are the degrees of the cones of the lines of B passing through a general point in, respectively, F_1 and F_2 . The class follows from Lemma 4.1. \square

From this, we easily obtain Theorem 0.1, case (2).

REFERENCES

- [AF1] S. I. AGAFONOV AND E. V. FERAPONTOV, *Systems of conservation laws from the point of view of the projective theory of congruences*, Izv. Ross. Akad. Nauk Ser. Mat., 60 (1996), pp. 3–30.
- [AF2] ———, *Theory of congruences and systems of conservation laws*, J. Math. Sci. (New York), 94 (1999), pp. 1748–1794, Geometry, 4.
- [AF3] ———, *Systems of conservation laws of Temple class, equations of associativity and linear congruences in \mathbb{P}^4* , Manuscripta Math., 106 (2001), pp. 461–488.
- [De 99] P. DE POI, *On First Order Congruences of Lines*, Ph.D. thesis, SISSA-ISAS, October 1999.
- [De 01] ———, *On first order congruences of lines of \mathbb{P}^4 with a fundamental curve*, Manuscripta Math., 106 (2001), pp. 101–116.
- [De 03] ———, *Threefolds in \mathbb{P}^5 with one apparent quadruple point*, Comm. Algebra, 31 (2003), pp. 1927–1947.
- [De 04] ———, *Congruences of lines with one-dimensional focal locus*, Portugaliae Mathematica (N.S.), 61 (2004), pp. 329–338.
- [De 05] ———, *On first order congruences of lines in \mathbb{P}^4 with irreducible fundamental surface*, Mathematische Nachrichten, 278 (2005), pp. 363–378.
- [GH78] P. GRIFFITHS AND J. HARRIS, *Principles of Algebraic Geometry*, John Wiley & Sons, 1978.
- [Har77] R. HARTSHORNE, *Algebraic Geometry*, Graduate Texts in Mathematics, no. 52, Springer-Verlag, 1977.
- [Kum66] E. E. KUMMER, *Über die algebraischen Strahlensysteme, insbesondere über die der ersten und zweiten Ordnung*, Abh. K. Preuss. Akad. Wiss. Berlin (1866), pp. 1–120, also in E. E. Kummer, *Collected Papers*, Springer Verlag, 1975.
- [Mar09a] G. MARLETTA, *Sopra i complessi d'ordine uno dell' S_4* , Atti Accad. Gioenia, Serie V, Catania, III (1909), pp. 1–15, Memoria II.
- [Mar09b] ———, *Sui complessi di rette del primo ordine dello spazio a quattro dimensioni*, Rend. Circ. Mat. Palermo, XXVIII (1909), pp. 353–399.
- [MP97] E. MEZZETTI AND D. PORTELLI, *A tour through some classical theorems on algebraic surfaces*, An. Șt. Univ. Ovidius Constanța, 5 (1997), pp. 51–78.
- [O] A. OBLOMKOV, *private communication*.
- [Ran86] Z. RAN, *Surfaces of order 1 in Grassmannians*, J. Reine Angew. Math., 368 (1986), pp. 119–126.
- [Seg88] C. SEGRE, *Un'osservazione sui sistemi di rette degli spazî superiori*, Rend. Circ. Mat. Palermo, II (1888), pp. 148–149.
- [Zak93] F. L. ZAK, *Tangents and Secants of Algebraic Varieties*, American Mathematical Society, Providence, RI, 1993, translated from the Russian manuscript by the author.