

## EQUIVARIANT MINIMAL SURFACES IN $\mathbb{C}\mathbb{H}^2$ AND THEIR HIGGS BUNDLES\*

JOHN LOFTIN<sup>†</sup> AND IAN MCINTOSH<sup>‡</sup>

**Abstract.** This paper gives a construction for all minimal immersions  $f$  of the Poincaré disc into the complex hyperbolic plane  $\mathbb{C}\mathbb{H}^2$  which are equivariant with respect to an irreducible representation  $\rho$  of a hyperbolic surface group into  $PU(2, 1)$ . We exploit the fact that each such immersion is a twisted conformal harmonic map and therefore has a corresponding Higgs bundle. We identify the structure of these Higgs bundles and show how each is determined by properties of the map, including the induced metric and a holomorphic cubic differential on the surface. We show that the moduli space of pairs  $(\rho, f)$  is a disjoint union of finitely many complex manifolds, whose structure we fully describe. The holomorphic (or anti-holomorphic) maps provide multiple components of this union, as do the non-holomorphic maps. Each of the latter components has the same dimension as the representation variety for  $PU(2, 1)$ , and is indexed by the number of complex and anti-complex points of the immersion. These numbers determine the Toledo invariant and the Euler number of the normal bundle of the immersion. We show that there is an open set of quasi-Fuchsian representations of Toledo invariant zero for which the minimal surface is unique and Lagrangian.

**Key words.** Minimal surface, Higgs bundle, complex hyperbolic plane.

**Mathematics Subject Classification.** 20H10, 53C43, 58E20.

**1. Introduction.** In this article we provide a complete classification of  $\rho$ -equivariant minimal immersions  $f : \mathcal{D} \rightarrow \mathbb{C}\mathbb{H}^2$  and a parametrisation for their moduli space as a union of complex manifolds. Here  $\mathcal{D}$  is the Poincaré disc and  $\rho$  is an irreducible (or more generally, reductive) representation of a hyperbolic surface group (i.e., the fundamental group  $\pi_1\Sigma$  of a closed orientable surface  $\Sigma$  of genus at least two) into the group  $PU(2, 1) = U(2, 1)/\text{centre}$ . Recall that a minimal immersion of a surface is the same thing as a conformal harmonic map. To say  $f$  is  $\rho$ -equivariant means it intertwines the action of a Fuchsian group on  $\mathcal{D}$  with the action of  $\rho$  on the complex hyperbolic plane  $\mathbb{C}\mathbb{H}^2$  by holomorphic isometries. One can also think of  $f$  as a section of a  $\mathbb{C}\mathbb{H}^2$ -bundle over  $\Sigma$  or, when  $\rho$  is a discrete embedding, as a minimal immersion  $f : \Sigma \rightarrow \mathbb{C}\mathbb{H}^2/\rho$  into the quotient manifold.

We classify these pairs  $(\rho, f)$  up to  $PU(2, 1)$ -equivalence, i.e., up to the natural left action of  $PU(2, 1)$  by conjugation of  $\rho$  and the simultaneous ambient isometry of  $f$ . In particular, the space of such pairs has a natural “forgetful” map to the moduli space

$$\mathcal{R}(G) = \text{Hom}^+(\pi_1\Sigma, G)/G,$$

of conjugacy classes of reductive representations into  $G = PU(2, 1)$ . Recall that this is a real analytic variety whose connected components are indexed by the Toledo invariant  $\tau(\rho) \in \frac{2}{3}\mathbb{Z}$ ,  $|\tau(\rho)| \leq -\chi(\Sigma)$  (see, e.g., [19]). We show that there are families of pairs for every value of  $\tau$ .

We achieve this classification by exploiting the powerful machinery which links equivariant harmonic maps to the Yang-Mills-Higgs equations over a compact Riemann surface and thereby to Higgs bundles [10, 29, 9, 43]. Our starting point is two

---

\*Received May 8, 2017; accepted for publication July 27, 2017.

<sup>†</sup>Department of Mathematics and Computer Science, Rutgers-Newark, Newark, NJ 07102, USA (loftin@rutgers.edu).

<sup>‡</sup>Department of Mathematics, University of York, York YO10 5DD, UK (ian.mcintosh@york.ac.uk).

facts: (i) to each irreducible  $\rho$  and marked conformal structure on  $\Sigma$  (i.e., conjugacy class of Fuchsian representations) there is a  $\rho$ -equivariant harmonic map  $f$  [10, 9]; (ii) when this information is encoded into a  $G$ -Higgs bundle  $(E, \Phi)$   $f$  is weakly conformal precisely when  $\text{tr } \Phi^2 = 0$ . It is surprising that this approach has not, to our knowledge, been exploited before, since it provides a very effective way to understand the moduli as holomorphic data and avoids a direct analysis of the Gauss-Codazzi equations for minimal surfaces (cf. [44]).

When  $G = PU(2, 1)$  the structure of the Higgs bundles and their moduli space is quite well understood [52, 22, 3]. In this case the bundle  $E$  splits into a sum  $V \oplus L$  of a rank two sub-bundle  $V$  and a rank one sub-bundle  $L$  which are mapped to each other by the Higgs field, i.e., we can write  $\Phi = (\Phi_1, \Phi_2)$  where  $\Phi_1 : L \rightarrow KV$  and  $\Phi_2 : V \rightarrow KL$  for  $K$  the canonical bundle determined by the marked conformal structure. In fact by projective equivalence we may assume that  $L = 1$ , the trivial bundle. The Higgs field  $\Phi$  corresponds to the differential of  $f$ . To be precise, it corresponds to

$$\partial f : T^{1,0}\mathcal{D} \rightarrow T^{\mathbb{C}}\mathbb{C}\mathbb{H}^2,$$

and the components  $\Phi_1, \Phi_2$  correspond to the components of  $\partial f$  with respect to the type decomposition of  $T^{\mathbb{C}}\mathbb{C}\mathbb{H}^2$ . For a minimal surface there are two possibilities: (i)  $f$  is holomorphic ( $\Phi_2 = 0$ ) or anti-holomorphic ( $\Phi_1 = 0$ ), or (ii)  $f$  is neither and has isolated complex and anti-complex points which give finite divisors  $D_2$  and  $D_1$  over  $\Sigma$  (where, respectively,  $\Phi_2$  and  $\Phi_1$  have zeroes). We treat these two possibilities separately, and in fact it is the latter which we treat first, in §2 and §3. An important role is played by a cubic holomorphic differential  $\mathcal{Q}$  which can be naturally assigned to any minimal surface in a Kähler manifold of constant holomorphic section curvature [51]. It vanishes identically for holomorphic or anti-holomorphic immersions (but not only for them). When  $f$  is neither holomorphic nor anti-holomorphic we show that, with the two bilinear forms  $\gamma_j = \frac{1}{2} \text{tr } \Phi_j \Phi_j^\dagger$ , which carry the information of the metric  $\gamma$  induced by  $f$ , the data  $(\gamma_1, \gamma_2, \mathcal{Q})$  completely determines the minimal immersion up to ambient isometries. The principal results of §2 and §3 (Theorems 2.3 and 3.1) can be summarised as follows.

**THEOREM 1.1.** *Let  $\rho$  be irreducible and  $f$  be minimal and  $\rho$ -equivariant. If  $f$  is neither holomorphic nor anti-holomorphic then the pair  $(\rho, f)$  is faithfully determined, up to  $G$ -equivalence, by data  $(\Sigma_c, D_1, D_2, \xi)$  where:  $c$  is a marked conformal structure on  $\Sigma$ ,  $D_1$  and  $D_2$  are effective divisors on  $\Sigma$  whose degrees  $d_1, d_2$  satisfy*

$$2d_1 + d_2 < 6(g - 1) \text{ and } d_1 + 2d_2 < 6(g - 1),$$

and  $\xi \in H^1(\Sigma_c, K^{-2}(D_1 + D_2))$  represents an extension class which determines  $V$  as an extension of  $K^{-1}(D_1)$  by  $K(-D_2)$ . The Higgs bundle is then  $E = V \oplus 1$  equipped with a Higgs field determined by the extension class. Under the Dolbeault isomorphism this extension class corresponds to the cohomology class of  $-\bar{\mathcal{Q}}/\gamma_1\gamma_2$  in  $H^{0,1}(\Sigma_c, K^{-2}(D_1 + D_2))$ , and  $\xi = 0$  if and only if  $\mathcal{Q} = 0$ . In this correspondence,  $\rho$  has Toledo invariant  $\tau(\rho) = \frac{2}{3}(d_2 - d_1)$ , and the Euler number of the normal bundle of  $f$  is  $\chi(T\Sigma^\perp) = 2(g - 1) - d_1 - d_2$ .

In particular, the integers  $d_1, d_2$  determine, and are determined by, the Toledo invariant of  $\rho$  and the Euler number of the normal bundle of  $f$ .

The correspondence is constructive in the sense that, given data  $(\Sigma_c, D_1, D_2, \xi)$  satisfying the conditions above, we give an explicit construction of a stable Higgs

bundle  $(E, \Phi)$  with  $\text{tr } \Phi^2 = 0$ . The divisors  $D_1, D_2$  are precisely the divisors of anti-complex and complex points of the minimal immersion. For  $f$  to be strictly an immersion they must have no common points, but the construction still works to produce branched minimal immersions if they intersect, with branch points at the intersections.

This construction includes and greatly extends the construction of minimal Lagrangian embeddings we gave in [37]. Indeed, up to this time we were not aware of any other examples of non-holomorphic equivariant minimal immersions into  $\mathbb{C}\mathbb{H}^2$  (save for the Lagrangian examples whose existence is a consequence of the “mountain-pass” solutions to the Gauss equation described in [31]). The theorem above not only gives all examples for reductive representations (when we allow branch points) but allows us to describe the structure of the moduli space of these as a complex manifold (Theorem 1.3 below).

By a theorem of Wolfson [50],  $f$  is Lagrangian precisely when it has no complex or anti-complex points, and is therefore parametrised by the pair  $(\Sigma_c, \xi)$ . The embeddings constructed in [37] all have the property that the exponential map on the normal bundle provides a diffeomorphism between  $T\mathcal{D}^\perp$  and  $\mathbb{C}\mathbb{H}^2$ . It follows that  $\rho$  has a finite fundamental domain (given by the normal bundle over a finite fundamental domain for the action of  $\pi_1 \Sigma$  on  $\mathcal{D}$ ). Here we call such embeddings *almost  $\mathbb{R}$ -Fuchsian*, because they are deformations of the embedding  $\mathbb{R}\mathbb{H}^2 \rightarrow \mathbb{C}\mathbb{H}^2$ , which is equivariant with respect to every Fuchsian representation into  $SO(2, 1) \subset PU(2, 1)$ . In §4 we improve on the results in [37] by showing that whenever  $f$  is minimal Lagrangian with  $\|\mathcal{Q}\|_\gamma^2 < 2$  it is almost  $\mathbb{R}$ -Fuchsian and the unique  $\rho$ -equivariant minimal immersion. The  $\mathbb{R}$ -Fuchsian case corresponds to  $\mathcal{Q} = 0$ , and therefore  $\xi = 0$  by the theorem above. The uniqueness of  $f$  proved here implies that the data  $(\Sigma_c, \xi)$  also parametrises the almost  $\mathbb{R}$ -Fuchsian family, although at present we do not understand the appropriate bound on  $\xi$ . It is preferable to have the parametrisation in terms of  $(\Sigma_c, \xi)$  since that gets us directly to the Higgs bundle and therefore to  $\rho$ . The parametrisation in [37] using  $\mathcal{Q}$  requires an additional condition to provide a unique solution to the Gauss equation of the immersion. We describe the subtleties of existence and uniqueness for this equation in §4.1, which also draws on earlier work by Huang, Loftin & Lucia [31].

The minimal Lagrangian case suggests that it is important to understand those minimal immersions for which  $\mathcal{Q} = 0$ . These are treated in §5. We show that they have a very interesting interpretation in terms of the Higgs bundle, for  $\mathcal{Q} = 0$  exactly when the Higgs bundle is a *Hodge bundle* (or *variation of Hodge structure*). These are known to be the critical points of the Morse-Bott function  $\|\Phi\|_{L^2}^2$  [22] and come in two flavours: length two or length three. The length-two Hodge bundles all correspond to holomorphic or anti-holomorphic maps. The following theorem summarises our results regarding these.

**THEOREM 1.2.** *Let  $\rho$  be irreducible and  $f$  be branched holomorphic and  $\rho$ -equivariant. Then the pair  $(\rho, f)$  is faithfully determined by data  $(\Sigma_c, B, L, \mathbb{C}.\eta)$  where  $B$  is an effective divisor of degree  $b$ ,  $L$  is a holomorphic line bundle of degree  $l$  satisfying*

$$3(g-1) + \frac{1}{2}b < l < 6(g-1) - b, \quad 0 \leq b < 2(g-1),$$

*and  $\mathbb{C}.\eta \in \mathbb{P}H^1(\Sigma_c, KL^{-1}(B))$  determines the isomorphism class of  $V$  as an extension of  $K^{-1}(B)$  by  $K^{-2}L$ . The Toledo invariant of  $\rho$  is  $\frac{2}{3}(6g - 6 - b - l) > 0$ .*

This also accounts for anti-holomorphic immersions, since  $f$  is anti-holomorphic

and  $\rho$ -equivariant if and only if  $\bar{f}$  is holomorphic and  $\bar{\rho}$ -equivariant. The latter has the dual Higgs bundle to  $\rho$ .

As with the non-holomorphic case, the extension class  $\eta$  corresponds to the Dolbeault cohomology class of a tensor over  $\Sigma_c$  which has geometrical significance and is related to the second fundamental form of  $f$  (see Theorem 5.2). We explain how the limiting value  $\eta = 0$  corresponds to reducible representations which are not *maximal*, i.e., do not have  $\tau = \pm\chi(\Sigma)$ .

By contrast, the length-three Hodge bundles correspond to those pairs  $(\rho, f)$  coming from Theorem 1.1 with  $\xi = 0$ . By using the method of harmonic sequences [4, 7, 14, 15] we show that  $\xi = 0$  precisely when the harmonic sequence of  $f$  contains a holomorphic  $\rho$ -equivariant (and “timelike”) map into complex de Sitter 2-space. This is a pseudo-Hermitian symmetric space, the analogue of the real 2-dimensional de Sitter space which complements  $\mathbb{RH}^2$ .

In the final section, §6, we describe the moduli space

$$\mathcal{V} = \{(\rho, f) : \rho \text{ irreducible, } f \text{ branched minimal}\}/G.$$

By the results stated above it is a union of components  $\mathcal{V}(d_1, d_2)$  containing those pairs described by Theorem 1.1 and  $\mathcal{W}^+(b, l)$  (respectively,  $\mathcal{W}^-(b, l)$ ) containing the holomorphic (resp., anti-holomorphic) immersions described in Theorem 1.2. Of these we prove:

**THEOREM 1.3.** *Each  $\mathcal{V}(d_1, d_2)$  is a complex manifold of dimension  $8g - 8$ , while each  $\mathcal{W}^\pm(b, l)$  is a complex manifold of dimension  $3(g - 1) + l$ . All of these are diffeomorphic to bundles over the Teichmüller space of  $\Sigma$ , and the fibres are complex analytic submanifolds. For  $\mathcal{V}(d_1, d_2)$  each fibre  $\mathcal{V}_c(d_1, d_2)$  is a rank  $5g - 5 - d_1 - d_2$  vector bundle over  $S^{d_1}\Sigma_c \times S^{d_2}\Sigma_c$ , while for  $\mathcal{W}^\pm(b, l)$  each fibre is a  $\mathbb{CP}^{l-b-g}$ -bundle over  $S^b\Sigma_c \times \text{Pic}_l(\Sigma_c)$ .*

We finish in §6.3 with a brief discussion of the map  $\mathcal{V} \rightarrow \mathcal{R}(G)$  given by forgetting the immersion. There is much yet to be understood about this map. For example, we do not know if this map is onto for non-maximal representations, or the dimension of its image on components of  $\mathcal{V}$ . Nevertheless, we can make some salient remarks about its restriction to any fibre over Teichmüller space. We point out that on the image of  $\mathcal{V}$  the  $L^2$ -norm of the Higgs fields equals the area of the minimal immersion. The critical points of  $\|\Phi\|_{L^2}^2$  are all accounted for by the Hodge bundles, and therefore lie in the image of  $\mathcal{V}$ . A comparison of the structure of  $\mathcal{V}_c(d_1, d_2)$  with what is known of the Morse index from [22], together with an area bound established in §3, suggests that the fibres of the vector bundle  $\mathcal{V}_c(d_1, d_2)$  map onto the downward Morse flow of  $\|\Phi\|_{L^2}^2$ .

For us, one of the outstanding challenges is to use this construction to study the *quasi-Fuchsian* representations, where “quasi-Fuchsian” is meant in the sense of Parker & Platis [41], i.e., a convex cocompact, totally loxodromic, discrete embedding. Recent work of Guichard & Wienhard [25, Thm 1.8] has shown that  $\rho$  is a convex cocompact embedding precisely when it is an *Anosov embedding*. Since the latter are totally loxodromic, the notions “quasi-Fuchsian”, “convex cocompact embedding” and “Anosov embedding” coincide for  $PU(2, 1)$  (and more generally, any semisimple Lie group of real rank one). One knows from [24] that quasi-Fuchsian representations comprise an open subset of the representation variety  $\mathcal{R}(G)$ . Examples for every even value of  $\tau(\rho)$  were constructed by Goldman, Kapovich & Leeb [20], while Parker & Platis [40] constructed a open family of quasi-Fuchsian representations

in the Toledo invariant zero component. The latter family are perturbations of the Fuchsian representations corresponding to the totally geodesic and Lagrangian embedding  $\mathbb{R}\mathbb{H}^2 \subset \mathbb{C}\mathbb{H}^2$ , so they must overlap with the almost  $\mathbb{R}$ -Fuchsian representations constructed in [37] (all of which are quasi-Fuchsian).

Beyond these examples, there is very little known; it is not even known whether there are any quasi-Fuchsian representations for non-integral values of the Toledo invariant. One compelling reason for looking to minimal immersions to provide more insight is the theorem of Goldman & Wentworth [21], that for convex cocompact representations the harmonic map energy functional on Teichmüller space is a proper function. It therefore has at least one critical point, and it is a well-known result of Sacks & Uhlenbeck [42] that each critical point corresponds to a weakly conformal harmonic (i.e., branched minimal) map. Since our construction includes branched minimal maps, it follows that the map  $\mathcal{V} \rightarrow \mathcal{R}(G)$  has all quasi-Fuchsian representations in its image.

**Acknowledgements.** The authors gratefully acknowledge support from U.S. National Science Foundation grants DMS 1107452, 1107263, 1107367 *RNMS: Geometric Structures and Representation Varieties* (the GEAR Network). The first author gratefully acknowledges support from a Simons Collaboration Grant for Mathematicians 210124. The first author is grateful to Olivier Guichard, Zeno Huang, and Marcello Lucia for useful conversations. The second author thanks Peter Gothen for his remarks regarding the Morse flow on the moduli space of Higgs bundles.

**2. Minimal surfaces and their Higgs bundles.** We begin by setting up the notation and standard constructions for the minimal surfaces and the Higgs bundles we will be working with.

**2.1. Equivariant minimal surfaces in  $\mathbb{C}\mathbb{H}^2$ .** Our model for  $\mathbb{C}\mathbb{H}^2$  will be the projective model, as follows. Let  $\mathbb{C}^{2,1}$  denote the vector space  $\mathbb{C}^3$  equipped with the (indefinite) Hermitian metric

$$\langle v, v \rangle = v_1 \bar{v}_1 + v_2 \bar{v}_2 - v_3 \bar{v}_3.$$

Let  $\mathbb{C}_-^{2,1} = \{v \in \mathbb{C}^{2,1} : \langle v, v \rangle < 0\}$ , so that  $\mathbb{C}\mathbb{H}^2 \simeq \mathbb{P}\mathbb{C}_-^{2,1}$ . Thus we consider  $\mathbb{C}\mathbb{H}^2$  as the orbit of the line  $[0, 0, 1] \in \mathbb{P}\mathbb{C}_-^{2,1}$  under the standard action of  $G = PU(2, 1)$ . Consequently  $\mathbb{C}\mathbb{H}^2 \simeq G/H$ , where  $H \simeq P(U(2) \times U(1))$  is a maximal compact subgroup of  $G$ . We equip  $\mathbb{C}\mathbb{H}^2$  with its Hermitian metric of constant holomorphic sectional curvature  $-4$ ; so that its sectional curvature has bounds  $-4 \leq \kappa \leq -1$ . We write the Hermitian metric on  $\mathbb{C}\mathbb{H}^2$  as  $h = g - i\omega$ , where  $\omega(X, Y) = g(JX, Y)$ , and recall that  $(\mathbb{C}\mathbb{H}^2, h)$  is a Kähler-Einstein manifold.

We will always think of a minimal immersion  $f : \mathcal{D} \rightarrow \mathbb{C}\mathbb{H}^2$  as a conformal harmonic immersion, so the induced metric  $\gamma = f^*g$  is conformally equivalent to the hyperbolic metric  $\mu$ , with  $\gamma = e^u \mu$  for a smooth function  $u : \Sigma \rightarrow \mathbb{R}$ . To say that  $f$  is  $\rho$ -equivariant means it intertwines  $\rho$  with a Fuchsian representation  $\pi_1 \Sigma \rightarrow \text{Isom}(\mathcal{D})$ . The conjugacy class of such a representation is equivalent to a choice of a *marked conformal structure* on  $\Sigma$ , i.e., a point  $c \in \mathcal{T}_g$  in the Teichmüller space of  $\Sigma$ . We will write  $\Sigma_c$  to denote the surface with this structure. From now on we will assume  $f$  is  $\rho$ -equivariant.

To understand the properties of such minimal immersions we need some notation for the type decomposition of the (complexified) differential  $df : T^{\mathbb{C}}\mathcal{D} \rightarrow T^{\mathbb{C}}\mathbb{C}\mathbb{H}^2$ .

Both the domain and the codomain are complex manifolds, so to distinguish between the type decompositions of their tangent vectors we will write

$$T^{\mathbb{C}}\mathcal{D} = T^{1,0}\mathcal{D} \oplus T^{0,1}\mathcal{D}, \quad T^{\mathbb{C}}\mathbb{C}\mathbb{H}^2 = T'\mathbb{C}\mathbb{H}^2 + T''\mathbb{C}\mathbb{H}^2.$$

The projections will be such that  $X = X^{1,0} + \overline{X^{1,0}}$  (or  $X' + \overline{X'}$  as appropriate) whenever  $X$  is real. Our primary model for  $T^{\mathbb{C}}\mathbb{C}\mathbb{H}^2$  will be the projective model: viz, at any point  $\ell \in \mathbb{P}\mathbb{C}^{2,1}_-$  we use the isomorphism

$$\begin{aligned} T'_\ell\mathbb{C}\mathbb{H}^2 \oplus T''_\ell\mathbb{C}\mathbb{H}^2 &\rightarrow \text{Hom}(\ell, \ell^\perp) \oplus \text{Hom}(\ell^\perp, \ell) \subset \text{End}(\mathbb{C}^{2,1}); \\ (Z, W) &\mapsto (\pi_\ell^\perp \circ Z, \pi_\ell \circ W), \end{aligned} \quad (2.1)$$

where  $\pi_\ell : \mathbb{C}^{2,1} \rightarrow \ell$  is the orthogonal projection and we think of  $Z, W$  as operations of differentiation on local sections of  $\mathbb{C}\mathbb{H}^2 \times \mathbb{C}^3$ . In particular, conjugation in  $T^{\mathbb{C}}\mathbb{C}\mathbb{H}^2$  corresponds to taking the Hermitian transpose in  $\text{End}(\mathbb{C}^{2,1})$ , i.e., whose fixed subspace is  $\mathfrak{u}(2, 1)$ . The isomorphism can be derived from the symmetric space model for  $\mathbb{C}\mathbb{H}^2$ , which we will occasionally need to use (cf. the related model for  $\mathbb{C}\mathbb{P}^n$  in, for example, [5]). For that model, let  $\mathfrak{g} = \mathfrak{su}(2, 1)$  and let  $\mathfrak{h} \subset \mathfrak{g}$  denote the Lie subalgebra for  $H$ . Then the symmetric space decomposition is  $\mathfrak{g} = \mathfrak{h} + \mathfrak{m}$  where  $\mathfrak{m} = \mathfrak{h}^\perp$  with respect to the Killing form, and  $T\mathbb{C}\mathbb{H}^2 \simeq [\mathfrak{m}] = G \times_H \mathfrak{m}$ , where the action of  $H$  on  $\mathfrak{m}$  is its right adjoint action. It is easy to check that the fibre of  $[\mathfrak{m}^{\mathbb{C}}]$  at  $\ell$  agrees with the codomain of (2.1), and that the metric corresponds to  $g(A, B) = \frac{1}{2} \text{tr}(AB)$  whenever  $A, B \in \mathfrak{m}$ .

By extending  $\ell$  to mean the tautological sub-bundle the Hermitian metric  $h$  on  $T'\mathbb{C}\mathbb{H}^2$  is then equivalent to the inner product

$$h(Z_1, Z_2) = \langle \pi_\ell^\perp Z_1 \sigma_0, \pi_\ell^\perp Z_2 \sigma_0 \rangle, \quad \sigma_0 \in \Gamma(\ell), \quad \langle \sigma_0, \sigma_0 \rangle = -1.$$

Then type decomposition induces an isometry  $T\mathbb{C}\mathbb{H}^2 \rightarrow T'\mathbb{C}\mathbb{H}^2$ . These type decompositions give four complex linear parts of  $df$ :

$$\partial f' : T^{1,0}\mathcal{D} \rightarrow T'\mathbb{C}\mathbb{H}^2, \quad \partial f'' : T^{1,0}\mathcal{D} \rightarrow T''\mathbb{C}\mathbb{H}^2, \quad (2.2)$$

$$\bar{\partial} f' : T^{0,1}\mathcal{D} \rightarrow T'\mathbb{C}\mathbb{H}^2, \quad \bar{\partial} f'' : T^{0,1}\mathcal{D} \rightarrow T''\mathbb{C}\mathbb{H}^2, \quad (2.3)$$

which are related by  $\bar{\partial} f'' = \overline{\partial f'}$  and  $\bar{\partial} f' = \overline{\partial f''}$  using simultaneously the conjugation in  $T^{\mathbb{C}}\mathcal{D}$  and  $T^{\mathbb{C}}\mathbb{C}\mathbb{H}^2$ . Since  $f$  is  $\rho$ -equivariant, so is  $df$ , i.e.,  $df d\delta = d\rho(\delta)df$  whenever  $\delta \in \pi_1\Sigma \subset \text{Isom}(\mathcal{D})$ . Therefore we can think of  $df$  as a section of the bundle  $T^{\mathbb{C}}\Sigma^* \otimes (f^{-1}T^{\mathbb{C}}\mathbb{C}\mathbb{H}^2/\rho)$  over  $\Sigma_c$ . In particular,

$$\partial f = \partial f' + \partial f'',$$

is a smooth section of the vector bundle  $K \otimes (f^{-1}T^{\mathbb{C}}\mathbb{C}\mathbb{H}^2/\rho)$  over  $\Sigma_c$ , where  $K$  is the canonical bundle of  $\Sigma_c$ .

One says that  $f$  has a *complex point* at  $p = f(z)$  when  $\partial f''$  vanishes at  $p$  (i.e., when  $df(T^{1,0}\mathcal{D}) \subset T'\mathbb{C}\mathbb{H}^2$ ), and an *anti-complex point* when  $\partial f'$  vanishes at  $p$ . The Levi-Civita connexion induces a holomorphic structure on  $f^{-1}T^{\mathbb{C}}\mathbb{C}\mathbb{H}^2/\rho$  which preserves type decomposition, and  $f$  is harmonic when  $\nabla_{\bar{Z}}^{\mathbb{C}\mathbb{H}^2} \partial f(Z) = 0$  for local holomorphic sections  $Z$  of  $T^{1,0}\mathcal{D}$ , i.e., when

$$\nabla_{\bar{Z}}^{\mathbb{C}\mathbb{H}^2} \partial f'(Z) = 0 \quad \text{and} \quad \nabla_{\bar{Z}}^{\mathbb{C}\mathbb{H}^2} \partial f''(Z) = 0.$$

Thus  $\partial f'$  and  $\partial f''$  are holomorphic sections of their respective bundles; so a harmonic immersion which is not holomorphic or anti-holomorphic must have isolated anti-complex and complex points. We will denote the divisors of zeroes of  $\partial f'$  and  $\partial f''$  on  $\Sigma$  by  $D_1$  and  $D_2$  respectively.

Through these identifications there is a sesqui-linear form  $h(df', df')$  on  $T^{\mathbb{C}}\Sigma$  which gives the induced metric as  $\gamma = \operatorname{Re} h(df', df') = g(df', df')$ . The map  $f$  is *weakly conformal* when

$$h(\partial f', \bar{\partial} f') = 0,$$

and conformal when additionally  $df'$  does not vanish. Therefore, for conformal maps the induced metric and the pull-back of the Kähler form are expressed, with respect to a local complex coordinate  $z$ , as

$$\gamma = f^*g = (u_1^2 + u_2^2)|dz|^2, \quad f^*\omega = \frac{i}{2}(u_1^2 - u_2^2)dz \wedge d\bar{z}. \quad (2.4)$$

where

$$u_1 = \|\partial f'(Z)\|, \quad u_2 = \|\partial f''(Z)\|, \quad Z = \partial/\partial z. \quad (2.5)$$

The functions  $u_1, u_2$  are locally real analytic and vanish precisely at the anti-complex and complex points, respectively. They correspond to Hermitian metrics

$$\gamma_1 = h(\partial f', \partial f') = u_1^2 dz d\bar{z}, \quad \gamma_2 = h(\partial f'', \partial f'') = u_2^2 dz d\bar{z} \quad (2.6)$$

on  $K^{-1}(D_1)$  and  $K^{-1}(D_2)$  respectively. Note that, to apply  $\gamma$  to all elements of  $T^{\mathbb{C}}\Sigma$  consistently, the meaning of “ $|dz|^2$ ” above is

$$|dz|^2 = \frac{1}{2}(dz d\bar{z} + d\bar{z} dz),$$

in terms of the local complex linear forms  $dz, d\bar{z}$ . For this reason we do not write  $\gamma = \gamma_1 + \gamma_2$ .

Because the forms  $\gamma, f^*\omega$  live on  $\Sigma$  we can use some of the arguments which apply to compact minimal surfaces in Kähler-Einstein manifolds [48, 50, 6] to relate numerical invariants of a minimal immersion.

**THEOREM 2.1.** *Let  $f : \mathcal{D} \rightarrow \mathbb{C}\mathbb{H}^2$  be a  $\rho$ -equivariant minimal immersion which is neither holomorphic nor anti-holomorphic. Let  $d_1, d_2$  be the degrees of the divisors  $D_1$  and  $D_2$  of anti-complex and complex points. Then*

$$c_1(\rho) = d_1 - d_2, \quad (2.7)$$

$$\chi(\Sigma) + \chi(T\Sigma^{\perp}) = -d_1 - d_2, \quad (2.8)$$

where  $c_1(\rho)$  is the first Chern class of the bundle  $f^{-1}T\mathbb{C}\mathbb{H}^2/\rho$  over  $\Sigma$  and  $T\Sigma^{\perp} \subset f^{-1}T\mathbb{C}\mathbb{H}^2/\rho$  is the normal sub-bundle.

Wolfson also showed that in the absence of complex or anti-complex points a minimal immersion into a Kähler-Einstein surface of negative scalar curvature must be Lagrangian [50, Thm 2.1]. His argument extends to  $\rho$ -equivariant maps, so that in the setting of the previous theorem  $f$  will be Lagrangian if and only if  $d_1 = 0 = d_2$ .

Now we recall that the Toledo invariant  $\tau(\rho)$  is the integer

$$\tau(\rho) = \frac{2}{\pi} \int_{\Sigma} f^*\omega. \quad (2.9)$$

This is the normalisation which fits with the metric of holomorphic sectional curvature  $-4$ . It is known that for any representation  $\rho$  into  $PU(2, 1)$ ,  $|\tau(\rho)| \leq -\chi(\Sigma)$  [11] and  $\tau(\rho) \in \frac{2}{3}\mathbb{Z}$  [20]. Since  $\mathbb{C}\mathbb{H}^2$  has Einstein constant  $-6$ , equation (2.7) tells us that

$$\tau(\rho) = -\frac{2}{3}c_1(\rho) = \frac{2}{3}(d_2 - d_1). \quad (2.10)$$

Finally, as well as the degenerate metrics  $\gamma_j$  above, there is a third important invariant of minimal equivariant immersions [51, Cor 2.7], the cubic holomorphic differential  $\mathcal{Q} \in H^0(\Sigma, K^3)$  defined by

$$\mathcal{Q}(Z, Z, Z) = h(\nabla_Z^{\text{CH}^2} \partial f'(Z), \bar{\partial} f'(\bar{Z})) = -h(\partial f'(Z), \nabla_{\bar{Z}}^{\text{CH}^2} \bar{\partial} f'(\bar{Z})), \quad (2.11)$$

for  $Z \in T^{1,0}\mathcal{D}$ . It follows at once from this that  $\mathcal{Q}$  vanishes identically for holomorphic or anti-holomorphic immersions (but not only for them, as we shall see). When  $f$  is neither holomorphic nor anti-holomorphic we will see later that the quantities  $\gamma_1, \gamma_2, \mathcal{Q}$  uniquely determine  $f$  up to ambient isometries.

**2.2.  $G$ -Higgs bundles and representations.** As well as fixing our notation for Higgs bundles, we also need to summarise their correspondence with (projectively) flat connexions, and hence representations and harmonic maps, since we will be making explicit use of this correspondence for most of this article. A good general introduction to Higgs bundles can be found in, for example, [16, 17], while details for their moduli spaces in the case  $G = PU(2, 1)$  can be found in [22, 52].

Suppose  $\Sigma$  has been given a fixed conformal structure. With the notation of the previous section, a  $G$ -Higgs bundle for  $G = PU(2, 1)$  is a projective equivalence class of  $U(2, 1)$ -Higgs bundles. The latter are pairs  $(E, \Phi)$  consisting of a holomorphic rank three vector bundle  $E$  over  $\Sigma$  equipped with a splitting  $E = V \oplus L$  into a rank two sub-bundle  $V$  and a line sub-bundle  $L$  (both holomorphic) together with a holomorphic section

$$\Phi \in H^0(K \otimes [\text{Hom}(L, V) \oplus \text{Hom}(V, L)]), \quad (2.12)$$

called the *Higgs field*. Projective equivalence identifies pairs  $(E, \Phi)$  and  $(E', \Phi')$  when there is a holomorphic line bundle  $\mathcal{L}$  for which  $E' = E \otimes \mathcal{L}$  and  $\Phi' = \Phi$ .

We will write  $\Phi = (\Phi_1, \Phi_2)$  to denote the two summands implied by the direct sum (2.12). It is also convenient to write the holomorphic structure on  $E$  as a  $\bar{\partial}$ -operator on smooth sections,  $\bar{\partial}_E : \mathcal{E}^0(E) \rightarrow \mathcal{E}^{0,1}(E)$ . A Higgs bundle is *stable* if for any proper (non-zero)  $\Phi$ -invariant sub-bundle  $W \subset E$  the slope condition

$$\frac{\deg(W)}{\text{rk}(W)} < \frac{1}{3} \deg(E), \quad (2.13)$$

is satisfied. It is *polystable* when it is either stable or the direct sum of stable proper Higgs sub-bundles all having the same slope (these latter type are called *strictly polystable*). These properties are independent of the choice of pair  $(E, \Phi)$  representing the projective equivalence class. Now we recall that each polystable  $U(2, 1)$  Higgs bundle admits a  $\mathbb{C}^{2,1}$  metric and compatible projectively flat connexion, and thereby produces a flat  $PU(2, 1)$ -bundle.

**THEOREM 2.2** (Prop 3.9, [3]). *Fix a Kähler metric on  $\Sigma_c$  in the conformal class  $c$ , with Kähler 2-form  $\omega_c$ , normalised so that  $\Sigma_c$  has area  $2\pi$ . For each polystable stable  $U(2, 1)$ -Higgs bundle  $(E, \Phi)$  of slope  $\mu$  there is a  $\mathbb{C}^{2,1}$  metric on  $E$  for which  $L = V^\perp$  and the corresponding Chern connexion  $\nabla_E$  and Hermitian adjoint  $\Phi^\dagger$  yield a projectively flat connexion  $\nabla = \nabla_E + \Phi - \Phi^\dagger$ , i.e.,*

$$R^\nabla = R^{\nabla_E} + [\Phi \wedge \Phi^\dagger] = -i\mu\omega_c I_E. \quad (2.14)$$

The connexion  $\nabla$  induces a flat connexion on the principal  $PU(2, 1)$ -bundle whose associated bundle is  $\mathbb{P}E$ . The holonomy of this flat connexion yields a reductive



representation  $\rho : \pi_1 \Sigma \rightarrow PU(2, 1)$  (determined up to conjugacy) which is irreducible precisely when  $(E, \Phi)$  is stable (hence strictly polystable Higgs bundles correspond to reducible reductive representations).

REMARK 2.1. This theorem can be thought of as an example of the Donaldson-Uhlenbeck-Yau correspondence between stable bundles over compact Kähler manifolds and Hermitian-Einstein connections (Garcia-Prada gives a good overview which fits our context in [16]). For  $G$ -Higgs bundles such a correspondence is due to Hitchin for  $G = SL(2, \mathbb{C})$  [29] and Simpson [43] when  $G$  is a complex reductive algebraic group.

Note that the usual statement of the previous theorem gives the existence of a Hermitian metric on  $E$ . This is equivalent, since one can simply swap the sign of the metric on  $L$ , and the condition (2.12) on the Higgs field ensures that the adjoint remains the same. This makes the  $\mathbb{C}^{2,1}$  metric negative definite on  $L$ , and therefore  $L$  determines a smooth section of the  $\mathbb{C}\mathbb{H}^2$  bundle  $\mathbb{P}E_-$  (where  $E_-$  denotes the bundle of negative length vectors in  $E$ ). Since  $\mathbb{P}E_- \simeq \mathcal{D} \times_{\rho} \mathbb{C}\mathbb{H}^2$  this section is equivalent to a  $\rho$ -equivariant map  $f : \mathcal{D} \rightarrow \mathbb{C}\mathbb{H}^2$  in such a way that  $\Phi = \partial f$ . Moreover,  $\nabla_E$  induces a metric connexion on  $f^{-1}T\mathbb{C}\mathbb{H}^2$  which agrees with the pull-back of the Levi-Civita connexion, so that the equation  $\bar{\partial}_E \Phi = 0$  is the harmonic map condition for  $f$ .

In the reverse direction, a representation  $\rho : \pi_1 \Sigma \rightarrow G$  determines the projective bundle  $\mathbb{P}E$  uniquely and therefore a class of projectively equivalent  $\mathbb{C}^{2,1}$  bundles, each with a projectively flat connexion. By Corlette's results [9], there is a  $\rho$ -equivariant harmonic map  $f : \mathcal{D} \rightarrow \mathbb{C}\mathbb{H}^2$  precisely when  $\rho$  is reductive, and this map is unique when  $\rho$  is irreducible. The map corresponds to a line sub-bundle  $L \subset E$  and therefore a splitting  $E = L^{\perp} \oplus L$ . The splitting determines a bundle automorphism  $\sigma \in \text{End}(E, E)$  for which  $\sigma|_{L^{\perp}} = 1$  and  $\sigma|_L = -1$ , and therefore a decomposition  $\nabla = \nabla_E + \Psi$ , where

$$\nabla_E = \frac{1}{2}(\nabla + \sigma\nabla\sigma), \quad \Psi = \frac{1}{2}(\nabla - \sigma\nabla\sigma).$$

The Higgs field is  $\Phi = \Psi^{1,0}$ . The harmonic map equation, when paired with the projective flatness of  $\nabla$ , asserts that  $\bar{\partial}_E \Psi^{1,0} = 0$ , and thus the Higgs field satisfies (2.12) when we take  $V = L^{\perp}$ .

Two such bundles,  $(E, \nabla)$  and  $(E', \nabla')$ , are projectively equivalent when there is a line bundle  $\mathcal{L}$  equipped with a connexion  $\nabla_{\mathcal{L}}$  for which  $E' \simeq E \otimes \mathcal{L}$  and  $\nabla' \simeq \nabla \otimes \nabla_{\mathcal{L}}$  (the induced connexion on the tensor product). In particular, by taking  $\mathcal{L} = L^{-1}$  equipped with the connexion obtained from the restriction of  $\nabla$  to  $L$ , we may assume without loss of generality that  $E = V \oplus 1$ , where  $1$  denotes the trivial bundle, and the restriction of  $\nabla$  to  $1$  is the canonical flat connexion. In that case the Toledo invariant of  $\rho$  is  $-\frac{2}{3} \deg(V)$ .

REMARK 2.2. The alternative normalisation, used by Xia [52], is to note that since  $\deg(E \otimes \mathcal{L}) = \deg(E) + 3 \deg(\mathcal{L})$  one can normalise by degree, i.e., insist that  $0 \leq \deg(E) < 3$ . In particular, the topological type of  $\mathbb{P}E$  is determined by  $\deg(E) \bmod 3$ , and the representation  $\rho$  only lifts to  $SU(2, 1)$  when there exists an  $\mathcal{L}$  for which  $E \otimes \mathcal{L} \simeq \mathcal{D} \times_{\hat{\rho}} \mathbb{C}^{2,1}$  for some representation  $\hat{\rho} : \pi_1 \Sigma \rightarrow U(2, 1)$ . This happens if and only if  $\deg(E) \equiv 0 \bmod 3$ , i.e., when  $\tau \in 2\mathbb{Z}$ .

REMARK 2.3. From the Higgs bundle perspective, the Toledo invariant is sometimes defined to be  $\frac{2}{3} \deg(VL^{-1})$  (see, for example, [52]). This differs by a sign from

our convention, since

$$\frac{2}{3} \deg(VL^{-1}) = \frac{2}{3} \deg \operatorname{Hom}(L, V) = \frac{2}{3} c_1(\rho) = -\tau(\rho). \quad (2.15)$$

**2.3. Minimal surfaces and their Higgs bundles.** We are now in a position to classify, in terms of Higgs bundle data, the minimal  $\rho$ -equivariant surfaces which are neither holomorphic nor anti-holomorphic, when  $\rho$  is irreducible.

**THEOREM 2.3.** *An irreducible representation  $\rho \in \operatorname{Hom}(\pi_1 \Sigma, G)$  admits a  $\rho$ -equivariant minimal immersion  $f : \mathcal{D} \rightarrow \mathbb{C}\mathbb{H}^2$  which is neither holomorphic nor anti-holomorphic if and only if it corresponds to a Higgs bundle  $(E, \Phi)$  for which  $E = V \oplus 1$  where  $V$  is a rank 2 holomorphic extension*

$$0 \rightarrow K^{-1}(D_1) \xrightarrow{\Phi_1} V \xrightarrow{\Phi_2} K(-D_2) \rightarrow 0. \quad (2.16)$$

Here  $D_1, D_2$  are effective divisors with no common points, whose degrees  $d_1, d_2$  satisfy the stability inequalities

$$2d_1 + d_2 < 6(g-1) \text{ and } d_1 + 2d_2 < 6(g-1), \quad (2.17)$$

where  $g$  is the genus of  $\Sigma$ . The Higgs field is given by  $\Phi = (\Phi_1, \Phi_2)$ , after making the canonical identifications

$$\begin{aligned} \operatorname{Hom}(K^{-1}(D_1), V) &\simeq K(-D_1) \otimes \operatorname{Hom}(1, V), \\ \operatorname{Hom}(V, K(-D_2)) &\simeq K(-D_2) \otimes \operatorname{Hom}(V, 1). \end{aligned} \quad (2.18)$$

These divisors are, respectively, the divisors of anti-complex and complex points of the minimal immersion  $f$ . The representation  $\rho$  has  $\tau(\rho) = \frac{2}{3}(d_2 - d_1)$ .

*Proof.* Given  $\rho$  we obtain a projectively flat  $\mathbb{C}^{2,1}$  bundle of the form  $E = V \oplus 1$ , and then  $f$  provides holomorphic sections

$$\begin{aligned} \Phi_1 &= \partial f' \in H^0(K \otimes \operatorname{Hom}(1, V)) \simeq H^0(\operatorname{Hom}(K^{-1}, V)), \\ \Phi_2 &= \partial f'' \in H^0(K \otimes \operatorname{Hom}(V, 1)) \simeq H^0(\operatorname{Hom}(V, K)), \end{aligned}$$

whose sum is  $\Phi = \partial f$ . Since  $f$  is conformal we have

$$0 = g(\partial f, \partial f) = \frac{1}{2} \operatorname{tr}(\Phi^2).$$

From this we can show that  $\Phi_2 \circ \Phi_1 = 0$ , by the following local frame argument. Neither of  $\Phi_1, \Phi_2$  are identically zero since  $f$  is not  $\pm$ -holomorphic. So, away from its zero locus, the image of  $\Phi_1$  is a rank one sub-bundle  $V_1 \subset V$ . We can locally frame  $E$  by sections  $\sigma_1, \sigma_2$  of  $V$  and  $\sigma_3$  of  $1$  such that  $\sigma_1$  generates  $V_1$ , and we may assume this is a  $U(2, 1)$  frame with respect to the metric on  $E$ . It follows that there are locally holomorphic sections  $a, b, c$  of  $K$  for which

$$\Phi_2(\sigma_1) = a\sigma_3, \quad \Phi_2(\sigma_2) = b\sigma_3, \quad \Phi_1(\sigma_3) = c\sigma_1.$$

Since  $f$  is not anti-holomorphic  $c \neq 0$ . Now

$$\begin{aligned} \operatorname{tr}(\Phi^2) &= \operatorname{tr}(\Phi_1 \circ \Phi_2 + \Phi_2 \circ \Phi_1) \\ &= \langle \Phi_1 \circ \Phi_2(\sigma_1), \sigma_1 \rangle + \langle \Phi_1 \circ \Phi_2(\sigma_2), \sigma_2 \rangle - \langle \Phi_2 \circ \Phi_1(\sigma_3), \sigma_3 \rangle \\ &= 2ac. \end{aligned}$$

Therefore  $\text{tr}(\Phi^2) = 0$  implies  $a = 0$ , i.e.,  $\Phi_2 \circ \Phi_1 = 0$ . Thus

$$K^{-1} \xrightarrow{\Phi_1} V \xrightarrow{\Phi_2} K,$$

has the image of  $\Phi_1$  in the kernel of  $\Phi_2$ . Now  $\Phi_1$  vanishes precisely on anti-complex points, while  $\Phi_2$  vanishes precisely on complex points, so we have

$$0 \rightarrow K^{-1}(D_1) \xrightarrow{\Phi_1} V \xrightarrow{\Phi_2} K(-D_2) \rightarrow 0.$$

This must be exact at the middle since  $V$  has rank 2.

For stability we need to identify the  $\Phi$ -invariant sub-bundles. With respect to the local frame  $\sigma_1, \sigma_2, \sigma_3$  above  $\Phi$  is represented by the matrix

$$\begin{pmatrix} 0 & 0 & c \\ 0 & 0 & 0 \\ 0 & b & 0 \end{pmatrix}.$$

It follows that the only  $\Phi$ -invariant proper sub-bundles of  $E$  are the image  $V_1$  of  $\Phi_1$  and  $V_1 \oplus 1$ . So stability requires

$$\deg(V_1) < \frac{1}{3} \deg(E), \quad \frac{1}{2} \deg(V_1) < \frac{1}{3} \deg(E).$$

Since  $\deg(V_1) = d_1 - 2(g - 1)$  and  $\deg(E) = d_1 - d_2$  these inequalities are equivalent to the inequalities (2.17).

Reversing the argument is straightforward: when the Higgs bundle has this form we have  $\Phi_2 \circ \Phi_1 = 0$ , hence  $\text{tr}(\Phi^2) = 0$ . So when the Higgs bundle is stable the sub-bundle  $1 \subset E$  provides a conformal harmonic  $\rho$ -equivariant map  $f$  for some irreducible  $\rho$ .  $\square$

The proof works perfectly well in the case where  $D_1, D_2$  have common points, in which case the map  $f$  is a branched minimal immersion with branch points on  $D_1 \cap D_2$ . We will follow common usage and say  $D_1, D_2$  are *co-prime* when they are disjoint.

The Higgs bundle data which appears in the previous theorem can be written as a quadruple  $(\Sigma_c, D_1, D_2, \xi)$ , where  $\xi \in H^1(\Sigma_c, K^{-2}(D_1 + D_2))$  is the extension class of (2.16). The theorem shows that this data determines the pair  $(\rho, f)$  up to  $G$ -equivalence, i.e., up to the simultaneous action of  $G$  by conjugacy of  $\rho$  and ambient isometry of  $f$ . The exact sequence (2.16), which gives us  $V$  and  $\Phi_1, \Phi_2$ , is completely and uniquely determined by its extension class  $\xi$  [26, Ch. 5]. Moreover, since the isomorphisms (2.18) require  $D_1, D_2$ , not just their linear equivalence classes, the assignment from  $(\Sigma_c, D_1, D_2, \xi)$  to the  $G$ -equivalence class of  $(\rho, f)$  is bijective. The moduli space parametrised by this data will be described in §6.

REMARK 2.4. The proof above shows that when  $(E, \Phi)$  has  $\text{tr}(\Phi^2) = 0$  it cannot split into a direct sum of  $\Phi$ -invariant sub-bundles unless one of  $\Phi_1$  or  $\Phi_2$  is identically zero. It follows that a reducible  $\rho$  can only admit  $\rho$ -equivariant minimal surfaces which are holomorphic or anti-holomorphic. We describe all these reducible representations in Remark 5.1 below. However, there are also irreducible representations which admit holomorphic or anti-holomorphic surfaces: we give a complete classification in §5.

**3. The Higgs bundle data in terms of minimal surface data.** In this section we will give the explicit correspondence between the minimal surface data  $(\gamma_1, \gamma_2, \mathcal{Q})$  and the Higgs bundle data  $(\Sigma_c, D_1, D_2, \xi)$  of the previous section. This is achieved by exploiting the harmonic sequence for a minimal surface in  $\mathbb{C}\mathbb{H}^2$ . It provides us with a preferred system of local  $U(2, 1)$  frames for the bundle  $E$  in Theorem 2.3, and which we will call *Toda frames*. These frames are explicitly determined by the minimal surface data, and through them we calculate the extension class  $\xi$  of the bundle  $V$  in Theorem 2.3. The correspondence between the Higgs data and the minimal surface data then comes through the Dolbeault isomorphism

$$H^1(\Sigma_c, K^{-2}(D_1 + D_2)) \simeq H^{0,1}(\Sigma_c, K^{-2}(D_1 + D_2)).$$

**THEOREM 3.1.** *Let the pair  $(\rho, f)$  correspond to the Higgs data  $(\Sigma_c, D_1, D_2, \xi)$  as in the previous section. Let  $\gamma_1, \gamma_2, \mathcal{Q}$  be the minimal surface data determined by  $f$  through (2.6) and (2.11). Then the extension class  $\xi$  corresponds, under the Dolbeault isomorphism, to the cohomology class*

$$-\left[ \frac{\bar{\mathcal{Q}}}{\gamma_1 \gamma_2} \right] \in H^{0,1}(\Sigma_c, K^{-2}(D_1 + D_2)).$$

Moreover  $\xi = 0$  if and only if  $\mathcal{Q} = 0$ .

In particular, this means that  $(\gamma_1, \gamma_2, \mathcal{Q})$  determines  $(\Sigma_c, D_1, D_2, \xi)$ , since we get the conformal structure of the induced metric and the divisors  $D_1, D_2$  from  $\gamma_1, \gamma_2$ . Therefore Theorems 2.3 and 3.1 together have the following corollary.

**COROLLARY 3.2.** *If two  $\rho$ -equivariant minimal surfaces have the same data  $(\gamma_1, \gamma_2, \mathcal{Q})$  then they are identical up to ambient isometry.*

Before we begin the proof of Theorem 3.1 we describe the local Toda frames which link the minimal surface data to the local geometry of the Higgs bundle. Fix a pair  $(\rho, f)$  and let  $E = V \oplus 1$  be the Higgs bundle over  $\Sigma_c$  corresponding to them as above, equipped with its Higgs field  $\Phi$ , its  $\mathbb{C}^{2,1}$  metric and the Chern connexion  $\nabla_E$ .

**LEMMA 3.3.** *Let  $(U, z)$  be holomorphic chart on  $\Sigma_c$  for which  $U$  contains no complex or anti-complex points of  $f$ . Then over  $U$  there is a local trivialisation  $\varphi : E|_U \rightarrow U \times \mathbb{C}^3$  for which*

$$\varphi \circ \bar{\partial}_E \circ \varphi^{-1} = d\bar{z} \left[ \frac{\partial}{\partial \bar{z}} + \begin{pmatrix} -\bar{Z} \log u_1 & -\bar{Q}/u_1 u_2 & 0 \\ 0 & \bar{Z} \log u_2 & 0 \\ 0 & 0 & 0 \end{pmatrix} \right] \quad (3.1)$$

and

$$\varphi \circ \Phi \circ \varphi^{-1} = \begin{pmatrix} 0 & 0 & u_1 \\ 0 & 0 & 0 \\ 0 & u_2 & 0 \end{pmatrix} dz, \quad (3.2)$$

where  $\bar{Z} = \partial/\partial \bar{z}$  and  $u_1, u_2, Q$  are given by (2.5) and (2.11).

*Proof.* For notational convenience, let us set  $\ell_0 = 1 \subset E$ . Then  $V = \ell_0^\perp$  with respect to the  $\mathbb{C}^{2,1}$  metric, and  $\ell_0$  is the section of  $\mathbb{P}E_-$  which represents  $f$ . Thus

$\partial f', \bar{\partial} f' \in \Omega_{\Sigma}^1(\text{Hom}(\ell_0, \ell_0^\perp))$  and they determine line sub-bundles  $\ell_1, \ell_2 \subset \ell_0^\perp$  via their images, i.e.,

$$\ell_2 \otimes \bar{K} \xleftarrow{\bar{\partial} f'} \ell_0 \xrightarrow{\partial f'} \ell_1 \otimes K. \quad (3.3)$$

These two sub-bundles are orthogonal since  $f$  is conformal. We give each of these line bundles the holomorphic structure it inherits from  $\bar{\partial}_E$ , i.e., a local section  $\sigma_j$  of  $\ell_j$  is holomorphic when  $\pi_j \bar{\partial}_E \sigma_j = 0$ , where  $\pi_j : E \rightarrow \ell_j$  is the orthogonal projection. Note that since  $\Phi \in \text{Hom}(\ell_0, \ell_0^\perp)$  this holomorphic structure on each  $\ell_j$  agrees with the one induced by the projectively flat connexion  $\nabla = \nabla_E + \Phi + \Phi^\dagger$ . Then  $\partial f' = \Phi_1$  is a holomorphic map, while  $\bar{\partial} f' = \Phi_2^\dagger$  is anti-holomorphic, since  $f$  is harmonic. Since  $\nabla$  induces the canonical flat connexion on 1 we can choose a globally flat section  $f_0$  of 1, i.e.,  $\langle \nabla f_0, f_0 \rangle = 0$  with  $\langle f_0, f_0 \rangle = -1$ .

Now fix a chart  $(U, z)$  and for any section  $\sigma$  of  $E|U$  write  $X\sigma$  to mean  $\nabla_X \sigma$  with respect to the projectively flat connexion  $\nabla$ . Then the maps above can be written locally as

$$\ell_2 \xleftarrow{\pi_0^\perp \bar{Z}} \ell_0 \xrightarrow{\pi_0^\perp Z} \ell_1.$$

Define local sections  $\sigma_j \in \Gamma(U, \ell_j)$ ,  $j = 1, 2$  by

$$\sigma_1 = \partial f'(Z)f_0 = \pi_0^\perp Z f_0, \quad \sigma_2 = \bar{\partial} f'(\bar{Z})f_0 = \pi_0^\perp \bar{Z} f_0.$$

We assume that  $U$  does not contain any complex or anti-complex points, and therefore the functions  $u_1, u_2$  in (2.5) are non-vanishing. Clearly  $|\sigma_j| = u_j$ , and we set  $f_j = \sigma_j/u_j$ . Then  $f_1, f_2, f_0$  is a  $U(2, 1)$  frame for  $E$ . We claim that these satisfy the equations

$$Z f_1 = (Z \log u_1) f_1 + (Q/u_1 u_2) f_2, \quad (3.4)$$

$$Z f_2 = -(Z \log u_2) f_2 + u_2 f_0, \quad (3.5)$$

$$Z f_0 = u_1 f_1. \quad (3.6)$$

The last of these is obvious, since  $Z f_0 = \pi_0^\perp Z f_0$ .

Next consider

$$Z f_1 = \langle Z f_1, f_1 \rangle f_1 + \langle Z f_1, f_2 \rangle f_2.$$

Since  $f_0$  is holomorphic, so is  $\sigma_1$ , and therefore  $Z \langle \sigma_1, \sigma_1 \rangle = \langle Z \sigma_1, \sigma_1 \rangle$ . Thus

$$\begin{aligned} \langle Z f_1, f_1 \rangle &= u_1^{-1} \langle Z(u_1^{-1} \sigma_1), \sigma_1 \rangle \\ &= -Z \log(u_1) + u_1^{-2} Z \langle \sigma_1, \sigma_1 \rangle \\ &= Z \log(u_1). \end{aligned}$$

Now

$$\begin{aligned} \langle Z f_1, f_2 \rangle &= \langle Z(\sigma_1/u_1), \sigma_2/u_2 \rangle \\ &= \langle Z \sigma_1, \sigma_2 \rangle / u_1 u_2 \\ &= \langle \pi_0^\perp Z \pi_0^\perp Z f_0, \pi_0^\perp \bar{Z} f_0 \rangle / u_1 u_2. \end{aligned}$$

On the other hand, using (2.11) and the fact that  $\nabla^{\text{CH}^2}$  is the connexion on  $\text{Hom}(\ell_0, \ell_0^\perp)$  induced by the connexions on each bundle  $\ell_0, \ell_0^\perp$ , we have

$$\begin{aligned} Q &= \langle [\nabla_Z^{\text{CH}^2} \partial f'(Z)]f_0, \bar{\partial} f'(\bar{Z})f_0 \rangle \\ &= \langle \pi_0^\perp Z[\partial f'(Z)f_0] - \partial f'(Z)[\pi_0 Z f_0], \bar{\partial} f'(\bar{Z})f_0 \rangle \\ &= \langle \pi_0^\perp Z \pi_0^\perp Z f_0, \pi_0^\perp \bar{Z} f_0 \rangle - \langle \pi_0^\perp Z \pi_0 Z f_0, \pi_0^\perp \bar{Z} f_0 \rangle. \end{aligned} \quad (3.7)$$

The second term in the last line vanishes since  $\ell_{-1}$  is orthogonal to  $\ell_1$ . Thus  $\langle Z f_1, f_2 \rangle = Q/u_1 u_2$ .

Finally, consider

$$Z f_2 = \langle Z f_2, f_2 \rangle f_2 - \langle Z f_2, f_0 \rangle f_0.$$

For the first term we note that  $\sigma_2$  is anti-holomorphic since  $f_0$  is, so

$$0 = \langle Z \sigma_2, f_2 \rangle = Z u_2 + u_2 \langle Z f_2, f_2 \rangle.$$

Since  $\langle f_2, f_0 \rangle = 0$  the second term yields

$$-\langle Z f_2, f_0 \rangle = \langle f_2, \bar{Z} f_0 \rangle = \langle f_2, \sigma_2 \rangle = u_2.$$

Now that we have established the equations for the frame  $f_1, f_2, f_0$ , we take  $\varphi$  to be the corresponding trivialisation. In this frame the equations (3.4)-(3.6) show us that

$$\varphi \circ \nabla^{1,0} \circ \varphi^{-1} = dz \left( \frac{\partial}{\partial z} + \begin{pmatrix} Z \log u_1 & 0 & u_1 \\ Q/u_1 u_2 & -Z \log u_2 & 0 \\ 0 & u_2 & 0 \end{pmatrix} \right) \quad (3.8)$$

Thus (3.1) and (3.2) follow.  $\square$

At complex or anti-complex points we require a slight adjustment of the frame above. We may choose the chart  $(U, z)$  so that  $U$  contains precisely one complex or anti-complex point, at  $z = 0$ . Thus one of  $\sigma_1, \sigma_2$  vanishes at  $z = 0$ . To treat all cases simultaneously, let  $p, q$  be the non-negative integers for which  $z^{-q}\sigma_1$  and  $\bar{z}^{-p}\sigma_2$  are respectively holomorphic and anti-holomorphic, and do not vanish at  $z = 0$ : at most one of  $p, q$  is non-zero. It follows that the functions  $u_1/|z|^q = |z^{-q}\sigma_1|$  and  $u_2/|z|^p = |\bar{z}^{-p}\sigma_2|$  do not vanish. By setting

$$\tilde{f}_1 = \frac{z^{-q}\sigma_1}{|z^{-q}\sigma_1|} = (|z|/z)^q f_1, \quad \tilde{f}_2 = (|z|/\bar{z})^p f_2 = (z/|z|)^p f_2,$$

we obtain a  $U(2)$  frame  $\tilde{f}_1, \tilde{f}_2$  for  $V$  throughout  $U$ , with corresponding trivialisation  $\tilde{\varphi}$ . This is easiest to work with by writing it as

$$\tilde{\varphi} = S(\varphi|V), \quad S = \begin{pmatrix} (z/|z|)^q & 0 \\ 0 & (|z|/\bar{z})^p \end{pmatrix}. \quad (3.9)$$

The next step towards Theorem 3.1 is to represent the extension class  $\xi$  of  $V$  using a 1-cocycle with values in  $K^{-2}(D_1 + D_2)$ . For this computation, we choose an atlas  $\mathcal{U} = \{(U_j, z_j)\}$  for  $\Sigma_c$  of contractible charts for which each complex or anti-complex point lies only in one chart, at  $z_j = 0$ , and for simplicity assume charts containing distinct complex or anti-complex points are disjoint.

LEMMA 3.4. *There is a holomorphic atlas  $\mathcal{U} = \{(U_j, z_j)\}$  for  $\Sigma_c$  in which  $\xi$  is represented by the 1-cocycle  $\{(\xi_{jk}, U_j, U_k)\} \in H^1(\mathcal{U}, K^{-2}(D_1 + D_2))$  for which*

$$\xi_{jk} = a_j z^{-(p_j+q_j)} dz_j^{-2} - a_k dz_k^{-2} \text{ on } U_j \cap U_k, \quad (3.10)$$

for smooth functions  $a_j$  on  $U_j$  satisfying

$$\partial a_j / \partial \bar{z}_j = -\frac{\bar{Q}_j z_j^{p_j+q_j}}{u_{1j}^2 u_{2j}^2},$$

(assuming  $U_k$  contains no complex or anti-complex points).

*Proof.* For simplicity, set  $w = z/|z|$ . Using the previous lemma and the transformation (3.9) we have, on  $V$  over a single chart  $(U, z)$ ,

$$\tilde{\varphi} \circ \bar{\partial}_E \circ \tilde{\varphi}^{-1} = d\bar{z} \left[ \frac{\partial}{\partial \bar{z}} + \begin{pmatrix} -\bar{Z} \log(u_1/|z|^q) & -\bar{Q} w^{p+q}/u_1 u_2 \\ 0 & \bar{Z} \log(u_2/|z|^p) \end{pmatrix} \right].$$

Now suppose  $U_j \cap U_k \neq \emptyset$ , and assume without loss of generality that  $U_k$  contains no complex or anti-complex points, so that  $p_k = 0 = q_k$  and  $\tilde{\varphi}_k = \varphi_k|_V$ . Then we have transition relations  $\tilde{\varphi}_j = c_{jk} \tilde{\varphi}_k$  where

$$c_{jk} = \begin{pmatrix} w_j^{q_j} \frac{dz_j/dz_k}{|dz_j/dz_k|} & 0 \\ 0 & w_j^{-p_j} \frac{|dz_j/dz_k|}{dz_j/dz_k} \end{pmatrix},$$

where we have used that fact that

$$\frac{d\bar{z}_j/d\bar{z}_k}{|d\bar{z}_j/d\bar{z}_k|} = \overline{\frac{dz_j/dz_k}{|dz_j/dz_k|}} = \frac{|dz_j/dz_k|}{dz_j/dz_k}.$$

It follows that, as a smooth bundle,  $V \simeq K^{-1}(D_1) \oplus K(-D_2)$ .

To elucidate its holomorphic structure we find local trivialisations  $\chi_j$  in which

$$\chi_j \circ \bar{\partial}_E \circ \chi_j^{-1} = d\bar{z}_j \frac{\partial}{\partial \bar{z}_j},$$

i.e., we seek local gauge transformations  $\chi_j = R_j \tilde{\varphi}_j$  for which

$$R_j [\tilde{\varphi} \circ \bar{\partial}_E \circ \tilde{\varphi}^{-1}] R_j^{-1} = d\bar{z}_j \frac{\partial}{\partial \bar{z}_j}.$$

A straightforward calculation shows that this is achieved by taking

$$R_j = \begin{pmatrix} 1 & a_j \\ 0 & 1 \end{pmatrix} \begin{pmatrix} (u_{1j}/|z_j|^{q_j})^{-1} & 0 \\ 0 & u_{2j}/|z_j|^{p_j} \end{pmatrix},$$

where

$$\frac{\partial a_j}{\partial \bar{z}_j} = -\frac{\bar{Q}_j z_j^{p_j+q_j}}{u_{1j}^2 u_{2j}^2}, \quad (3.11)$$

throughout  $U_j$ . Such a function  $a_j$  exists, by the  $\bar{\partial}$ -Poincaré Lemma, because the right hand side is smooth throughout  $U_j$  since  $Q_j$  vanishes to at least order  $p_j + q_j$  at  $z_j = 0$  by (2.11). Thus the transition between two such charts  $\chi_j = b_{jk} \chi_k$  is given by

$$b_{jk} = R_j c_{jk} R_k^{-1} = \begin{pmatrix} z_j^{q_j} dz_j/dz_k & \lambda_{jk} \\ 0 & z_j^{-p_j} dz_k/dz_j \end{pmatrix},$$

where

$$\lambda_{jk} = a_j z_j^{-p_j} \frac{dz_k}{dz_j} - a_k z_j^{q_j} \frac{dz_j}{dz_k},$$

and this is holomorphic on  $U_j \cap U_k$  (this follows easily from (3.11)). Now using the same convention as [26, p74] this determines the 1-cocycle with values in  $K^{-2}(D_1 + D_2)$  given by

$$\begin{aligned} \xi_{jk} &= \left( z_j^{q_j} \frac{dz_j}{dz_k} \right)^{-1} \lambda_{jk} dz_k^{-2} \\ &= a_j z_j^{-(p_j+q_j)} dz_j^{-2} - a_k dz_k^{-2}. \end{aligned}$$

□

*Proof of Theorem 3.1.* For notational simplicity, set  $\mathcal{L} = K^{-2}(D_1 + D_2)$ . In  $U_j$  the quantity

$$z_j^{-(p_j+q_j)} dz_j^{-2}$$

is a local holomorphic section of  $\mathcal{L}$ . Therefore we have local smooth sections

$$\eta_j = a_j z_j^{-(p_j+q_j)} dz_j^{-2} \in \Gamma(U_j, \mathcal{E}^0(\mathcal{L})),$$

which provide a 0-cochain  $\eta$  for the sheaf  $\mathcal{E}^0(\mathcal{L})$  of locally smooth sections of  $\mathcal{L}$ . By (3.10) the 1-cocycle  $\{(\xi_{jk}, U_j, U_k)\}$  is, as a smooth cocycle, the coboundary of  $\eta$ . Now recall that the Dolbeault isomorphism  $H^1(\mathcal{U}, \mathcal{L}) \rightarrow H^{0,1}(\mathcal{U}, \mathcal{L})$  is derived from the short exact sequence

$$0 \rightarrow \mathcal{O}(\mathcal{L}) \rightarrow \mathcal{E}^0(\mathcal{L}) \xrightarrow{\bar{\partial}} \mathcal{E}^{0,1}(\mathcal{L}) \xrightarrow{\bar{\partial}} 0,$$

by taking the cohomology class of  $\bar{\partial}\eta$ . But

$$\bar{\partial}\eta_j = -\frac{\bar{Q}_j}{u_{1j}^2 u_{2j}^2} dz_j^{-2} \bar{d}z_j.$$

So  $\bar{\partial}\eta$  is represented by the cohomology class of the smooth form

$$-\bar{Q}/\gamma_1\gamma_2 \in \Gamma(\Sigma_c, \mathcal{E}^{0,1}(\mathcal{L})).$$

Finally, we show that this vanishes when it is  $\bar{\partial}$ -exact, by showing that it is harmonic with respect to Hermitian metric  $B = \gamma_1\gamma_2$  on  $\mathcal{L}$ . With respect to the Hodge inner product on  $\mathcal{E}^{*,*}(\mathcal{L})$  determined by  $B$  on  $\mathcal{L}$  and the induced metric  $\gamma$  on  $\Sigma_c$ , the adjoint of  $\bar{\partial} : \mathcal{E}^0(\mathcal{L}) \rightarrow \mathcal{E}^{0,1}(\mathcal{L})$  is given by

$$\mathcal{E}^{0,1}(\mathcal{L}) \xrightarrow{\bar{\partial}^*} \mathcal{E}^0(\mathcal{L}); \quad \bar{\partial}^* = -\bar{\star}\bar{\partial}\bar{\star},$$

where  $\bar{\star}$  is the conjugate linear Hodge star operator for our choice of metrics (see, for example, [49, p168]). It therefore suffices for us to show that

$$\bar{\partial}\bar{\star}(\bar{Q}/\gamma_1\gamma_2) = 0.$$



Let  $(U, z)$  be a chart in which the divisor  $D_1 + D_2$  has at most one point, of degree  $d$  at  $z = 0$ . Then  $\tau = z^{-d}dz^{-2}$  is a local holomorphic trivialising section of  $\mathcal{L}$  over  $U$  and

$$\mathcal{E}^{0,1}(U, \mathcal{L}) \xrightarrow{\bar{\kappa}} \mathcal{E}^{1,0}(U, \mathcal{L}^*); \quad a\tau d\bar{z} \mapsto -i\bar{a}B(\cdot, \tau)dz,$$

for any locally smooth complex function  $a$  over  $U$ . Now in  $U$  we write

$$\bar{Q}/\gamma_1\gamma_2 = \frac{\bar{Q}z^d}{u_1^2u_2^2}\tau d\bar{z},$$

and therefore

$$\bar{\partial}\bar{\kappa}(\bar{Q}/\gamma_1\gamma_2) = -i\frac{\partial}{\partial\bar{z}}\left(\frac{Q\bar{z}^d\|\tau\|^2}{u_1^2u_2^2}\right)(z^d dz^2)d\bar{z} \wedge dz.$$

But

$$\|\tau\|^2 = B(\tau, \tau) = |z|^{-2d}u_1^2u_2^2,$$

and therefore  $\bar{\partial}\bar{\kappa}(\bar{Q}/\gamma_1\gamma_2) = 0$  throughout  $U$  since  $Q$  is holomorphic.  $\square$

We finish this section by giving a global expression which relates the curvatures of the immersion  $f$  to the norms of  $Q$  and  $Q/\gamma_1\gamma_2$  with respect to the induced metric. This in turn provides an area bound for such immersions.

First we note that a straightforward calculation using (3.8) shows that the projective flatness of the connexion  $\nabla$  is equivalent to the local equations

$$Z\bar{Z}\log u_1^2 = 2u_1^2 + |Q|^2/u_1^2u_2^2 - u_2^2, \quad (3.12)$$

$$Z\bar{Z}\log u_2^2 = 2u_2^2 + |Q|^2/u_1^2u_2^2 - u_1^2, \quad (3.13)$$

in a chart  $(U, z)$  containing no complex or anti-complex points. These are the appropriate version of the Toda equations for this geometry (cf. [4] for the  $\mathbb{C}\mathbb{P}^n$  version). They are also related to the two equations Wolfson derived for the *Kähler angle* in [50]. Recall that the Kähler angle is a continuous function  $\theta : \Sigma_c \rightarrow \mathbb{R}$  for which  $f^*\omega = \cos(\theta)v_\gamma$ , where  $v_\gamma$  is the area form for the induced metric. Wolfson showed that, except at complex or anti-complex points where  $\theta$  is not differentiable,

$$i\partial\bar{\partial}\log \tan^2(\theta/2) = f^*\text{Ric}, \quad (3.14)$$

$$i\partial\bar{\partial}\log \sin^2(\theta) = (\kappa_\gamma + \kappa^\perp)v_\gamma, \quad (3.15)$$

where  $\text{Ric}$  is the Ricci form and  $\kappa_\gamma, \kappa^\perp$  are the Gaussian and normal curvatures of the immersion respectively. In our case we have, from (2.4),  $\cos(\theta) = (u_1^2 - u_2^2)/(u_1^2 + u_2^2)$ . Therefore

$$\tan^2(\theta/2) = \frac{u_2^2}{u_1^2}, \quad \sin^2(\theta) = \frac{4u_1^2u_2^2}{(u_1^2 + u_2^2)^2},$$

and (3.14) is just the difference of (3.12) and (3.13) since  $f^*\text{Ric} = -6f^*\omega$ . Now

$$\begin{aligned} Z\bar{Z}\log \sin^2 \theta &= Z\bar{Z}u_1^2 + Z\bar{Z}u_2^2 - 2Z\bar{Z}\log(u_1^2 + u_2^2) \\ &= u_1^2 + \frac{2|Q|^2}{u_1^2u_2^2} + u_2^2 - \frac{2}{u_1^2 + u_2^2}[Z\bar{Z}\log(u_1^2 + u_2^2)](u_1^2 + u_2^2), \end{aligned}$$

using (3.12) and (3.13). The second term on the right contains the local expression for  $\kappa_\gamma$ , so substituting this equation into (3.15) reveals that

$$\kappa^\perp - \kappa_\gamma = 2 \left( 1 + \frac{2|Q|^2}{u_1^2 u_2^2 (u_1^2 + u_2^2)} \right).$$

The right hand side can be written in terms of the quantities

$$\|\mathcal{Q}\|_\gamma = \frac{2\sqrt{2}|Q|}{(u_1^2 + u_2^2)^{3/2}}, \quad \left\| \frac{\mathcal{Q}}{\gamma_1 \gamma_2} \right\|_\gamma = \frac{|Q| \sqrt{u_1^2 + u_2^2}}{\sqrt{2} u_1^2 u_2^2}.$$

It is easy to check that their product is smooth everywhere, so that we obtain the global identity

$$\kappa^\perp - \kappa_\gamma = 2 \left( 1 + \left\| \frac{\mathcal{Q}}{\gamma_1 \gamma_2} \right\|_\gamma \|\mathcal{Q}\|_\gamma \right). \quad (3.16)$$

By integration over  $\Sigma$ , and using (2.8), we arrive at the following conclusion.

**PROPOSITION 3.5.** *Let  $f$  be a  $\rho$ -equivariant minimal immersion which is neither holomorphic nor anti-holomorphic, with induced metric  $\gamma$ , cubic holomorphic differential  $\mathcal{Q}$ ,  $d_1$  anti-complex points and  $d_2$  complex points. Then*

$$(4(g-1) - d_1 - d_2)\pi \geq \text{Area}_\gamma(\Sigma) + \int_\Sigma \|\mathcal{Q}\|_\gamma^2 v_\gamma, \quad (3.17)$$

with equality if and only if either  $\mathcal{Q} = 0$  or when  $f$  is Lagrangian.

Note that the stability inequalities (2.17) confirm that the left hand side is positive. The last statement follows because if  $\mathcal{Q} \neq 0$  equality requires  $\|\mathcal{Q}/\gamma_1 \gamma_2\|_\gamma = \|\mathcal{Q}\|_\gamma$ , which in turn requires  $u_1 = u_2$ , whence  $\cos(\theta) = 0$ .

**REMARK 3.1.** The local equations (3.12), (3.13) are clearly invariant under any unimodular scaling  $Q \mapsto e^{i\alpha}Q$ . Globally this corresponds to the symmetry  $\mathcal{Q} \mapsto e^{i\alpha}\mathcal{Q}$ , and by Theorem 3.1 this corresponds in turn to  $\xi \mapsto e^{-i\alpha}\xi$ . In fact this is equivalent to the well-known symmetry of the equations (2.14)  $\Phi \mapsto e^{i\psi}\Phi$  (taking  $\alpha = 2\psi$ ) which Hitchin showed is a Hamiltonian circle action on the moduli space of  $SL(2, \mathbb{C})$ -Higgs bundles [29]. To see this equivalence it is enough to perform the following gauge transformation for  $\partial_E + e^{i\psi}\Phi$  using the local gauge (3.8):

$$\begin{aligned} & \text{Ad} \left( \begin{pmatrix} e^{-2i\psi} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & e^{-i\psi} \end{pmatrix} \cdot \left( \frac{\partial}{\partial z} + \begin{pmatrix} Z \log u_1 & 0 & e^{i\psi} u_1 \\ Q/u_1 u_2 & -Z \log u_2 & 0 \\ 0 & e^{i\psi} u_2 & 0 \end{pmatrix} \right) \right) \\ &= \frac{\partial}{\partial z} + \begin{pmatrix} Z \log u_1 & 0 & u_1 \\ e^{2i\psi} Q/u_1 u_2 & -Z \log u_2 & 0 \\ 0 & u_2 & 0 \end{pmatrix} \end{aligned} \quad (3.18)$$

Note in particular that, unlike the  $SL(2, \mathbb{C})$  case, the map  $(E, \Phi) \mapsto (E, -\Phi)$  fixes every  $PU(2, 1)$ -Higgs bundle since  $\partial_E + \Phi$  and  $\partial_E - \Phi$  are gauge equivalent (by the symmetric space involution).

**4. Minimal Lagrangian immersions.** By Wolfson's theorem [50, Thm 2.1] Theorem 2.3 yields minimal Lagrangian immersions when both divisors  $D_1, D_2$  are zero. Therefore Theorem 2.3 implies the following characterisation of all equivariant minimal Lagrangian immersions.

**COROLLARY 4.1.** *Given a closed oriented surface  $\Sigma$  of hyperbolic type, minimal Lagrangian immersions  $f : \mathcal{D} \rightarrow \mathbb{C}\mathbb{H}^2$  which are equivariant with respect to an irreducible representation of  $\pi_1\Sigma$  into  $PU(2, 1)$  are in one-to-one correspondence with the Higgs data  $(\Sigma_c, \xi)$ , where  $c$  is a point in Teichmüller space and  $\xi \in H^{0,1}(\Sigma_c, K^{-2})$ .*

It is not strictly necessary to say that  $\rho$  is irreducible in this statement: this is implied by a combination of Corlette's result that twisted harmonic maps correspond to reductive representations [9] and Remark 2.4.

For a Lagrangian immersion the Kähler angle satisfies  $\sin\theta = 1$ , so that (3.15) implies  $\kappa^\perp = -\kappa_\gamma$ . Moreover,  $\|\mathcal{Q}\|_\gamma^2 = 2\|A_f\|^2$ , where  $A_f$  is the shape operator of  $f$  [38, Lemma 2.8] and

$$\|A_f\|^2 = \sup\{\frac{1}{2}(\text{tr}_\gamma A_f(\nu)^2) : \nu \in T_p\mathcal{D}^\perp, |\nu| = 1\}.$$

Since  $\mathbb{C}\mathbb{H}^2$  has constant Lagrangian sectional curvature  $-1$ , (3.16) reduces to the Gauss (and Ricci) equation for minimal Lagrangian immersions:

$$-1 = \kappa_\gamma + 2\|A_f\|^2. \quad (4.1)$$

In [37] we wrote this as an equation for the conformal factor  $\gamma = e^u\mu$  of the induced metric with respect to the hyperbolic metric  $\mu$ :

$$\Delta_\mu u - 2\|\mathcal{Q}\|_\mu^2 e^{-2u} - 2e^u + 2 = 0. \quad (4.2)$$

We gave existence results for this in terms of pairs  $(\Sigma_c, \mathcal{Q})$ , and showed that there is a constant  $k$ , independent of  $c \in \mathcal{T}_g$ , for which  $\|\mathcal{Q}\|_\mu \leq k$  yields a minimal  $\rho$ -equivariant embedding for which the normal bundle exponential map  $\exp^\perp : T\mathcal{D}^\perp \rightarrow \mathbb{C}\mathbb{H}^2$  is a diffeomorphism. We then showed that  $\rho$  is quasi-Fuchsian, since the image under  $\exp^\perp$  of a fundamental domain for  $\pi_1\Sigma$  gives a globally finite fundamental domain for  $\rho$ . Taking  $\mathcal{Q} = 0$  gives the  $\mathbb{R}$ -Fuchsian representations, i.e., those which factor through the canonical inclusion  $SO(2, 1) \rightarrow PU(2, 1)$ . It is therefore convenient to adopt here the following terminology (similar terminology is used in the study of minimal surfaces in  $\mathbb{R}\mathbb{H}^3$ , where it was introduced in [34]).

**DEFINITION 4.2.** *A representation  $\rho \in \text{Hom}(\pi_1, PU(2, 1))$  will be called almost  $\mathbb{R}$ -Fuchsian if it admits a  $\rho$ -equivariant minimal Lagrangian embedding  $f : \mathcal{D} \rightarrow \mathbb{C}\mathbb{H}^2$  whose normal bundle exponential map  $\exp^\perp$  is a diffeomorphism. We will call  $f$  an almost  $\mathbb{R}$ -Fuchsian embedding.*

A necessary and sufficient condition for  $\exp^\perp$  to be an immersion is that  $\|\mathcal{Q}\|_\gamma^2 \leq 2$  [37, Thm 7.1]. The following theorem improves substantially on the existence results in [37] by showing that almost  $\mathbb{R}$ -Fuchsian immersions exist, and are the unique minimal immersion, up to this optimal bound on  $\mathcal{Q}$ : this is the direct analogue of Uhlenbeck's result [46, Thm 3.3] for almost Fuchsian minimal surfaces in  $\mathbb{R}\mathbb{H}^3$ .

**THEOREM 4.3.** *Let  $f$  be a  $\rho$ -equivariant minimal Lagrangian immersion for which  $\|\mathcal{Q}\|_\gamma^2 < 2$ . Then  $f$  is an almost  $\mathbb{R}$ -Fuchsian embedding (and  $\rho$  is almost  $\mathbb{R}$ -Fuchsian). Moreover,  $f$  is the unique  $\rho$ -equivariant minimal immersion.*

REMARK 4.1. This theorem should also be compared with the parametrisation of hyperbolic affine sphere immersions  $\mathcal{D} \rightarrow \mathbb{R}^3$  equivariant with respect to an irreducible representation into  $PSL(3, \mathbb{R})$ , or equally, to the parametrisation of all convex real projective structures on  $\Sigma$  [47, 36, 35]. The data is a pair  $(\Sigma_c, \mathcal{Q})$  where  $\mathcal{Q}$  is a cubic holomorphic differential. By a theorem of Choi & Goldman [8] this data parametrises an entire component, the Hitchin component, of the representation space  $\mathcal{R}(SL(3, \mathbb{R}))$ . Labourie [35] directly related the hyperbolic affine sphere data to the Higgs bundles identified by Hitchin in [30]. In that case the Hitchin component is parametrized by Higgs bundles over  $\Sigma_c$  with bundle  $K^{-1} \oplus 1 \oplus K$  and Higgs field

$$\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ \mathcal{Q} & 0 & 0 \end{pmatrix}.$$

Theorem 4.3 shows that Corollary 4.1 provides a similar parametrisation of the almost Fuchsian locus inside the  $\tau = 0$  component  $\mathcal{R}(PU(2, 1))$ . It is unlikely, however, to parametrise the entire component. In the analogous case of equivariant minimal surfaces in  $\mathbb{RH}^3$ , there are examples of quasi-Fuchsian hyperbolic 3-manifolds which admit more than one minimal surface [1, 32].

We will break the proof of Theorem 4.3 into two parts, starting with the proof that  $f$  is an embedding. For this we need to recall from [37, §7] an explicit expression for  $\exp^\perp : T\mathcal{D}^\perp \rightarrow \mathbb{CH}^2$  when  $f$  is minimal Lagrangian. In this case the frame  $f_1, f_2, f_0$  used in the proof of Theorem 3.1 has the simple form

$$f_1 = \frac{1}{s}(f_0)_z, \quad f_2 = \frac{1}{s}(f_0)_{\bar{z}}, \quad s = u_1 = u_2.$$

Let  $S_- \subset \mathbb{C}_-^{2,1}$  be the pseudo-sphere into which  $f_0$  maps, and  $\pi : S_- \rightarrow \mathbb{CH}^2$  be the projection. When  $f$  is Lagrangian  $T\mathcal{D}^\perp = Jf_*T\mathcal{D}$  is a Lagrangian 2-plane in  $T\mathbb{CH}^2$ . This implies that  $\exp^\perp(T_z\mathcal{D}^\perp)$  is a totally geodesic Lagrangian disc normal to  $f(\mathcal{D})$  at  $f(z)$ . This Lagrangian disc has the form

$$\{\pi[(ia - b)f_1(z) + (ia + b)f_2(z) + f_0(z)] : a^2 + b^2 < \frac{1}{2}\}.$$

Let  $\Delta \subset \mathbb{C}$  denote the open disc of radius  $1/2$ . Then we have an identification between  $T\mathcal{D}^\perp$  and  $\mathcal{D} \times \Delta$  for which  $\exp^\perp$  is represented by the map

$$\Theta : \mathcal{D} \times \Delta \rightarrow \mathbb{CH}^2; \quad \Theta(z, w) = \pi(-\bar{w}f_1(z) + wf_2(z) + f_0(z)).$$

The following result improves [37, Thm 8.1].

LEMMA 4.4. *When  $\|\mathcal{Q}\|_\gamma^2 < 2$  the pullback metric  $\Theta^*g$  is complete, and therefore  $\Theta$  is a proper map and hence a diffeomorphism. In that case  $f$  is an embedding and  $\rho$  is almost  $\mathbb{R}$ -Fuchsian.*

*Proof.* Fix a point  $p \in \mathcal{D}$ . We can normalise the frame  $f_1, f_2, f_0$  so that these are the standard basis vectors  $e_1, e_2, e_3$  at  $p$ , and choose a conformal normal coordinate  $z$  centred at  $p$  so that  $\gamma(p) = |dz|^2$  (i.e.,  $s(0) = 1/\sqrt{2}$ ) with  $s_z = 0 = s_{\bar{z}}$  at this point. We may also rotate  $z$  so that  $Q_0 = Q(p)$  is real and non-negative. Since  $\mathcal{Q} = Q_0 dz^3$  and  $\|dz\| = \sqrt{2}$  we have  $0 \leq Q_0 < \frac{1}{2}$ . With such choices, in [37, §7], we computed the differential of  $\Theta$  (in affine coordinates) at  $p$  to be given by

$$\begin{pmatrix} d\Theta_1 \\ d\bar{\Theta}_1 \\ d\Theta_2 \\ d\bar{\Theta}_2 \end{pmatrix} = \begin{pmatrix} l & k & 0 & -1 \\ \bar{k} & l & -1 & 0 \\ \bar{k} & l & 1 & 0 \\ l & k & 0 & 1 \end{pmatrix} \begin{pmatrix} dz \\ d\bar{z} \\ dw \\ d\bar{w} \end{pmatrix},$$

where

$$l = \frac{1}{\sqrt{2}}(1 + |w|^2), \quad k = -2Q_0w - \frac{1}{\sqrt{2}}\bar{w}^2.$$

Now set  $\phi = ldz + kd\bar{z}$  and notice that in affine coordinates  $\Theta(p) = (-\bar{w}, w)$ . Then at  $p$  we compute the pull-back of  $g$  to be

$$\begin{aligned} (\Theta^*g)_p &= \sum_{i,j=1}^2 \frac{1}{1 - \|\Theta\|^2} \left( \delta_{ij} + \frac{\bar{\Theta}_i \Theta_j}{1 - \|\Theta\|^2} \right) d\Theta_i d\bar{\Theta}_j \\ &= \frac{1}{1 - 2|w|^2} \left[ \left( 1 + \frac{(-w)(-\bar{w})}{1 - 2|w|^2} \right) (\phi - d\bar{w})(\bar{\phi} - dw) \right. \\ &\quad + \left( 0 + \frac{(-w)w}{1 - 2|w|^2} \right) (\phi - d\bar{w})(\bar{\phi} + d\bar{w}) \\ &\quad + \left( 0 + \frac{\bar{w}(-\bar{w})}{1 - 2|w|^2} \right) (\bar{\phi} + dw)(\bar{\phi} - dw) \\ &\quad \left. + \left( 1 + \frac{\bar{w}w}{1 - 2|w|^2} \right) (\bar{\phi} + dw)(\phi + d\bar{w}) \right] \\ &= \frac{1}{(1 - 2|w|^2)^2} ([2|\phi|^2 - (w\phi + \bar{w}\bar{\phi})^2] + [2|dw|^2 + (wd\bar{w} - \bar{w}dw)^2]). \end{aligned} \quad (4.3)$$

Now consider the two terms in this expression:

$$\theta_1 = \frac{1}{(1 - 2|w|^2)^2} (2|\phi|^2 - (w\phi + \bar{w}\bar{\phi})^2), \quad \theta_2 = \frac{1}{(1 - 2|w|^2)^2} (2|dw|^2 + (wd\bar{w} - \bar{w}dw)^2).$$

The term  $\theta_2$ , which is the induced metric on  $\Delta$ , is just the Klein model for the hyperbolic plane and reflects the fact that the fibres of  $\exp^\perp$  are totally geodesic. The first term  $\theta_1$  is the expression at  $p$  for the metric induced by the immersion  $\varphi_w : \mathcal{D} \rightarrow \mathbb{C}\mathbb{H}^2$ ,  $\varphi_w(z) = \Theta(z, w)$ . We can think of each vector  $(-\bar{w}, w)$  as determining a section of  $T\mathcal{D}^\perp$ , and  $\varphi_w$  is the image of this under  $\exp^\perp$ . We claim that there is a constant  $\varepsilon_1 > 0$  for which, for every  $w$ ,

$$\varphi_w^*g(X, X) \geq \varepsilon_1\gamma(X, X), \quad \forall X \in T_p\mathcal{D}.$$

It follows that  $\Theta^*g = \theta_1 + \theta_2$  is bounded below by  $\varepsilon_2\gamma + \theta_2$ . The latter is a product of complete metrics on  $\mathcal{D} \times \Delta$  and therefore  $\Theta^*g$  is also complete.

To prove the claim, write  $w = w_1 + iw_2$  and  $\phi = \phi_1 + i\phi_2$  for real and imaginary parts, and set  $r^2 = |w|^2 < 1/2$ . Then

$$\theta_1 = \frac{1}{(1 - 2r^2)^2} \begin{pmatrix} \phi_1 & \phi_2 \end{pmatrix} \begin{pmatrix} 2 - 4w_1^2 & 4w_1w_2 \\ 4w_1w_2 & 2 - 4w_2^2 \end{pmatrix} \begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix}. \quad (4.4)$$

The eigenvalues of the matrix are 2 and  $2 - 4r^2$ . Therefore, using the smaller eigenvalue  $2 - 4r^2$ ,

$$\theta_1 \geq \frac{2}{1 - 2r^2} |\phi|^2 \geq 2|\phi|^2.$$

It now suffices to show that

$$|\phi|^2 \geq \varepsilon_2\gamma = \varepsilon_2(dx^2 + dy^2),$$

for a constant  $\varepsilon_2 > 0$  independent of  $w$ , where  $z = x + iy$ . For this, write  $k = k_1 + ik_2$  so that

$$\begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix} = B \begin{pmatrix} dx \\ dy \end{pmatrix} \text{ for } B = \begin{pmatrix} l + k_1 & k_2 \\ k_2 & l - k_1 \end{pmatrix}. \quad (4.5)$$

The components of the metric  $|\phi|^2$  are the entries of  $B^t B = B^2$ , and the eigenvalues are the roots of

$$\lambda^2 - 2(l^2 + |k|^2)\lambda + (l^2 - |k|^2)^2 = 0,$$

and are therefore  $(l \pm |k|)^2$ . Thus for any unit vector  $X \in T_p \mathcal{D}$ ,

$$|\phi|^2(X, X) \geq (l - |k|)^2.$$

Now

$$l - |k| = \frac{l^2 - |k|^2}{l + |k|},$$

and  $l + |k|$  is clearly bounded above, so it suffices to show that  $l^2 - |k|^2$  is bounded below by a positive constant independent of  $w$ . We compute

$$\begin{aligned} l^2 - |k|^2 &= \frac{1}{2}(1 + r^2)^2 - (2Q_0 w + \frac{1}{\sqrt{2}}\bar{w}^2)(2Q_0 \bar{w} + \frac{1}{\sqrt{2}}w^2) \\ &= r^2(1 - 4Q_0^2) + \frac{1}{2} - Q_0(\sqrt{2}r)^3 \cos(3\alpha), \end{aligned} \quad (4.6)$$

where  $w = re^{i\alpha}$ . Now  $Q_0 < \frac{1}{2}$  and  $\sqrt{2}r < 1$ , so  $1 - 4Q_0^2 > 0$  and  $\frac{1}{2} - 2\sqrt{2}Q_0 r^3 \cos(3\alpha) > 0$ . With  $\mathcal{Q}$  fixed we get a uniform positive lower bound over  $r, \alpha$ . Thus the metric  $\Theta^*g$  is complete on  $\mathcal{D} \times \Delta$ . We conclude, as in [37], that  $\Theta$  is a proper map and a diffeomorphism, whence  $f$  is an embedding and  $\rho$  is almost  $\mathbb{R}$ -Fuchsian.  $\square$

To prove that  $f$  is unique we first need a result which can be given in greater generality than our current situation. Let  $(N, g)$  be a complete Riemannian manifold.

**PROPOSITION 4.5.** *Let  $f : M \rightarrow (N, g)$  be a compact embedded minimal submanifold for which  $\exp^\perp : TM^\perp \rightarrow N$  is a diffeomorphism. For a local section  $\eta$  of  $TM^\perp$  of unit length and a positive constant  $r$ , set  $\varphi_r = \exp^\perp(r\eta)$  and let  $v_r$  be the volume form for the metric  $\varphi_r^*g$  on an open subset of  $M$ . Suppose  $dv_r/dr > 0$  for all  $r$  and for every  $\eta$ . Then  $f$  is the unique minimal immersion of  $M$  transverse to the fibres of  $\exp^\perp$ .*

The proof is given in Appendix A.

Now we can complete the proof of Theorem 4.3. Since  $\rho$  is quasi-Fuchsian the quotient  $\mathbb{C}\mathbb{H}^2/\rho$  is a manifold, and by the previous lemma  $f : \Sigma \rightarrow \mathbb{C}\mathbb{H}^2/\rho$  is a minimal embedding such that  $\exp^\perp : T\Sigma^\perp \rightarrow \mathbb{C}\mathbb{H}^2/\rho$  is a diffeomorphism. Now if  $\varphi : \mathcal{D} \rightarrow \mathbb{C}\mathbb{H}^2$  is any  $\rho$ -equivariant immersion then it must be transverse to the fibres of  $\exp^\perp$ , since  $d\varphi \circ d\delta = d\rho(\delta) \circ d\varphi$  for every  $\delta \in \pi_1\Sigma$  and the action of  $\rho$  is transverse to the fibres. Therefore the uniqueness claim in Theorem 4.3 follows from the previous proposition and the following lemma.

**LEMMA 4.6.** *Under the assumptions of Theorem 4.3, if  $v_r$  is the area form for the metric  $\varphi_r^*g$  induced by any local immersion of the form  $\varphi_r = \exp^\perp(r\eta)$  for a local section  $\eta$  of  $TM^\perp$  of unit length and a positive constant  $r$ , then  $dv_r/dr > 0$ .*

*Proof.* Since  $\varphi_r(z) = \Theta(z, re^{i\alpha})$  for some fixed  $r$  and  $\alpha$ , the induced metric  $\varphi_r^*g$  at a point  $p \in \Sigma$  is given by  $\theta_1$  from the proof of Lemma 4.4. Using the expression (4.4) we can write

$$\varphi_r^*g = \frac{1}{(1-2r^2)^2} \begin{pmatrix} dx & dy \end{pmatrix} B^t \begin{pmatrix} 2-4w_1^2 & 4w_1w_2 \\ 4w_1w_2 & 2-4w_2^2 \end{pmatrix} B \begin{pmatrix} dx \\ dy \end{pmatrix},$$

where  $B$  is given by (4.5). The determinant of the matrix in the middle is  $2(2-4r^2)$ , so that  $v_r = a(r)dx \wedge dy$  where

$$\begin{aligned} a(r) &= \frac{1}{(1-2r^2)^2} 2\sqrt{1-2r^2}(l^2 - |k|^2) \\ &= \frac{1}{(1-2r^2)^{3/2}} \left( 1 + 2r^2(1-4Q_0^2) - 4\sqrt{2}Q_0r^3 \cos(3\alpha) \right), \end{aligned}$$

using (4.6). A calculation shows that

$$\frac{da}{dr} = \frac{4r}{(1-2r^2)^{3/2}} \left[ (1+r^2)(1-4Q_0^2) + \frac{3}{2}(1-2\sqrt{2}Q_0r \cos(3\alpha)) \right].$$

Now  $Q_0 < \frac{1}{2}$ ,  $r < 1/\sqrt{2}$  and  $\cos(3\alpha) \leq 1$ , so  $da/dr > 0$ .  $\square$

**4.1. Families of solutions to the Gauss equation.** Theorem 4.3 shows that the norm  $\|\mathcal{Q}\|_\gamma$  gives control over uniqueness of minimal Lagrangian immersions, but at present we have no clear way of relating it to the parametrisation by the extension class  $\xi$ . Moreover, since this norm depends upon the solution to (4.2) it is difficult to control *a priori*. On the other hand, the combined results of [37] and [31] show that a bound on  $\|\mathcal{Q}\|_\mu$  must be combined with a condition the solution of (4.2) to get existence and uniqueness. One knows that the zero solution  $u \equiv 0$  is the unique solution for  $\mathcal{Q} = 0$ , and that for  $\|\mathcal{Q}\|_\mu$  small and non-zero there are always solutions [37], but these are not unique [31]. The challenge is to understand how solutions behave as one moves along a ray  $t\mathcal{Q}_0$ ,  $t \geq 0$ , given a fixed cubic holomorphic differential  $\mathcal{Q}_0$ . To study solutions along such rays, Huang et. al [31] introduced the following terminology.

DEFINITION 4.7. *A solution  $u$  to (4.2) is  $\mathcal{F}$ -stable if the linearised operator*

$$\mathcal{L} = -\Delta_\mu + 2e^u - 4\|\mathcal{Q}\|_\mu^2 e^{-2u}, \quad (4.7)$$

*is positive.*

This condition ensures, by the Implicit Function Theorem in the appropriate Sobolov spaces, that there is locally a smooth curve  $u(t)$  of solutions to

$$\mathcal{H}(u, t) = \Delta_\mu u - 2\|t\mathcal{Q}_0\|_\mu^2 e^{-2u} - 2e^u + 2 = 0, \quad (4.8)$$

nearby any  $\mathcal{F}$ -stable solution  $u(t_0)$ . Our aim here is to show that, given  $\mathcal{Q}_0$ , the  $\mathcal{F}$ -stable solutions form a continuous curve terminated at one end by the zero solution and at the other end by the first solution which is not  $\mathcal{F}$ -stable. This, together with the results of [37, 31], gives the following summary of the behaviour of solutions to (4.2) as the cubic differential is scaled.

THEOREM 4.8. *Fix a non-zero cubic holomorphic differential  $\mathcal{Q}_0$  on  $\Sigma_c$ . Set*

$$T_0 = \sqrt{4/27}(\sup_\Sigma \|\mathcal{Q}_0\|_\mu)^{-1}.$$

*Then:*

- (i) there exists a  $T_2 > T_0$  such that (4.8) has no solutions for  $t \geq T_2$ ;
- (ii) there exists  $T_0 \leq T_1 < T_2$  such that for  $t < T_1$  there is a unique continuous family of  $\mathcal{F}$ -stable solutions to (4.8). All  $\mathcal{F}$ -stable solutions lie on this family and the limiting solution  $u(T_1)$  is not  $\mathcal{F}$ -stable;
- (iii) for  $t < T_0$  the  $\mathcal{F}$ -stable solutions yield almost Fuchsian embeddings;
- (iv) for  $0 < t < T_1$  there is at least one solution which is not  $\mathcal{F}$ -stable.

REMARK 4.2. This result is analogous to Uhlenbeck's description of the bifurcation in families of solutions to the Gauss equation for minimal surfaces in  $\mathbb{R}\mathbb{H}^3$  [46, Thm 4.4]. We note that for Uhlenbeck the right notion of stability was stability with respect to the area functional. In our case that gives no extra control, since all minimal Lagrangian surfaces in  $\mathbb{C}\mathbb{H}^2/\rho$  are stable by a theorem of Oh [39].

Part (i) comes from [37], while (iii) comes from [37] and Theorem 4.3 above. Part (iv) and the existence of a local family of unique  $\mathcal{F}$ -stable solutions for  $t < T_1$  come from [31]. Here we provide the rest of (ii) via the following lemma.

LEMMA 4.9. *Let  $\mathcal{Q}_0$  be a holomorphic cubic differential on  $\Sigma_c$ , and let  $\tau > 0$  be such that  $u(\tau)$  is an  $\mathcal{F}$ -stable solution. Let  $u(t)$  be the local family of  $\mathcal{F}$ -stable solutions to (4.8) through  $u(\tau)$ . Then  $\dot{u}(\tau) \leq 0$  on all of  $\Sigma_c$ .*

*Proof.* By differentiating (4.8) we find that  $\dot{u}$  satisfies

$$-\mathcal{L}_\tau \dot{u} = 4\tau \|\mathcal{Q}_0\|_\mu^2 e^{-2u}.$$

Elliptic regularity implies  $\dot{u}$  is  $C^\infty$ . Now define  $\dot{u}^+ = \max\{\dot{u}(\tau), 0\}$ .

Then  $\dot{u}^+$  is in the Sobolev space  $H(\Sigma)$  and  $d\dot{u}^+ = 0$  wherever  $\dot{u} \leq 0$  (see e.g. [18]). Recall we define

$$\langle -\Delta_\mu \dot{u}^+, \dot{u}^+ \rangle = \int_\Sigma \|d\dot{u}^+\|_\mu^2 dA_\mu$$

in this case. Let  $v = 4\tau \|\mathcal{Q}_0\|_\mu^2 e^{-2u}$ , and let  $\epsilon_j \searrow 0$  be regular values of  $\dot{u}$  (as guaranteed by Sard's Theorem). Thus we can integrate by parts, as each  $\{\dot{u} = \epsilon_j\} = \partial\{\dot{u} > \epsilon_j\}$  is a smooth 1-manifold. Let  $w_j = \max\{\dot{u} - \epsilon_j, 0\}$ . Now since  $\mathcal{L} > 0$ ,

$$\begin{aligned} 0 &\geq \langle -\mathcal{L}\dot{u}^+, \dot{u}^+ \rangle \\ &= \int_\Sigma [-\|d\dot{u}^+\|_\mu^2 + (-2e^u + 4\|\mathcal{Q}_0\|_\mu^2 e^{-2u})(\dot{u}^+)^2] dA_\mu \\ &= \int_{\{\dot{u} > 0\}} [-\|d\dot{u}\|_\mu^2 + (-2e^u + 4\|\mathcal{Q}_0\|_\mu^2 e^{-2u})\dot{u}^2] dA_\mu \\ &= \lim_{j \rightarrow \infty} \int_{\{\dot{u} > \epsilon_j\}} [-\|dw_j\|_\mu^2 + (-2e^u + 4\|\mathcal{Q}_0\|_\mu^2 e^{-2u})\dot{u}^2] dA_\mu \\ &= \lim_{j \rightarrow \infty} \int_{\{\dot{u} > \epsilon_j\}} [w_j \Delta_\mu w_j + (-2e^u + 4\|\mathcal{Q}_0\|_\mu^2 e^{-2u})\dot{u}^2] dA_\mu \\ &= \lim_{j \rightarrow \infty} \int_{\{\dot{u} > \epsilon_j\}} [(\dot{u} - \epsilon_j) \Delta_\mu \dot{u} + (-2e^u + 4\|\mathcal{Q}_0\|_\mu^2 e^{-2u})\dot{u}^2] dA_\mu \\ &= \int_{\{\dot{u} > 0\}} (-\mathcal{L}\dot{u})\dot{u} dA_\mu \\ &= \int_{\{\dot{u} > 0\}} v\dot{u} dA_\mu \geq 0. \end{aligned}$$



(The limits above are valid by the Dominated Convergence Theorem.)

Since  $v$  is positive almost everywhere, the last inequality is strict if  $\dot{u} > 0$  anywhere on  $\Sigma$ . Thus by contradiction  $\dot{u}(\tau) \leq 0$  everywhere on  $\Sigma$ .  $\square$

*Proof of Thm 4.8(ii).* Let  $u(\tau)$  be an  $\mathcal{F}$ -stable solution, with local family  $u(t)$  and let  $\mathcal{L}_t$  be the corresponding family of linearised operators (4.7). Now

$$\dot{\mathcal{L}} = 2\dot{u}e^u + 8t\|\mathcal{Q}\|_\mu^2 e^{-2u}(\dot{u} - t),$$

which is nonpositive for  $t > 0$  by the previous Lemma. Thus  $\mathcal{L}_\tau > 0$  implies  $\mathcal{L}_t > 0$  for all  $t \in [0, \tau]$  in the path of solutions.

The Maximum Principle shows that every solution to (4.8) is nonpositive. Thus for any  $t$  in an interval of the form  $(\tau_0, \tau]$  the proposition implies  $0 \geq u_t \geq u_\tau$ , and thus we have uniform  $L^\infty$  bounds on  $u_t$  for all  $t \in (\tau_0, \tau]$ . Then the  $L^p$  theory, applied to (4.8), and standard bootstrapping show that the limit

$$\lim_{t \rightarrow \tau_0^+} u_t$$

exists and is a solution  $u_{\tau_0}$  to the equation. This provides a closedness argument for the continuity method. On the other hand the  $\mathcal{L}_t > 0$  condition, verified in the previous paragraph, provides openness as well, and thus we can extend the solution space back down to  $t = 0$ .  $\square$

**5. Surfaces with zero cubic holomorphic differential.** Minimal (possibly branched) immersions for which  $\mathcal{Q} = 0$  have particularly important properties. They include all the holomorphic and anti-holomorphic immersions and, by Theorem 3.1, the extension class zero cases when  $f$  is not holomorphic or anti-holomorphic. In all such cases, the Higgs bundle  $E$  is a *Hodge bundle* (or *variation of Hodge structure*), i.e.,  $E = \oplus_{i=1}^m E_i$  for proper sub-bundles  $E_i$  for which  $\Phi : E_i \rightarrow E_{i+1} \otimes K$ , with  $E_{m+1} = 0$ . For  $PU(2,1)$  the length  $m$  of the Hodge bundle must be either two or three [22]. We will show below that the length-two Hodge bundles correspond to holomorphic or anti-holomorphic immersions, while immersions which arise from Theorem 3.1 with  $\xi = 0$  give length-three Hodge bundles.

Hodge bundles play the central role in calculating the Betti numbers of the smooth components of the moduli space  $\mathcal{H}(\Sigma_c, G)$  of polystable Higgs bundles, and therefore the Betti numbers of the representation space  $\mathcal{R}(G)$ . For on smooth components the *Hitchin function*  $\|\Phi\|_{L^2}^2 : \mathcal{H}(\Sigma_c, G) \rightarrow \mathbb{R}$ , is a perfect Morse-Bott function, whose critical points are the Hodge bundles. The length-two Hodge bundles are minima, while the length-three Hodge bundles have non-zero Morse index [22].

**5.1. Holomorphic and anti-holomorphic surfaces.** By Toledo's theorem [45] every *maximal representation* (those for which  $\tau(\rho) = \pm\chi(\Sigma)$ ) leaves invariant a complex line and acts on that line as a Fuchsian representation. Such representations are reducible. To understand the non-maximal  $\rho$ -equivariant holomorphic or anti-holomorphic immersions, we will start by describing their Hodge bundles. First we note that for any holomorphic  $\rho$ -equivariant immersion  $f : \mathcal{D} \rightarrow \mathbb{C}\mathbb{H}^2$  the area form  $v_\gamma$  for the induced metric equals  $f^*\omega$ . It follows from the definition (2.9) that  $\tau(\rho) > 0$ . For anti-holomorphic immersions  $f^*\omega = -v_\gamma$  so that  $\tau(\rho) < 0$ . The next lemma completely characterises the Higgs bundle data for representations which admit either holomorphic or anti-holomorphic branched immersions.

**THEOREM 5.1.** *An irreducible representation  $\rho$  admits a branched holomorphic  $\rho$ -equivariant immersion if and only if it corresponds to a Hodge bundle  $(V \oplus 1, \Phi)$*

with  $\Phi = (\Phi_1, 0)$  and  $V$  a non-trivial extension bundle of the form

$$0 \rightarrow K^{-1}(B) \xrightarrow{\Phi_1} V \rightarrow K^{-2}L \rightarrow 0. \quad (5.1)$$

Here  $B$  is an effective divisor of degree  $b \geq 0$  (the divisor of branch points of the immersion) and  $L$  is a line bundle of degree  $l$ , only determined up to isomorphism, satisfying the inequalities

$$3(g-1) + \frac{1}{2}b < l < 6(g-1) - b, \quad 0 \leq b < 2(g-1). \quad (5.2)$$

In particular,  $\tau(\rho) = \frac{2}{3}(6g-6-b-l)$  and  $0 < \tau(\rho) < 2(g-1) - b$ .

Moreover,  $(E, \Phi)$  corresponds to a branched anti-holomorphic immersion  $f$  if and only if  $\bar{f}$  is the branched holomorphic immersion determined by  $(E^*, \Phi^t)$ .

Note that by  $\bar{f}$  we mean the post-composition of  $f$  with the natural anti-holomorphic involution on  $\mathbb{C}\mathbb{H}^2$  which descends from complex conjugation on  $\mathbb{C}^{2,1}$ . Clearly  $f$  is  $\rho$ -equivariant precisely when  $\bar{f}$  is  $\bar{\rho}$ -equivariant. The map  $\rho \mapsto \bar{\rho}$  is an involution on  $\text{Hom}(\pi_\Sigma, G)/G$  for which  $\tau(\bar{\rho}) = -\tau(\rho)$ : it fixes the representations with values in  $SO(2, 1)$ .

*Proof.* First suppose  $(E, \Phi)$ , with  $E = V \oplus 1$  and  $\Phi = (\Phi_1, \Phi_2)$ , is a length-two Hodge bundle with  $\deg(V) < 0$ . In this case by [22, §3] we have  $\Phi_2 = 0$ . The corresponding representation  $\rho$  admits a  $\rho$ -equivariant harmonic map  $f : \mathcal{D} \rightarrow \mathbb{C}\mathbb{H}^2$  determined by the sub-bundle  $1$  as a section  $\mathbb{P}E_-$  with  $df' = \Phi$ . Therefore  $\bar{\partial}f' = 0$  and  $f$  is holomorphic. Conversely, suppose  $\rho$  admits a holomorphic immersion  $f$ . Taking  $V = f^{-1}T'\mathbb{C}\mathbb{H}^2$  and  $\Phi = \partial f' : 1 \rightarrow KV$  gives a length-two Hodge bundle with  $\deg(V) < 0$ . The involution  $(E, \Phi) \rightarrow (E^*, \Phi^t)$  on Higgs bundles maps length-two Hodge bundles with  $\Phi_2 = 0$  to those with  $\Phi_1 = 0$ . In the latter case  $V \simeq f^{-1}T''\mathbb{C}\mathbb{H}^2$  with the opposite complex structure, and the map  $f$  is anti-holomorphic.

Now the structure of  $(V \oplus 1, \Phi)$  follows a simplified version of the argument in the proof of Theorem 2.3. We can think of  $\Phi_1$  as a holomorphic section of  $KV$ , with divisor  $B \geq 0$  corresponding to the branch divisor of  $f$ . The bundle injection  $\Phi_1 : K^{-1}(B) \rightarrow V$  has image  $V_1$  and quotient line bundle  $V_2 = V/V_1$ . Provided  $V$  is not the direct sum  $V_1 \oplus V_2$  the  $\Phi$ -invariant sub-bundles of  $E$  are  $V_1, V_1 \oplus 1$  and  $V$ . The stability inequalities are therefore

$$\deg(V_1) < \frac{1}{3}\deg(V), \quad \frac{1}{2}\deg(V_1) < \frac{1}{3}\deg(V), \quad \frac{1}{2}\deg(V) < \frac{1}{3}\deg(V).$$

On the other hand, if  $V$  is the direct sum then  $V_2$  is also  $\Phi$ -invariant and stability requires the additional inequality

$$\deg(V_2) < \frac{1}{3}\deg(V), \text{ i.e., } \frac{2}{3}\deg(V) < \deg(V_1),$$

so this is not possible. For later convenience we write  $V_2 = K^{-2}L$  and the inequalities (5.2) follow from  $\deg(V_1) = b - 2(g-1)$  and  $\deg(V_2) = l - 4(g-1)$ .  $\square$

Note that while the splitting of  $V \simeq f^{-1}T'\mathbb{C}\mathbb{H}^2/\rho$  into  $T\Sigma \oplus T\Sigma^\perp$  is  $J$ -invariant, the sub-bundle  $T\Sigma^\perp$  is not  $\bar{\partial}_E$ -invariant unless  $\rho$  is reducible. Indeed, the normal bundle is  $\partial_E$ -invariant (since it is paired with  $T\Sigma$  by the Hermitian metric) so the induced structure of this splitting makes the normal bundle *anti-holomorphic*.

**REMARK 5.1 (Reducible representations).** Although  $E = V \oplus 1$  cannot be stable when  $V$  is a trivial extension, it can be polystable. This corresponds to a reducible reductive representation. Such representations either:

- (i) factor through a maximal compact subgroup, or,
- (ii) factor through  $P(U(1, 1) \times U(1))$ .

This is easy to see. We may simplify things, by replacing  $\rho$  by  $\bar{\rho}$  if necessary, to assume that  $\tau(\rho) \geq 0$  and thus  $\Phi_2 = 0$ . To be strictly polystable  $(E, \Phi)$  must decompose into either

$$(i) (V, 0) \oplus (1, 0), \text{ or } (ii) (1 \oplus V_1, \Phi_1) \oplus (V_2, 0),$$

where  $V = V_1 \oplus V_2$  and  $\Phi_1 : 1 \rightarrow V_1$ . In the first case  $V$  must be a stable rank two bundle of degree zero (to have the same slope as 1) and therefore the representation lies in a maximal compact subgroup and has  $\tau(\rho) = 0$ . We note that the corresponding harmonic map  $f$  is constant. In the second case  $(1 \oplus V_1, \Phi_1)$  corresponds to a representation into  $U(1, 1)$ . In this case polystability requires

$$\deg(V_1) < \frac{1}{2} \deg(V_1 \oplus 1) = \frac{1}{2} \deg(V_1), \text{ i.e., } \deg(V_1) < 0,$$

together with the ‘‘same slope’’ condition  $\frac{1}{2} \deg(V_1) = \deg(V_2)$ . When we write  $V_1 = K^{-1}(B)$  as above we deduce that

$$b < 2(g - 1), \quad \deg(V_2) = \frac{1}{2}b - g + 1.$$

Thus  $b$  is even and

$$\tau(\rho) = -\frac{2}{3} \deg(V) = -\deg(V_1) = 2g - 2 - b \in 2\mathbb{Z}.$$

In particular, such  $\rho$  can only admit an unbranched holomorphic map  $f$  when  $\tau(\rho)$  is maximal, i.e., when  $\rho$  factors through a Fuchsian representation. The map  $f : \mathcal{D} \rightarrow \mathbb{C}\mathbb{H}^2$  is a totally geodesic embedding onto a complex line. More generally, the  $PU(1, 1)$  representation corresponding to the rank two Higgs bundle  $(1 \oplus V_1, \Phi_1)$  has Toledo invariant  $-\deg(\text{Hom}(1, V_1))$ , which therefore equals  $\tau(\rho)$ . Every irreducible representation into  $PU(1, 1)$  of even Toledo invariant lifts to  $SU(1, 1)$ , and therefore provides a representation into  $P(U(1, 1) \times U(1))$ . Thus the whole structure of Higgs bundles for irreducible representations in  $SU(1, 1)$  [29] lifts up to provide reducible representations into  $PU(2, 1)$ , and this is what we are seeing above. Note that those which are non-maximal cannot be convex cocompact, since they preserve a complex line but act non-cocompactly on this line.

Let  $\eta \neq 0$  be the extension class of the extension (5.1) for  $\rho$  irreducible. Since  $L$  is only determined up to isomorphism the Higgs bundle only determines  $\eta$  up to scale. Therefore each Higgs bundle in Theorem 5.1 corresponds to a quadruple  $(\Sigma_c, B, L, \mathbb{C}\eta)$  where  $\mathbb{C}\eta$  is the line generated by  $\eta$ , i.e., a point in  $\mathbb{P}H^1(\Sigma_c, KL^{-1}(B))$ . In fact, it is not hard to show that the rescaling of  $\eta$  corresponds to the  $\mathbb{C}^*$ -action on the Higgs bundle  $(E, \Phi) \mapsto (E, t\Phi)$ . Since Hodge bundles are invariant under this action, this gives another way of interpreting the independence of the data on the scale of  $\eta$ . However, once  $(\rho, f)$  is known there is a preferred representative for  $\eta$  given by the geometric invariants of  $f$  via the Dolbeault isomorphism. First we need to introduce the tensor

$$S \in \mathcal{E}^0(\Sigma_c, K^2 \bar{K}^2), \quad S = h(\mathbb{I}^{2,0}, \mathbb{I}^{2,0}),$$

where  $\mathbb{I}^{2,0} = \pi^\perp \nabla' \partial f'$  is the  $(2, 0)$  component of the second fundamental form of  $f$  (here  $\pi^\perp : f^{-1}T'\mathbb{C}\mathbb{H}^2/\rho \rightarrow T\Sigma^\perp$  is projection onto the normal bundle). We will show that  $\mathbb{I}^{2,0}$  is a holomorphic section of  $K^2 \otimes T\Sigma^\perp$ .

**THEOREM 5.2.** *Let  $f : \mathcal{D} \rightarrow \mathbb{C}\mathbb{H}^2$  be a  $\rho$ -equivariant branched holomorphic immersion, with  $\rho$  irreducible and data  $(\Sigma_c, B, L, \mathbb{C}\eta)$ . Then  $L \simeq \mathcal{O}(D)$ , where  $D$  is the divisor of zeroes of  $\mathbb{I}^{2,0}$ , and we can choose  $\eta$  so that under the Dolbeault isomorphism it maps to the cohomology class*

$$-[S/\gamma] \in H^{0,1}(\Sigma_c, K(B - D)).$$

*Proof.* We follow the steps in the proof of Theorem 3.1 but using a local frame more suited to holomorphic maps. As before, use  $\ell_0 \subset E$  to denote 1 and write its  $\partial f'$  transform as  $\ell_1 \otimes K$ . But now take the further transform of  $\ell_1$ , so we have a harmonic sequence

$$\ell_0 \xrightarrow{\pi_0^\perp Z} \ell_1 \xrightarrow{\pi_1^\perp Z} \ell_2 \xrightarrow{\pi_2^\perp Z} 0,$$

in each chart  $(U, z)$ . The last step terminates the sequence because  $\pi_2^\perp Z : \ell_2 \rightarrow \ell_0$  is the adjoint to  $\pi_0^\perp \bar{Z} : \ell_0 \rightarrow \ell_2$ , which represents  $\bar{\partial} f' = 0$ . As before, let  $f_0 \in \Gamma(\ell_0)$  be global and parallel with  $\langle f_0, f_0 \rangle = -1$ . Set

$$\sigma_1 = \pi_0^\perp Z f_0 = Z f_0, \quad \sigma_2 = \pi_1^\perp Z \sigma_1 = \pi_1^\perp Z Z f_0.$$

Since  $f_0$  is a holomorphic section of  $\ell_0$ ,  $\sigma_j$  is a holomorphic section of  $\ell_j$  (by standard harmonic sequence theory [4]). Under the isomorphism  $f^{-1}T'\mathbb{C}\mathbb{H}^2/\rho \simeq \text{Hom}(1, V) \simeq V$  the image  $\partial f'(T^{1,0}\Sigma)$  of the holomorphic tangent bundle of  $\Sigma$  is identified with  $\ell_1$ , i.e.,  $\ell_1 \simeq K^{-1}(B)$ . Clearly the induced metric of  $f$  is  $u_1^2 |dz|^2$  for  $u_1 = |\sigma_1|$ . Further, since

$$(\nabla_Z Z)f_0 = \pi_0^\perp Z \pi_0^\perp Z f_0 - \pi_0^\perp Z \pi_0 Z f_0 = \pi_0^\perp Z Z f_0,$$

we have

$$\mathbb{I}(Z, Z) = [\pi^\perp(\nabla_Z Z)]f_0 = \pi_1^\perp Z Z f_0 = \sigma_2.$$

In particular,  $\mathbb{I}^{2,0}$  is a holomorphic section of  $K^2 \otimes \ell_2$ . Set  $s_2 = |\sigma_2|$ . Then

$$S(Z, Z, \bar{Z}, \bar{Z}) = \langle \mathbb{I}(Z, Z), \mathbb{I}(Z, Z) \rangle = |\sigma_2|^2 = s_2^2.$$

Thus in a chart  $U$  in which neither  $u_1$  nor  $s_2$  vanishes we have a local  $U(2, 1)$  frame given by  $f_1, f_2, f_0$  where  $f_1 = \sigma_1/u_1$ ,  $f_2 = \sigma_2/s_2$ . Straightforward calculations as before give

$$\begin{aligned} Z f_1 &= (Z \log u_1) f_1 + u_1^{-1} s_2 f_2, \\ Z f_2 &= (Z \log s_2) f_2, \\ Z f_0 &= u_1 f_1. \end{aligned} \tag{5.3}$$

From this we can read off the connexion 1-form for the projectively flat connexion  $\nabla$  in this frame. Now let  $\varphi : V|U \rightarrow U \times \mathbb{C}^2$  be the local trivialisation corresponding to  $f_1, f_2$ . Then

$$\varphi \circ \bar{\partial}_E \circ \varphi^{-1} = d\bar{z} \left[ \frac{\partial}{\partial \bar{z}} - \begin{pmatrix} \bar{Z} \log u_1 & s_2/u_1 \\ 0 & \bar{Z} \log s_2 \end{pmatrix} \right].$$

To deal with zeroes of  $u_1$  and  $s_2$ , we may assume  $U$  only has these at  $z = 0$ , to order  $p$  and  $q$  respectively. In such a chart we take the local frame  $\tilde{f}_1 = w^{-p}f_1$ ,  $\tilde{f}_2 = w^{-q}f_2$ , where  $w = z/|z|$ , and in the corresponding trivialisation  $\tilde{\varphi}$  we have

$$\tilde{\varphi} \circ \nabla^{0,1} \circ \tilde{\varphi}^{-1} = d\bar{z} \begin{bmatrix} \frac{\partial}{\partial \bar{z}} - \left( \bar{Z} \log(u_1/|z|^p) & -s_2 w^{p-q}/u_1 \right) \\ 0 & \bar{Z} \log(s_2/|z|^q) \end{bmatrix}.$$

We obtain a local holomorphic trivialisation, with respect to  $\bar{\partial}_E$ , by applying a gauge transformation of the form

$$R = \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix} \begin{pmatrix} |z|^p/u_1 & 0 \\ 0 & |z|^q/s_2 \end{pmatrix},$$

i.e., for  $\chi = R\tilde{\varphi}$  we have  $\chi \circ \bar{\partial}_E \circ \chi^{-1} = d\bar{z}(\partial/\partial \bar{z})$ . This requires

$$\partial a / \partial \bar{z} = -s_2^2 z^{p-q} / u_1^2. \quad (5.4)$$

When  $U$  is contractible this equation has a solution since  $s_2^2/u_1^2 \sim |z|^{2(q-p)}$  near  $z = 0$ , therefore the right hand side of (5.4) is smooth throughout  $U$ .

Now cover  $\Sigma_c$  by charts  $(U_j, z_j)$  of the type used above, and index the local objects living over  $U_j$  by  $j$ . Thus for  $V$  we have transition relations  $\tilde{\varphi}_j = \tilde{c}_{jk}\tilde{\varphi}_k$  where

$$\tilde{c}_{jk} = \begin{pmatrix} \frac{dz_j/dz_k}{|dz_j/dz_k|} & 0 \\ 0 & \frac{(dz_j/dz_k)^2}{|dz_j/dz_k|^2} \end{pmatrix} \begin{pmatrix} w_j^{p_j} w_k^{-p_k} & 0 \\ 0 & w_j^{q_j} w_k^{-q_k} \end{pmatrix}.$$

Therefore  $\chi_j = b_{jk}\chi_k$  where

$$b_{jk} = R_j \tilde{c}_{jk} R_k^{-1} = \begin{pmatrix} z_j^{p_j} z_k^{-p_k} dz_j/dz_k & \lambda_{jk} \\ 0 & z_j^{q_j} z_k^{-q_k} (dz_j/dz_k)^2 \end{pmatrix}, \quad (5.5)$$

for

$$\lambda_{jk} = a_j z_j^{q_j} z_k^{-q_k} (dz_j/dz_k)^2 - a_k z_j^{p_j} z_k^{-p_k} dz_j/dz_k, \quad (5.6)$$

and we have used the fact that

$$u_{1k}/u_{1j} = |dz_j/dz_k|, \quad s_{2k}/s_{2j} = |dz_j/dz_k|^2.$$

In particular, this shows that  $V$  is an extension of the line bundle  $K^{-1}(B)$  by the line bundle  $K^{-2}(D)$  where  $D$  is the divisor of zeroes of  $\Pi^{2,0}$ , and therefore we have fixed an isomorphism  $L \simeq \mathcal{O}(D)$ . As earlier, the extension class of  $V$  is given by the Čech cohomology class  $\eta$  of the 1-cocycle  $\{(\eta_{jk}, U_j, U_k)\}$  where

$$\begin{aligned} \eta_{jk} &= z_j^{-p_j} z_k^{p_k} (dz_k/dz_j) \lambda_{jk} dz_k \\ &= a_j z_j^{q_j - p_j} dz_j - a_k z^{q_k - p_k} dz_k. \end{aligned}$$

This is plainly a coboundary for Čech cohomology in smooth sections of  $K(B-D)$  of the form  $\delta\tau$  where  $\{(\tau_j, U_j)\}$  has  $\tau_j = a_j z^{q_j - p_j} dz_j$ . Under the Dolbeault isomorphism this corresponds to

$$\bar{\partial}\tau_j = -\frac{s_{2j}^2}{u_{1j}^2} dz_j d\bar{z}_j,$$

which is the local expression for  $-S/\gamma$ .  $\square$

**5.2. Surfaces arising from zero extension class.** Theorems 2.3 and 3.1 imply that Higgs bundles for non-holomorphic minimal surfaces with  $\mathcal{Q} = 0$  are exactly the length-three Hodge bundles. When  $\mathcal{Q} = 0$  we have a trivial extension bundle and can take

$$E_1 \oplus E_2 \oplus E_3 = K(-D_2) \oplus 1 \oplus K^{-1}(D_1),$$

to get  $\Phi : E_i \rightarrow KE_{i+1}$ . Conversely, any length-three Hodge bundle is projectively equivalent to one of this form and has  $\text{tr } \Phi^2 = 0$ . We will show that  $f$  is still related to a holomorphic map, via its harmonic sequence (in the sense of Erdem & Glazebrook [15],  $f$  is isotropic but non-holomorphic). To explain this, we first recall (from e.g., [4, 14]), the notion of the *Gauss transforms*  $\varphi_1, \varphi_2$  of  $f$ .

The line bundles  $\ell_1, \ell_2$  defined by (3.3) both lie inside the subset  $E_+$  consisting of fibre vectors which have positive length with respect to the  $\mathbb{C}^{2,1}$  metric. Let

$$\mathbb{C}_+^{2,1} = \{v \in \mathbb{C}^{2,1} : \langle v, v \rangle > 0\}.$$

Then its projective space  $\mathbb{P}\mathbb{C}_+^{2,1}$  is an open submanifold of  $\mathbb{C}\mathbb{P}^2$  on which  $G = PU(2, 1)$  acts transitively. We identify it with the orbit of the line  $[e_2]$ , with isotropy subgroup  $H_2 \simeq P(U(1, 1) \times U(1))$ , so that  $\mathbb{P}\mathbb{C}_+^{2,1} \simeq G/H_2$ . In fact we can think of it as the complex version of two dimensional de Sitter space, and will henceforth denote it by  $\mathbb{C}dS^2$ . Its tangent space at the base point  $[e_2]$  is identified with the orthogonal complement  $\mathfrak{m}_2 = \mathfrak{h}_2^\perp \subset \mathfrak{su}(2, 1)$  with respect to the Killing form  $-\frac{1}{2} \text{tr}(AB)$ , and the latter gives  $\mathfrak{m}_2$  an indefinite Hermitian metric. This extends to the tangent bundle, isomorphic to  $G \times_{H_2} \mathfrak{m}_2$ , and makes  $\mathbb{C}dS^2$  a pseudo-Hermitian symmetric space. Clearly  $\mathbb{P}E_+ \simeq \mathcal{D} \times_\rho \mathbb{C}dS^2$  and therefore  $\ell_1, \ell_2$  each determine a smooth  $\rho$ -equivariant map  $\varphi_1, \varphi_2 : \mathcal{D} \rightarrow \mathbb{C}dS^2$ , and these will be conformal harmonic with respect to the pseudo-Hermitian metric (they are isotropic in the sense of [15]). Following the terminology of harmonic sequences, we call  $\varphi_1$  the  $\partial$ -Gauss transform of  $f$ , and  $\varphi_2$  the  $\bar{\partial}$ -Gauss transform of  $f$ . An immersion into  $\mathbb{C}dS^2$  is *timelike* when its induced metric is negative definite (away from branch points).

**PROPOSITION 5.3.** *Let  $f : \mathcal{D} \rightarrow \mathbb{C}\mathbb{H}^2$  be  $\rho$ -equivariant and not  $\pm$ -holomorphic. Then  $\mathcal{Q} = 0$  if and only if the  $\bar{\partial}$ -transform  $\varphi_2 : \mathcal{D} \rightarrow \mathbb{C}dS^2$  of  $f$  is a timelike  $\rho$ -equivariant holomorphic map, branched at the divisor of complex points  $D_2$  of  $f$ .*

*Proof.* Let  $\varphi_2 : \mathcal{D} \rightarrow \mathbb{C}dS^2$  be the  $\bar{\partial}$ -Gauss transform  $f$ . Write the differential of  $\varphi_2$  as  $d\varphi_2' = \partial\varphi_2' + \bar{\partial}\varphi_2'$  in terms of the type decomposition

$$\varphi_2'^{-1}T_{\ell_2}\mathbb{C}dS^2 = \text{Hom}(\ell_2, \ell_2^\perp) \oplus \text{Hom}(\ell_2^\perp, \ell_2).$$

In local coordinates

$$\bar{\partial}\varphi_2'(\bar{Z}) = \pi_2^\perp \bar{Z}.$$

But from (3.7) and the fact that  $\ell_0, \ell_1, \ell_2$  are mutually orthogonal we have

$$\begin{aligned} \mathcal{Q} &= \langle \pi_0^\perp Z \pi_0^\perp Z f_0, \pi_0^\perp \bar{Z} f_0 \rangle, \\ &= -\langle \pi_0^\perp Z f_0, \pi_2^\perp \bar{Z} \pi_0^\perp \bar{Z} f_0 \rangle. \end{aligned}$$

Therefore if neither  $\pi_0^\perp Z f_0$  nor  $\pi_0^\perp \bar{Z} f_0$  is zero,  $\mathcal{Q}$  vanishes if and only if  $\pi_2^\perp \bar{Z}$  is identically zero on  $\ell_2$ . Hence  $\bar{\partial}\varphi_2'$  vanishes and  $\varphi_2$  is holomorphic.

Finally, we claim that when  $\mathcal{Q} = 0$  the induced metric for  $\varphi_2$  is  $-u_2^2|dz|^2$ . To see this, let  $(f_0, f_1, f_2)$  be a local Toda frame for  $f$ . Then by definition  $\varphi_2$  is given locally by the family of lines  $[f_2]$ . To calculate the differential we use (3.8) to deduce that in this frame  $d\varphi_2$  is represented by

$$\begin{pmatrix} & 0 & \\ Q/u_1u_2 & & 0 \\ & u_2 & \end{pmatrix} dz + \begin{pmatrix} & -\bar{Q}/u_1u_2 & \\ 0 & & u_2 \\ & 0 & \end{pmatrix} d\bar{z}.$$

Here we use blank spaces to indicate the Lie subalgebra  $\mathfrak{h}_2 \subset \mathfrak{g}$  of isotropy group  $H_2$ : relative to the frame  $d\varphi_2$  takes values in  $\mathfrak{h}_2^\perp$ . Therefore the induced metric is

$$-\frac{1}{2} \operatorname{tr}(d\varphi_2)^2 = \left(\frac{|Q|^2}{u_1^2u_2^2} - u_2^2\right)|dz|^2.$$

□

Note that, since  $\partial\varphi_2' : \ell_2 \rightarrow K\ell_0$ ,  $f$  is the  $\partial$ -Gauss transform of  $\varphi_2$ .

**6. Moduli.** Theorems 2.3 and 5.1 provide parameterisations for different components of the set

$$\mathcal{V} = \{(\rho, f) : \rho \text{ irreducible, } f \text{ branched minimal}\}/G,$$

where the quotient is by the simultaneous action of  $G$  as conjugation of  $\rho$  and ambient isometry of  $f$ . By those theorems it is natural to write  $\mathcal{V}$  as a disjoint union of the sets

$$\mathcal{V}(d_1, d_2) = \{(\rho, f) \in \mathcal{V} : f \text{ has } d_1 \text{ anti-complex points and } d_2 \text{ complex points}\}$$

and

$$\mathcal{W}^+(b, l) = \{(\rho, f) \in \mathcal{V} : f \text{ holomorphic with } \deg(B) = b, \deg(L) = l\},$$

$$\mathcal{W}^-(b, l) = \{(\rho, f) \in \mathcal{V} : (\bar{\rho}, \bar{f}) \in \mathcal{W}^+(b, l)\}.$$

These last two spaces are bijective under  $(\rho, f) \mapsto (\bar{\rho}, \bar{f})$ . We will now show that the parameterisations give each component the structure of a complex manifold.

**6.1. The structure of  $\mathcal{V}(d_1, d_2)$ .** As explained at the end of §2, each point of  $\mathcal{V}(d_1, d_2)$  is parametrised by a quadruple  $(\Sigma_c, D_1, D_2, \xi)$ . To understand the space of these quadruples we must understand how  $H^1(\Sigma_c, K^{-2}(D_1 + D_2))$  varies with  $(\Sigma_c, D_1, D_2)$ . Note that

$$\deg(K^{-2}(D_1 + D_2)) = d_1 + d_2 - 4(g - 1) < 0,$$

by the inequalities (2.17). Whenever a holomorphic line bundle  $\mathcal{L}$  over  $\Sigma_c$  has negative degree  $d$  its first cohomology has, by the Riemann-Roch theorem, dimension

$$h^1(\mathcal{L}) = g - 1 - d.$$

Therefore as  $\mathcal{L}$  moves over  $\operatorname{Pic}_d(\Sigma_c)$ , the Picard component of degree  $d$  line bundles, the dimension of  $H^1(\Sigma_c, \mathcal{L})$  is constant. Now  $\Sigma_c \times \operatorname{Pic}_d(\Sigma_c)$  carries a tautological line bundle  $\mathcal{P}$  (sometimes called a Poincaré line bundle) whose fibre over  $(p, \mathcal{L})$  is the fibre of  $\mathcal{L}$  at  $p$ . The vector spaces  $H^1(\Sigma_c, \mathcal{L})$  are the fibres of the higher direct image

$R^1\pi_*(\mathcal{P})$  for the projection  $\pi : \Sigma_c \times \text{Pic}_d(\Sigma_c) \rightarrow \text{Pic}_d(\Sigma_c)$  to the second factor. By a theorem of Grauert [27, III, Cor 12.9] their constant dimension implies they form a vector bundle over  $\text{Pic}_d(\Sigma_c)$ . In particular, for  $d = d_1 + d_2 - 4(g-1)$  this bundle has rank

$$h^1(K^{-2}(D_1 + D_2)) = 5g - 5 - d_1 - d_2.$$

The pairs  $(D_1, D_2)$  are parametrised by the product of symmetric products  $S^{d_1}\Sigma_c \times S^{d_2}\Sigma_c$  (in which the co-prime pairs occupy an open subvariety). The bundle can be pulled back along the holomorphic map

$$S^{d_1}\Sigma_c \times S^{d_2}\Sigma_c \rightarrow \text{Pic}_d(\Sigma_c); \quad (D_1, D_2) \mapsto K^{-2}(D_1 + D_2),$$

and the total space of the pullback parametrises the data  $(D_1, D_2, \xi)$ . It is a connected non-singular complex manifold of dimension  $5g - 5$ .

As  $c$  varies over the Teichmüller space  $\mathcal{T}_g$  of  $\Sigma$  we can take the disjoint union  $\mathcal{C}_g = \cup_{c \in \mathcal{T}_g} \Sigma_c$ , and likewise for any of the objects  $S^d\Sigma_c$  or  $\text{Pic}(\Sigma_d)$  above. In each of these cases we obtain a *complex analytic family over  $\mathcal{T}_g$* , i.e., the total space is a complex manifold for which the projection onto  $\mathcal{T}_g$  is holomorphic map, and although only a fibre bundle in the smooth category the fibre over  $c$  is a complex submanifold biholomorphic to the structure determined by  $c$ . In particular,  $\pi_c : \mathcal{C}_g \rightarrow \mathcal{T}_g$  is called the (*universal*) *Teichmüller curve*. About each point  $\mathcal{C}_g$  has a *permanent uniformising local parameter*, i.e., a complex chart  $(\mathcal{U}, \mathfrak{z})$  for which  $\mathfrak{z} = (z_1, \dots, z_{3g-3}, \zeta)$  has the properties that: (i) each non-empty intersection  $\mathcal{U}_c = \pi_c^{-1}(c) \cap \mathcal{U}$  is such that  $(\mathcal{U}_c, \zeta)$  is a chart on  $\Sigma_c$ ; (ii) the coordinates  $z_j$  are constant on the fibres. The existence of such a chart is an immediate consequence of Bers' construction of  $\mathcal{C}_g$  as a quotient of an open submanifold  $\mathcal{F}_g \subset \mathcal{T}_g \times \mathbb{C}$  by a properly discontinuous action of  $\pi_1\Sigma$  which preserves the fibres over  $\mathcal{T}_g$  [2] (this is also just the standard picture of Kodaira-Spencer for unobstructed deformations of complex structure [33]). It follows that one can put complex charts on the symmetric fibre-products of  $\mathcal{C}_g$  over  $\mathcal{T}_g$ , whose fibres are  $S^d\Sigma_c$ , to obtain non-singular complex analytic families over  $\mathcal{T}_g$ . The corresponding families of Picard components have been constructed by Earle [12].

Thus for each pair  $d_1, d_2$  satisfying (2.17) we obtain a manifold parametrising the data  $(\Sigma_c, D_1, D_2, \xi)$  with  $\deg(D_j) = d_j$ . Each is clearly a connected non-singular complex manifold of dimension  $8g - 8$ . Therefore we have proved the following theorem.

**THEOREM 6.1.** *Each set  $\mathcal{V}(d_1, d_2)$  can be given the structure of a non-singular complex manifold of dimension  $8g - 8$ . With this structure  $\mathcal{V}(d_1, d_2)$  is complex analytic family over Teichmüller space  $\mathcal{T}_g$ . The fibre over  $c \in \mathcal{T}_g$  is a complex submanifold biholomorphic to a holomorphic vector bundle over  $S^{d_1}\Sigma_c \times S^{d_2}\Sigma_c$  of rank  $5(g-1) - d_1 - d_2$ .*

**6.2. The structure of  $\mathcal{W}^\pm(b, l)$ .** By Theorem 5.1 the set  $\mathcal{W}^+(b, l)$  is parametrised by quadruples  $(\Sigma_c, B, L, \mathbb{C}\eta)$  with  $\eta \in H^1(\Sigma_c, KL^{-1}(B))$  and  $\eta \neq 0$ . We want to show this cohomology space has constant dimension as the pair  $(B, L)$  moves over  $S^b\Sigma_c \times \text{Pic}_l(\Sigma_c)$ . It suffices to see that, by the stability inequalities (5.2),

$$\deg(KL^{-1}(B)) = 2g - 2 + b - l < \frac{1}{2}b - (g-1) < 0,$$

and therefore

$$h^1(KL^{-1}(B)) = l + 1 - b - g.$$



By much the same argument as above, for each marked conformal structure  $c \in \mathcal{T}_g$  the triple  $(B, L, \mathbb{C}\eta)$  lies in a holomorphic  $\mathbb{C}\mathbb{P}^{l-b-g}$ -bundle over  $S^b\Sigma_c \times \text{Pic}_l(\Sigma_c)$ . The total space of this bundle has dimension

$$(b + g) + l - b - g = l.$$

As  $c$  moves through  $\mathcal{T}_g$  we obtain a complex analytic manifold, fibred over  $\mathcal{T}_g$ , of total dimension  $3(g - 1) + l$ .

**THEOREM 6.2.** *Each set  $\mathcal{W}^\pm(b, l)$  can be given the structure of a non-singular complex manifold of dimension  $3(g - 1) + l$ . It is a complex analytic family over  $\mathcal{T}_g$  whose fibre at  $c \in \mathcal{T}_g$  is a complex analytic submanifold biholomorphic to a holomorphic  $\mathbb{C}\mathbb{P}^{l-b-g}$ -bundle over  $S^b\Sigma_c \times \text{Pic}_l(\Sigma_c)$ .*

**6.3. Map from  $\mathcal{V}$  to  $\mathcal{R}(G)$ .** In order to understand when we can use minimal surface data to parametrise representations, we must understand the map

$$R : \mathcal{V} \rightarrow \mathcal{R}(G), \quad (\rho, f) \mapsto \rho. \tag{6.1}$$

We can expect this to be smooth. From the results above, this is likely to be most interesting on the components  $\mathcal{V}(d_1, d_2)$  since these have the same dimension as  $\mathcal{R}(G)$ . While a full understanding of this map will require further work, we can at least make some interesting comments about its behaviour on the fibres  $\mathcal{V}_c$  of  $\mathcal{V}$  over Teichmüller space. With a fixed conformal structure  $c$  we can identify  $\mathcal{R}(G)$  with the moduli space  $\mathcal{H}(\Sigma_c, G)$  of  $G$ -Higgs bundles. Then  $R$  is injective on  $\mathcal{V}_c$ , since it amounts to inclusion (equally, this is a consequence of the uniqueness theorem for twisted harmonic maps [9, 10]). Indeed

$$\mathcal{V}_c = \{(E, \Phi) \in \mathcal{H}(\Sigma_c, G) : \text{tr } \Phi^2 = 0\},$$

and so it plays the role of the *nilpotent cone* in  $\mathcal{H}(\Sigma_c, G)$ . Let us consider the structure of this in light of the discussion above. Recall that  $\|\Phi\|_{L^2}^2$  is a proper Morse-Bott function on  $\mathcal{H}(\Sigma_c, G)$  (at least at smooth points): we will normalise this by defining

$$\mathfrak{F}(E, \Phi) = \frac{i}{2} \int_{\Sigma} \text{tr}(\Phi \wedge \Phi^\dagger).$$

Whenever the twisted harmonic map determined by  $(E, \Phi)$  is conformal, we have

$$\mathfrak{F}(E, \Phi) = \int_{\Sigma} v_\gamma = \text{Area}_\gamma(\Sigma),$$

for the induced metric  $\gamma$ . Now fix a non-maximal value  $\tau$  for the Toledo invariant and consider the connected component  $\mathcal{H}(\Sigma_c, G)_\tau$ . Whenever  $d_2 = \frac{3}{2}\tau + d_1$  this component contains  $\mathcal{V}_c(d_1, d_2)$ . Inside the latter lies the locus  $\xi = 0$  consisting of length-three Hodge bundles, and this represents one connected critical manifold of  $\mathfrak{F}$  (cf. [22, §3]). Since  $\xi = 0$  exactly when  $\mathcal{Q} = 0$  we deduce from Prop. 3.5 that this is the level  $\mathfrak{F} = (4g - 4 - d_1 - d_2)\pi$ . As we move along the fibres of the bundle  $\mathcal{V}_c(d_1, d_2)$  Prop. 3.5 tells us that  $\mathfrak{F} < (4g - 4 - d_1 - d_2)\pi$ . Moreover, a comparison with [22, Prop 3.2] shows that the dimension of these fibres equals the Morse index of the critical manifold. Indeed, we conjecture that the bundle  $\mathcal{V}_c(d_1, d_2)$  is precisely the unstable manifold of the vector field  $-\text{grad } \mathcal{F}$  as a bundle over the critical manifold.

This seems to be a manifestation of Hausel’s theorem [28, Thm 5.2]. He proved that in the moduli space of stable  $GL(2, \mathbb{C})$ -Higgs bundles of odd degree, the downward Morse flow coincides with the nilpotent cone. Although Hausel only gave the proof for  $GL(2, \mathbb{C})$ , the ingredients hold equally well in the case of the smooth components of  $\mathcal{H}(\Sigma_c, G)$  for a real form  $G$  [23].

By contrast the image of  $\mathcal{W}_c^\pm(b, l)$  in  $\mathcal{H}(\Sigma_c, G)$  gives all the length-two Hodge bundles, which are the absolute minima of  $\mathcal{F}$  in their connected component. On the smooth locus of  $\mathcal{H}(\Sigma_c, G)$ , hence particularly when  $\tau \neq 2\mathbb{Z}$ , there is a single connected component of minima for fixed  $\tau$  (since  $\mathcal{F}$  is a perfect Morse-Bott function). Hence for fixed  $\tau$  the “components”  $\mathcal{W}_c^\pm(b, l)$  should really be thought of as strata of the space of holomorphic (or anti-holomorphic) branched immersions. Gothen [22, §3] explains how to view each critical manifold of length-two Hodge bundles as a moduli space of  $\alpha$ -stable Bradlow pairs. A Bradlow pair is a rank 2 vector bundle equipped with a holomorphic section. In our picture the Bradlow pair is  $(K \otimes V, \Phi)$ . The stratification of length-two Hodge bundles of fixed Toledo invariant by  $\mathcal{W}_c^\pm(b, l)$  amounts to fixing the degree  $b$  of  $\Phi$ .

**Appendix A. Uniqueness of minimal embeddings.** Here we give the proof of Proposition 4.5. We are assuming that  $\exp^\perp : TM^\perp \rightarrow N$  is a diffeomorphism, and therefore there is a radial distance function  $\rho : N \rightarrow \mathbb{R}_0^+$  given by  $\rho(p) = \|(\exp^\perp)^{-1}(p)\|$ . The idea of the proof is to show that the condition on  $dv_r/dr$  means each  $\varphi_r = \exp^\perp(r\eta)$  must have non-zero mean curvature, and that by local comparison every immersion must also have non-zero mean curvature at non-zero maximum values of  $\rho$ . Note that this is a local argument: we do not need the existence of global sections of  $TM^\perp$  of unit length (which will not, in general, exist).

First recall that for a family of immersions  $\varphi_t : M \times \mathbb{R} \rightarrow (N, g)$  with variational vector field  $V = \varphi_* \partial/\partial t$  a standard calculation gives

$$\frac{dv_\gamma(t)}{dt} \Big|_0 = d \star g(V, d\varphi) - g(V, H_\varphi)v_\gamma,$$

where  $H_\varphi = \text{tr}_\gamma \mathbb{I}_\varphi$  is the mean curvature for  $\gamma = \varphi^*g$ . In particular, for the mean curvature  $H_r$  of the map  $\varphi_r$ ,

$$d\rho(H_r) = g(\text{grad } \rho, H_r) = -\frac{1}{v_r} \frac{dv_r}{dr}, \tag{A.1}$$

since  $\text{grad } \rho$  is a normal variation.

Next we show that a local embedding  $\varphi$  which comes from an arbitrary non-vanishing local section  $\nu$  of  $TM^\perp$  must have non-zero mean curvature at any point at which  $|\nu|$  has a local maximum.

**LEMMA A.1.** *Let  $\varphi : \mathcal{U} \rightarrow N$  be an embedding of the form  $\varphi = \exp^\perp(\nu)$  for some local section  $\nu$  of  $TM^\perp$  which does not vanish on an open subset  $\mathcal{U} \subset M$ , and suppose  $u = \|\nu\| = \rho \circ \varphi$  has a local maximum at  $x \in \mathcal{U}$ . For each  $r > 0$  set  $\varphi_r = \exp^\perp(r\nu/u)$ , and let  $v_r$  be the volume form for  $\varphi_r^*g$ . Suppose that  $dv_r/dr > 0$  at  $x$  for each  $r$ , then  $\varphi$  must have non-zero mean curvature  $H_\varphi$  at  $x$ .*

*Proof.* Consider the expressions for the mean curvatures  $H_\varphi$  and  $H_r$ , considered as the tension fields  $\tau(\varphi)$  and  $\tau(\varphi_r)$ , in terms the tension fields for  $u = \rho \circ \varphi$  and the constant function  $r = \rho \circ \varphi_r$ . The composition formulas [13, 2.20] give

$$\begin{aligned} \tau(u) &= d\rho(H_\varphi) + \text{tr}_\gamma \nabla d\rho(d\varphi, d\varphi), \\ 0 = \tau(\rho \circ \varphi_r) &= d\rho(H_r) + \text{tr}_{\gamma_r} \nabla d\rho(d\varphi_r, d\varphi_r), \end{aligned}$$

where  $\gamma = \varphi^*g$  and  $\gamma_r = \varphi_r^*g$ . Now  $\tau(u) = \text{tr}_\gamma \text{Hess}(u)$  and at the local maximum  $x$  we have  $du = 0$ , which implies  $d\varphi = d\varphi_{u(x)}$  and  $\gamma = \gamma_{u(x)}$ . Therefore at  $x$  we have

$$d\rho(H_\varphi)|_x = \text{tr}_\gamma \text{Hess}(u)|_x + d\rho(H_{u(x)})|_x.$$

Since  $x$  is a local maximum we have  $\text{tr}_\gamma \text{Hess}(u)|_x \leq 0$ , and by assumption  $d\rho(H_r)|_x < 0$  for all  $r > 0$ , using (A.1). Thus  $H_\varphi$  cannot vanish at  $x$ .  $\square$

*Proof of Prop 4.5.* Suppose  $\psi : M \rightarrow N$  is any immersion transverse to the fibres of  $\exp^\perp$ , other than  $f$ . The function  $\rho \circ \psi$  must have a non-zero maximum at some  $y \in M$ . Then there is a local section  $\nu$  of  $TM^\perp$  and a local diffeomorphism  $\alpha$  on  $M$  for which  $\psi = \varphi \circ \alpha$  where  $\varphi = \exp^\perp(\nu)$ , and  $H_\psi|_y = H_\varphi|_{\alpha(y)}$  as elements of  $T_{\psi(y)}N$ . Now  $(\varphi, \alpha(y))$  satisfy the conditions of Lemma A.1, so  $H_\psi|_y \neq 0$ .  $\square$

## REFERENCES

- [1] M. ANDERSON, *Complete minimal hypersurfaces in hyperbolic  $n$ -manifolds*, Comment. Math. Helv., 58 (1983), pp. 264–290.
- [2] L. BERS, *Fiber spaces over Teichmüller spaces*, Acta. Math., 130 (1973), pp. 89–126.
- [3] S. B. BRADLOW, O. GARCIA-PRADA AND P. B. GOTHEN, *Surface group representations and  $U(p, q)$ -Higgs bundles*, J. Differential Geom., 64 (2003), pp. 111–170.
- [4] J. BOLTON AND L. M. WOODWARD, *Congruence theorems for harmonic maps from a Riemann surface into  $\mathbb{C}\mathbb{P}^n$  and  $S^n$* , J. London Math. Soc. (2), 45 (1992), pp. 363–376.
- [5] F. E. BURSTALL, *Harmonic tori in spheres and complex projective spaces*, J. Reine Angew. Math., 469 (1995), pp. 149–177.
- [6] J. CHEN AND G. TIAN, *Minimal surfaces in Riemannian 4-manifolds*, Geom. Funct. Anal., 7:5 (1997), pp. 873–916.
- [7] S. S. CHERN AND J. WOLFSON, *Minimal surfaces by moving frames*, Amer. J. Math., 105 (1983), pp. 59–83.
- [8] S. CHOI AND W. M. GOLDMAN, *Convex real projective structures on closed surfaces are closed*, Proc. Amer. Math. Soc., 118 (1993), pp. 657–661.
- [9] K. CORLETTE, *Flat  $G$ -bundles with canonical metrics*, J. Diff. Geom., 28 (1988), pp. 361–382.
- [10] S. K. DONALDSON, *Twisted harmonic maps and the self-duality equations*, Proc. London Math. Soc. (3), 55 (1987), pp. 127–131.
- [11] A. DOMIC AND D. TOLEDO, *The Gromov norm of the Kaehler class of symmetric domains*, Math. Ann., 276 (1987), pp. 425–432.
- [12] C. J. EARLE, *Families of Riemann surfaces and Jacobi varieties*, Ann. Math. (2), 107 (1978), pp. 255–286.
- [13] J. EELLS AND L. LEMAIRE, *Selected topics in harmonic maps*, CBMS Regional Conference Series in Mathematics, no. 50, AMS 1980.
- [14] J. EELLS AND J. WOOD, *Harmonic maps from surfaces to complex projective spaces*, Adv. in Math., 49 (1983), pp. 217–263.
- [15] S. ERDEM AND J. GLAZEBROOK, *Harmonic maps of Riemann surfaces to indefinite complex hyperbolic and projective spaces*, Proc. London Math. Soc. (3), 47 (1983), pp. 547–562.
- [16] O. GARCIA-PRADA, appendix in R. O. Wells, *Differential analysis on complex manifolds*, Springer Graduate Texts in Math. 65, Springer-Verlag, 2008.
- [17] O. GARCIA-PRADA, *Higgs bundles and surface group representations*, in “Moduli spaces and vector bundles”, pp. 265–310, LMS Lect. Note Ser., 359, Cambridge, 2009.
- [18] D. GILBARG AND N. TRUDINGER, *Elliptic partial differential equations of second order*, Classics in Mathematics, Springer-Verlag, Berlin, 2001.
- [19] W. M. GOLDMAN, *Higgs bundles and geometric structures on surfaces*, in “The many facets of geometry”, pp. 129–163, OUP, Oxford, 2010.
- [20] W. M. GOLDMAN, M. KAPOVICH AND B. LEEB, *Complex hyperbolic manifolds homotopy equivalent to a Riemann surface*, Comm. Anal. Geom., 9:1 (2001), pp. 61–95.
- [21] W. GOLDMAN AND R. WENTWORTH, *Energy of twisted harmonic maps of Riemann surfaces*, In the tradition of Ahlfors-Bers. IV, pp. 45–61, Contemp. Math., 432, Amer. Math. Soc., Providence, RI, 2007.
- [22] P. B. GOTHEN, *Topology of  $U(2, 1)$  representation spaces*, Bull. London Math. Soc., 34 (2002), pp. 729–738.
- [23] P. B. GOTHEN, private communication.

- [24] O. GUICHARD, *Groupes plongés quasi isométriquement dans un groupe de Lie*, Math. Ann., 330 (2004), pp. 331–351.
- [25] O. GUICHARD AND A. WIENHARD, *Anosov representations: domains of discontinuity and applications*, Invent. Math., 190 (2012), pp. 357–438.
- [26] R. GUNNING, *Lectures on vector bundles over Riemann surfaces*, Princeton University Press, 1967.
- [27] R. HARTSHORNE, *Algebraic geometry*, Graduate Texts in Mathematics 52, Springer-Verlag, Berlin, 1977.
- [28] T. HAUSEL, *Compactification of moduli of Higgs bundles*, J. Reine Angew. Math., 503 (1998), pp. 169–192.
- [29] N. J. HITCHIN, *The self-duality equations on a Riemann surface*, Proc. London Math. Soc. (3), 55 (1987), pp. 59–126.
- [30] N. J. HITCHIN, *Lie groups and Teichmüller space*, Topology, 31:3 (1992), pp. 449–473.
- [31] Z. HUANG, J. LOFTIN AND M. LUCIA, *Holomorphic cubic differentials and minimal Lagrangian surfaces in  $\mathbb{C}\mathbb{H}^2$* , Math. Res. Lett., 20 (2013), pp. 501–520.
- [32] Z. HUANG AND B. WANG, *Counting minimal surfaces in quasi-Fuchsian three-manifolds*, Trans. Amer. Math. Soc., 367 (2015), pp. 6063–6083.
- [33] K. KODAIRA, *Complex manifolds and deformation of complex structures*, Grundlehren der Math. Wissenschaften, 283. Springer-Verlag, New York, 1986.
- [34] K. KRASNOV AND J.-M. SCHLENKER, *Minimal surfaces and particles in 3-manifolds*, Geom. Dedicata, 126 (2007), pp. 187–254.
- [35] F. LABOURIE, *Flat projective structures on surfaces and cubic holomorphic differentials*, Pure Appl. Math. Q., 3 (2007), pp. 1057–1099.
- [36] J. C. LOFTIN, *Affine spheres and convex  $\mathbb{R}\mathbb{P}^n$ -manifolds*, Amer. J. Math., 123 (2001), pp. 255–274.
- [37] J. LOFTIN AND I. MCINTOSH, *Minimal Lagrangian surfaces in  $\mathbb{C}\mathbb{H}^2$  and representations of surface groups into  $SU(2, 1)$* , Geom. Dedicata, 162 (2013), pp. 67–93.
- [38] J. LOFTIN AND I. MCINTOSH, *Cubic differentials in the differential geometry of surfaces*, to appear in “Handbook of Teichmüller Theory” Vol. V.
- [39] Y.-G. OH, *Second variation and stabilities of minimal Lagrangian submanifolds in Kähler manifolds*, Invent. Math., 101 (1990), pp. 501–519.
- [40] J. R. PARKER AND I. PLATIS, *Open sets of maximal dimension in complex hyperbolic quasi-Fuchsian space*, J. Differential Geom., 73 (2006), pp. 319–350.
- [41] J. R. PARKER AND I. PLATIS, *Complex hyperbolic quasi-Fuchsian groups*, Geometry of Riemann surfaces, pp. 309–355, London Math. Soc. Lecture Note Ser., 368, CUP Cambridge, 2010.
- [42] J. SACKS AND K. UHLENBECK, *The existence of minimal immersions of 2-spheres*, Ann. Math., Second Series, 113 (1981), pp. 1–24.
- [43] C. SIMPSON, *Constructing variations of Hodge structure using Yang-Mills theory and applications to uniformization*, J. Amer. Math. Soc., 1 (1988), pp. 867–918.
- [44] C. H. TAUBES, *Minimal surfaces in germs of hyperbolic 3-manifolds*, Proceedings of the Casson Fest, 69100 (electronic), Geom. Topol. Monogr., 7 (2004).
- [45] D. TOLEDO, *Representations of surface groups in complex hyperbolic space*, J. Differential Geom., 29 (1989), pp. 125–133.
- [46] K. K. UHLENBECK, *Closed minimal surfaces in hyperbolic 3-manifolds*, Ann. Math. Stud., 103 (1983), pp. 147–168.
- [47] C. P. WANG, *Some examples of complete hyperbolic affine 2-spheres in  $\mathbb{R}^3$* , in “Global differential geometry and global analysis”, pp. 271–280, Lecture Notes in Math., 1481, Springer, Berlin, 1991.
- [48] S. WEBSTER, *Minimal surfaces in a Kähler surface*, J. Diff. Geometry, 20 (1984), pp. 463–470.
- [49] R. O. WELLS, *Differential analysis on complex manifolds*, Springer GTM 65, 1980.
- [50] J. WOLFSON, *Minimal surfaces in Kähler surfaces and Ricci curvature*, J. Differential Geom., 29 (1989), pp. 281–294.
- [51] J. C. WOOD, *Holomorphic differentials and classification theorems for harmonic maps and minimal immersions*, in “Global Riemannian geometry (Durham, 1983)”, pp. 168–175, Ellis Horwood Ser. Math. Appl., Horwood, Chichester, 1984.
- [52] E. XIA, *The moduli of flat  $PU(2, 1)$  structures on Riemann surfaces*, Pacific J. Math., 195 (2000), pp. 231–255.