

Regularity of CR-mappings of codimension one into Levi-degenerate hypersurfaces

ILYA KOSSOVSKIY, BERNHARD LAMEL, AND MING XIAO

We provide regularity results for CR-maps from a real hypersurface in n -dimensional complex space to a *Levi-degenerate target* hypersurface in a $n + 1$ -dimensional space. We address both the real-analytic and the smooth case. Our results allow immediate applications to the study of proper holomorphic maps between Bounded Symmetric Domains.

1	Introduction	151
2	Preliminaries	156
3	Normalization	158
4	Proof of Theorem 1	163
5	Proof of Theorem 2	165
6	Proof of Theorem 3	174
7	Proof of Theorem 4	177
	References	178

1. Introduction

This paper is devoted to establishing smooth and real-analytic versions of the *Schwarz reflection principle* for holomorphic maps in several complex variables. In the real-analytic version of the reflection principle, we investigate conditions under which a *CR-map* between real submanifolds in complex

space (or a holomorphic map between *wedges* attached to real submanifolds) extends holomorphically to an open neighborhood of the source manifold. In the smooth version, we ask for conditions under which a CR-map between real submanifolds in complex space has higher regularity than the given one. Problems of this type have attracted considerable attention since the work of Fefferman [Fe], Lewy [Le], and Pinchuk [Pi]. In the *equidimensional* case, the reflection principle is understood quite well due to the extensive research in this direction. We refer the reader to e.g. [BER, Fr1, KL1, KL2] for detailed surveys and references related to this research, as well as for the most up-to-date results.

In this paper, we study aspects of the regularity problem for CR-mappings between CR-manifolds M and M' of *different* dimension. This has been an extensively developing direction since the pioneering work of Webster [W], Faran [Fa], and Forstnerič [Fr1]. We shall note that the case of different dimensions is far more difficult than the equi-dimensional one, and much less is known in this setting. For an overview of existing results in the real-analytic case, we refer to the recent work of Berhanu and the first author [BX1].

The regularity problem in the *smooth category* rather than in the real-analytic one (in what follows, by “smooth” we refer to the C^∞ smoothness, if not otherwise stated) seems to be even more difficult due to lack of techniques. Starting from the work of Forstnerič [Fr1] and Huang [Hu1], [Hu2], the expected type of regularity of a finitely smooth CR-map between smooth CR-manifolds is *its C^∞ smoothness at a generic point*. One of the main tools for obtaining results in this line was introduced in the work [L1, L2, L3] by the second author, which is the notion of *k -nondegeneracy* of a CR-mapping. The latter is used for studying differential systems associated with CR-mappings. In particular, this tool was applied by Berhanu and the third author for studying the situation when the target manifold is *Levi-nondegenerate*. In the work [BX1], a smooth version of the reflection principle is established for CR-mappings from an abstract CR-manifold to a strongly pseudoconvex hypersurface. In particular, it solves a conjecture formulated earlier by Huang [Hu2] and also reproves a conjecture of Forstnerič [Fr1] consequently. In [BX2], this type of result is extended for CR-mapping into Levi-nondegenerate CR-submanifolds of hypersurface type with certain conditions on the signature. These results in particular show that if $F : M \rightarrow M'$ is a CR-transversal CR-mapping of class C^2 from a real-analytic (resp. smooth) strictly pseudoconvex hypersurface $M \subset \mathbb{C}^n$ into a real-analytic (resp. smooth) Levi-nondegenerate hypersurface $M' \subset \mathbb{C}^{n+1}$, then F is real-analytic (resp. smooth) on a dense open subset of M (we

mention that when F is assumed to be C^∞ , the result in the real-analytic case was proved in [EL]).

However, the case when the target is *Levi-degenerate* remains widely open, in both smooth and real-analytic categories, and very little is known in this setting. In the real-analytic case, a number of very interesting results in the latter direction were obtained in the recent paper of Mir [Mi1].

The main goal of this paper is to extend the reflection principle for CR-maps of real hypersurfaces in complex space to the setting when the target hypersurface $M' \subset \mathbb{C}^{n+1}$ is *Levi-degenerate*, while the source $M \subset \mathbb{C}^n$ is strictly pseudoconvex.

First, we obtain in the paper the generic analyticity property (resp. the generic smoothness property) for finitely smooth CR-maps between real-analytic (resp. smooth) real hypersurfaces of different dimensions with minimal assumptions for the target. Namely, in the real-analytic case, we assume the target M' to be merely *holomorphically nondegenerate*. Clearly, for any given source, the latter assumption can not be relaxed further (see Example 1.1 below). In the smooth case, we assume the *finite nondegeneracy* of the target. For definitions of different notions of nondegeneracy, see Section 2.

Second, we establish in the paper the *everywhere analyticity* (resp. *everywhere smoothness*) of CR-maps in the case when the target belongs to the class of *uniformly 2-nondegenerate hypersurfaces*. The latter class of hypersurfaces is of fundamental importance in Complex Analysis and Geometry. Uniformly 2-nondegenerate hypersurfaces have been recently studied intensively (e.g. Ebenfelt [E1, E2], Kaup and Zaitsev [KaZa], Fels and Kaup [FK1, FK2], Isaev and Zaitsev [IZ13], Medori and Spiro [MS], Kim and Zaitsev [KiZa], Beloshapka and the first author [BK]). These hypersurfaces naturally occur as boundaries of Bounded Symmetric Domains (see, e.g., [KaZa], [XY] for details), and in this way CR-maps into uniformly 2-nondegenerate hypersurfaces become important for understanding proper holomorphic maps between the respective Bounded Symmetric Domains (on the latter subject, see e.g. the work of Mok [Mo1, Mo2] and references therein). Uniformly 2-nondegenerate hypersurfaces occur as well as homogeneous holomorphically nondegenerate CR-manifolds [FK1, FK2]. We also note that the study of CR-embeddings of strictly-pseudoconvex hypersurfaces into 2-nondegenerate hypersurfaces performed in the present paper is important for understanding the geometry of the latter class of CR-manifolds (see, e.g., [BK]).

We shall now formulate our main results. Let us recall that a map $F: M \rightarrow M'$ between real hypersurfaces is said to be *CR-transversal* at

$p \in M$ if $T_{F(p)}^{(1,0)}M' + T_{F(p)}^{(0,1)}M' + dF(\mathbb{C}T_pM) = \mathbb{C}T_{F(p)}M'$. The definition of uniform 2-nondegeneracy is going to be given below in Definition 2.1.

Theorem 1. *Let $M \subset \mathbb{C}^n$ ($n \geq 2$) be a strongly pseudoconvex real-analytic (resp. smooth) hypersurface, and $M' \subset \mathbb{C}^{n+1}$ a uniformly 2-nondegenerate real-analytic (resp. smooth) hypersurface. Assume that $F = (F_1, \dots, F_{n+1}) : M \mapsto M'$ is a CR-transversal CR-mapping of class C^2 . Then F is real-analytic (resp. smooth) everywhere on M .*

We note that Theorem 1 has direct applications to the study of rigidity of proper holomorphic maps between bounded symmetric domains (see the work [XY] of Yuan and the third author, where certain rigidity results for holomorphic proper maps from the complex unit ball to the Type IV bounded symmetric domain D_m^{IV} are obtained). We also note that Theorem 1 somehow parallels a theorem proved by Mir [Mi1] and establishing the analyticity of CR-maps (at a generic point) in the situation when the source M is real-analytic and minimal while the target is the well known uniformly 2-nondegenerate hypersurface called *the tube over the future light cone*:

$$(1.1) \quad \mathbb{T}_{n+1} = \{(z_1, \dots, z_{n+1}) \in \mathbb{C}^{n+1} : (\operatorname{Im} z_1)^2 + \dots + (\operatorname{Im} z_n)^2 = (\operatorname{Im} z_{n+1})^2\}.$$

Next, in the more general setting of M' , we prove

Theorem 2. *Let $M \subset \mathbb{C}^n$ ($n \geq 2$) be a strongly pseudoconvex smooth hypersurface, and $M' \subset \mathbb{C}^{n+1}$ an everywhere finitely nondegenerate smooth hypersurface. Let $F = (F_1, \dots, F_{n+1}) : M \mapsto M'$ be a CR-transversal CR-mapping of class C^2 . Then F is smooth on a dense open subset of M .*

If the target hypersurface is real-analytic, a stronger assertion holds.

Theorem 3. *Let $M \subset \mathbb{C}^n$ ($n \geq 2$) be a strongly pseudoconvex smooth hypersurface, and $M' \subset \mathbb{C}^{n+1}$ a holomorphically nondegenerate real-analytic hypersurface. Let $F = (F_1, \dots, F_{n+1}) : M \mapsto M'$ be a CR-transversal CR-mapping of class C^2 . Then F is smooth on a dense open subset of M .*

A similar result holds in the real-analytic category: here we prove the stronger result, which does (only) require M' to be holomorphically nondegenerate.

Theorem 4. *Let $M \subset \mathbb{C}^n$ ($n \geq 2$) be a strongly pseudoconvex real-analytic hypersurface, and $M' \subset \mathbb{C}^{n+1}$ a holomorphically nondegenerate real-analytic hypersurface. Assume that $F = (F_1, \dots, F_{n+1}) : M \mapsto M'$ is a CR-transversal CR-mapping of class C^2 . Then F is real-analytic on a dense open subset of M .*

As was mentioned above, for any given M , one cannot drop the holomorphic nondegeneracy assumption when expecting the generic analyticity of CR-embeddings $F : M \mapsto M'$, $M' \in \mathbb{C}^{n+1}$ (see Example 1.1 below). The transversality assumption on F cannot be dropped either. See [BX2] for an example where F (being not transversal) is not smooth on any open subset of M . Thus, the assertion of Theorem 4 is in a sense optimal.

Example 1.1. Let $M \subset \mathbb{C}^n$, $n \geq 2$ be a strongly pseudoconvex hypersurface. Consider the holomorphically degenerate hypersurface $M' = M \times \mathbb{C} \subset \mathbb{C}^{n+1}$. Let f be a C^2 CR function on M which is not smooth on any open subset of M . Then $F(Z) := (Z, f(Z))$, $Z \in M$ is a CR-transversal map of class C^2 from M to M' . Clearly, F is not smooth on any open subset of M .

The following example shows also that one cannot expect F to be real-analytic everywhere on M in the setting of Theorems 3.

Example 1.2. Let $M \subset \mathbb{C}^2$ be the strongly pseudoconvex real hypersurface defined by

$$|z|^2 + |w|^2 + |1 - w|^{10} = 1$$

near $(0, 1)$, where (z, w) are the coordinates in \mathbb{C}^2 . Let $M' \subset \mathbb{C}^3$ be the holomorphically nondegenerate real hypersurface defined by

$$|z_1|^2 + |z_2|^2 + |z_3|^4 = 1,$$

where (z_1, z_2, z_3) are the coordinates in \mathbb{C}^3 . Consider the map

$$F = (z, w, (1 - w)^{5/2})$$

from one side of $M : \{|z|^2 + |w|^2 + |1 - w|^{10} < 1\}$ to \mathbb{C}^3 . It is easy to see F extends C^2 -smoothly up to M , sending M to M' . However, F is not even C^3 at the point $(0, 1)$.

We, however, hope that the following is true.

Conjecture 1.3. *For any integer $n \geq 2$, there exists an integer $k = k(n)$ such that the following holds. Let $M \subset \mathbb{C}^n$ ($n \geq 2$), $M' \subset \mathbb{C}^{n+1}$ be real-analytic (resp. smooth) hypersurfaces that are finitely nondegenerate (on some dense open subsets), and $F = (F_1, \dots, F_{n+1}) : M \rightarrow M'$ be a CR-transversal CR-mapping of class C^k . Then F is real-analytic (resp. smooth) on a dense open subset of M .*

(In the real-analytic version of the Conjecture, we may replace the condition on M by its holomorphic nondegeneracy).

The paper is organized as follows. In Section 2, we present some preliminaries on the degeneracy of CR-submanifolds and CR-mappings. Section 3 is devoted to a normalization result for a CR-map between hypersurfaces satisfying the assumptions of Theorems 1-3. It will be applied in later arguments. Theorems 1-4 will be proved in Sections 4-6.

Acknowledgement. The third author thanks Shiferaw Berhanu and Xiaojun Huang for helpful comments. All of the authors thank two anonymous referees for several helpful comments and suggestions.

The first author was supported by the FWF (Austrian Science Fund) Projects P29468 and I3472 and the GACR (Czech Grant Agency) grant 17-19437S. The second author is supported by the FWF (Austrian Science Fund) Project I3472. The third author is supported in part by National Science Foundation grant DMS-1800549.

2. Preliminaries

In this section, we recall various notions of degeneracy in CR geometry, and their relations. The following definition is introduced in [BHR].

Definition 2.1. Let M be a smooth generic submanifold in \mathbb{C}^N of CR-dimension n and real codimension d , and $p \in M$. Let $\rho = (\rho_1, \dots, \rho_d)$ be the defining function of M near p , and choose a basis L_1, \dots, L_n of CR vector fields near p . For a multiindex $\alpha = (\alpha_1, \dots, \alpha_n)$, write $L^\alpha = L_1^{\alpha_1} \dots L_n^{\alpha_n}$. Define the increasing sequence of subspaces $E_l(p)$ ($0 \leq l$) of \mathbb{C}^N by

$$E_l(p) = \text{Span}_{\mathbb{C}}\{L^\alpha \rho_{\mu,Z}(Z, \bar{Z})|_{Z=p} : 0 \leq |\alpha| \leq l, 1 \leq \mu \leq d\}.$$

Here $\rho_{\mu,Z} = (\frac{\partial \rho_\mu}{\partial z_1}, \dots, \frac{\partial \rho_\mu}{\partial z_N})$, and $Z = (z_1, \dots, z_N)$ are the coordinates in \mathbb{C}^N . We say that M is k -nondegenerate at p , $k \geq 1$ if

$$E_{k-1}(p) \neq E_k(p) = \mathbb{C}^N.$$

We say M is k -degenerate at p if $E_k(p) \neq \mathbb{C}^N$, and we say that M is uniformly k -nondegenerate if it is k -nondegenerate at every point $p \in M$.

We say M is (everywhere) finitely nondegenerate if M is $k(p)$ -nondegenerate at every $p \in M$ for some integer $k(p)$ depending on p . A smooth CR-manifold M of hypersurface type is Levi-nondegenerate at $p \in M$ if and only if M is 1-nondegenerate at p . This notion of degeneracy is then generalized to CR-mappings by the second author [La1] as follows.

Definition 2.2. Let $M \subset \mathbb{C}^N, M' \subset \mathbb{C}^{N'}$ be two generic CR-submanifolds of CR dimension n, n' , respectively. Let $H : M \rightarrow M'$ be a CR-mapping of class C^r near $p_0 \in M$. Let $\rho = (\rho_1, \dots, \rho_{d'})$ be local defining functions for M' near $H(p_0)$, and choose a basis L_1, \dots, L_n of CR vector fields for M near p_0 . If $\alpha = (\alpha_1, \dots, \alpha_n)$ is a multiindex, write $L^\alpha = L_1^{\alpha_1} \dots L_n^{\alpha_n}$. Define the increasing sequence of subspaces $E_l(p_0) (0 \leq l \leq r)$ of $\mathbb{C}^{N'}$ by

$$E_l(p_0) = \text{Span}_{\mathbb{C}}\{L^\alpha \rho_{\mu, Z'}(H(Z), \overline{H(\overline{Z})})|_{Z=p_0} : 0 \leq |\alpha| \leq l, 1 \leq \mu \leq d'\}.$$

Here $\rho_{\mu, Z'} = (\frac{\partial \rho_\mu}{\partial z'_1}, \dots, \frac{\partial \rho_\mu}{\partial z'_{N'}})$, and $Z' = (z'_1, \dots, z'_{N'})$ are the coordinates in $\mathbb{C}^{N'}$. We say that H is k_0 -nondegenerate at p_0 ($0 \leq k_0 \leq r$) if

$$E_{k_0-1}(p_0) \neq E_{k_0}(p_0) = \mathbb{C}^{N'}.$$

A manifold M is k_0 -nondegenerate if and only if the identity map from M to M is k_0 -nondegenerate. For a real-analytic submanifold, we also introduce the notion of holomorphic degeneracy.

Definition 2.3. A real-analytic submanifold $M \subset \mathbb{C}^N$ is *holomorphically nondegenerate* at $p \in M$ if there is no germ at p of a holomorphic vector field X tangent to M such that $X|_M \not\equiv 0$. We shall also say that M is *holomorphically nondegenerate* if it is so at every point of it.

We recall the following proposition about k -nondegeneracy and holomorphic nondegeneracy. For a proof of this, see [BER].

Proposition 2.4. *Let $M \subset \mathbb{C}^N$ be a connected real-analytic generic manifold with CR dimension n . Then the following conditions are equivalent:*

- M is holomorphically nondegenerate.
- M is holomorphically nondegenerate at some point $p \in M$.
- M is k -nondegenerate at some point $p \in M$ for some $k \geq 1$.

- *There exists V , a proper real-analytic subset of M and an integer $l = l(M)$, $1 \leq l(M) \leq n$, such that M is l -nondegenerate at every $p \in M \setminus V$.*

3. Normalization

In the section, we prove an auxiliary normalization result for CR-maps (Proposition 3.3 below), following the lines introduced by Huang [Hu3], in the following setting. Let $M \subset \mathbb{C}^n$ ($n \geq 2$) be a strongly pseudoconvex real-analytic (resp. smooth) hypersurface defined near a point $p_0 \in M$, and $M' \subset \mathbb{C}^{n+1}$ a real-analytic (resp. smooth) hypersurface which is Levi-degenerate at a point $q_0 \in M'$. Assume that $F = (F_1, \dots, F_{n+1}) : M \mapsto M'$ is a CR-transversal CR-mapping of class C^2 near p_0 with $F(p_0) = q_0$. We assume, after a holomorphic change of coordinates in \mathbb{C}^n , $p_0 = 0$ and that M is defined near 0 by

$$(3.1) \quad r(Z, \bar{Z}) = -\operatorname{Im}z_n + \sum_{i=1}^{n-1} |z_i|^2 + \psi(Z, \bar{Z}),$$

where $Z = (z_1, \dots, z_n)$ are the coordinates in \mathbb{C}^n , $\psi(Z, \bar{Z}) = O(|Z|^3)$ is real-analytic (resp. smooth) function defined near 0.

After a holomorphic change of coordinates in \mathbb{C}^{n+1} , we assume that $q_0 = F(p_0) = 0$ and that M' is locally defined near 0 by

$$(3.2) \quad \rho(W, \bar{W}) = -\operatorname{Im}w_{n+1} + \tilde{W}U\bar{W}^t + \phi(W, \bar{W}),$$

for some Hermitian $n \times n$ matrix U . Here

$$W = (\tilde{W}, w_{n+1}) = (w_1, \dots, w_n, w_{n+1})$$

are the coordinates in \mathbb{C}^{n+1} , $\phi(W, \bar{W}) = O(|W|^3)$ is a real-analytic (resp. smooth) function defined near 0.

If we write $F = (\tilde{F}, F_{n+1}) = (F_1, \dots, F_n, F_{n+1})$, then F satisfies:

$$(3.3) \quad -\frac{F_{n+1} - \overline{F_{n+1}}}{2i} + \tilde{F}U\bar{F}^t + \phi(F, \bar{F}) = 0,$$

along M . Since F is CR-transversal, we get $\lambda := \frac{\partial F_{n+1}}{\partial s}|_0 \neq 0$, where we write $z_n = s + it$ (cf. [BER]). Moreover, (3.3) shows that the imaginary part of F_{n+1} vanishes to second order at the origin, and so the number λ is

real. By applying the change of coordinates in \mathbb{C}^{n+1} : $\tau(w_1, \dots, w_n, w_{n+1}) = (w_1, \dots, w_n, -w_{n+1})$ if necessary, we may assume that $\lambda > 0$. Let us write

$$(3.4) \quad L_j = 2i \left(\frac{\partial r}{\partial \bar{z}_n} \frac{\partial}{\partial z_j} + \frac{\partial r}{\partial \bar{z}_j} \frac{\partial}{\partial z_n} \right), \quad 1 \leq j \leq n-1.$$

Then $\{L_j\}_{1 \leq j \leq n-1}$ forms a basis for the CR vector fields along M near p . By applying $\bar{L}_j, \bar{L}_j L_k, 1 \leq j, k \leq n-1$ to the equation (3.3) and evaluating at 0, we get:

$$\frac{\partial F_{n+1}}{\partial z_j}(0) = 0, \quad \frac{\partial^2 F_{n+1}}{\partial z_j \partial z_k}(0) = 0, \quad 1 \leq j, k \leq n-1.$$

Hence we have,

$$(3.5) \quad F_{n+1}(Z) = \lambda z_n + O(|Z|^2).$$

For $1 \leq j \leq n$, we write

$$(3.6) \quad F_j = a_j z_n + \sum_{i=1}^{n-1} a_{ij} z_i + O(|Z|^2),$$

for some $a_j \in \mathbb{C}, a_{ij} \in \mathbb{C}, 1 \leq i \leq n-1, 1 \leq j \leq n$. Or equivalently,

$$(3.7) \quad (F_1, \dots, F_n) = z_n(a_1, \dots, a_n) + (z_1, \dots, z_{n-1})A + (\hat{F}_1, \dots, \hat{F}_n),$$

where $A = (a_{ij})_{1 \leq i \leq n-1, 1 \leq j \leq n}$ is an $(n-1) \times n$ matrix, and $\hat{F}_j = O(|Z|^2), 1 \leq j \leq n$. We plug in (3.5) and (3.7) into (3.3) to get,

$$(3.8) \quad \begin{aligned} & \lambda |\tilde{Z}|^2 + O(|\tilde{Z}||z_n| + |z_n|^2) + o(|Z|^2) \\ & = \tilde{z} A U A^* \tilde{z}^t + O(|\tilde{Z}||z_n| + |z_n|^2) + o(|Z|^2), \end{aligned}$$

where we write $\tilde{Z} = (z_1, \dots, z_{n-1})$. Equip \tilde{Z} with weight 1, and z_n with weight 2. We then compare terms with weight 2 at both sides of (3.8) to get:

$$(3.9) \quad \lambda \mathbf{I}_{n-1} = A U A^*.$$

As a consequence, the matrix A has full rank $(n-1)$, and U has rank $(n-1)$ or n . Recall that M' is not 1-nondegenerate at $q=0$. We thus conclude that U has rank $(n-1)$. Moreover, note from (3.9) that U has $(n-1)$ positive eigenvalues. By a holomorphic change of coordinates in \mathbb{C}^{n+1} , we

may assume that $U = \text{diag}\{1, \dots, 1, 0\}$. M' is then of the following form near 0 :

$$(3.10) \quad \rho(W, \overline{W}) = -\text{Im}w_{n+1} + \sum_{j=1}^{n-1} |w_j|^2 + \phi(W, \overline{W}), \quad \phi = O(|W|^3).$$

Write $A = (B, \mathbf{b})$, where B is a $(n-1) \times (n-1)$ matrix, b is an $(n-1)$ -dimensional column vector. (3.9) yields that $B\overline{B}^t = \lambda \mathbf{I}_{n-1}$. We now apply the following holomorphic change of coordinates: $\widetilde{W} = WD$ or $W = \widetilde{W}D^{-1}$, where we set

$$D = \begin{pmatrix} \frac{1}{\sqrt{\lambda}}\overline{B}^t & \mathbf{c} & \mathbf{0} \\ \mathbf{0}^t & 1 & 0 \\ \mathbf{0}^t & 0 & 1 \end{pmatrix},$$

and $\mathbf{0}$ is the $(n-1)$ -dimensional zero column vector, \mathbf{c} is a $(n-1)$ -dimensional column vector to be determined. We compute

$$D^{-1} = \begin{pmatrix} \frac{1}{\sqrt{\lambda}}B & \mathbf{d} & \mathbf{0} \\ \mathbf{0}^t & 1 & 0 \\ \mathbf{0}^t & 0 & 1 \end{pmatrix},$$

where $\mathbf{d} = -\frac{1}{\sqrt{\lambda}}B\mathbf{c}$.

We write the new defining function of M' and the map as $\tilde{\rho}$ and $\tilde{F} = (\tilde{F}_1, \dots, \tilde{F}_{n+1})$ in the new coordinates $\widetilde{W} = (\tilde{w}_1, \dots, \tilde{w}_{n+1})$, respectively. We have

Lemma 3.1. *$\tilde{\rho}$ still has the form of (3.10). More precisely,*

$$\tilde{\rho}(\widetilde{W}, \overline{\widetilde{W}}) = -\text{Im}\tilde{w}_{n+1} + \sum_{j=1}^{n-1} |\tilde{w}_j|^2 + \tilde{\phi}(\widetilde{W}, \overline{\widetilde{W}}),$$

where $\tilde{\phi}(\widetilde{W}, \overline{\widetilde{W}}) = O(|\widetilde{W}|^3)$ is also a real-analytic (resp. smooth) function defined near 0.

Proof. This can be checked by a simple calculation and using the fact that

$$\begin{pmatrix} \frac{1}{\sqrt{\lambda}}B & \mathbf{d} \\ \mathbf{0}^t & 1 \end{pmatrix} \begin{pmatrix} \mathbf{I}_n & \mathbf{0} \\ \mathbf{0}^t & 0 \end{pmatrix} \begin{pmatrix} \frac{1}{\sqrt{\lambda}}\overline{B}^t & \mathbf{0} \\ \mathbf{d}^t & 1 \end{pmatrix} = \begin{pmatrix} \mathbf{I}_n & \mathbf{0} \\ \mathbf{0}^t & 0 \end{pmatrix}.$$

□

Moreover, since $\widetilde{F} = FD$, it is easy to see that

$$\frac{\partial \widetilde{F}_i}{\partial z_j}(0) = \delta_{ij}\sqrt{\lambda}, \quad 1 \leq i, j \leq n-1.$$

Here we denote by δ_{ij} the Kronecker symbol that takes value 1 when $i = j$ and 0 otherwise.

Lemma 3.2. *We can choose an appropriate \mathbf{c} such that*

$$(3.11) \quad \frac{\partial \widetilde{F}_n}{\partial z_j}(0) = 0, \quad 1 \leq j \leq n-1.$$

Proof: Note that $\widetilde{F}_n = (F_1, \dots, F_n) \begin{pmatrix} \mathbf{c} \\ 1 \end{pmatrix}$. Combining this with (3.7), we obtain,

$$\frac{\partial \widetilde{F}_n}{\partial z_j}(0) = 0, \quad 1 \leq j \leq n-1$$

is equivalent to $A \begin{pmatrix} \mathbf{c} \\ 1 \end{pmatrix} = \mathbf{0}$, where $\mathbf{0}$ is the $(n-1)$ -dimensional zero column vector. Recall $A = (B, \mathbf{b})$. We can thus choose $\mathbf{c} = -B^{-1}\mathbf{b}$.

In the following, for brevity, we still write W, F and ρ instead of $\widetilde{W}, \widetilde{F}$ and $\widetilde{\rho}$. We summarize the considerations of this section in the following

Proposition 3.3. *Let $M \subset \mathbb{C}^n (n \geq 2)$ be a strongly pseudoconvex real-analytic (resp. smooth) real hypersurface, $M' \subset \mathbb{C}^{n+1}$ a real-analytic (resp. smooth) real hypersurface. Assume that $F = (F_1, \dots, F_{n+1}) : M \mapsto M'$ is a CR-transversal CR-mapping of class C^2 near $p_0 \in M$ with $F(p_0) = q_0$, and that M' is Levi-degenerate at q_0 . Then, after appropriate holomorphic changes of coordinates in \mathbb{C}^n and \mathbb{C}^{n+1} respectively, we have $p_0 = 0, q_0 = 0$, and the following normalizations hold. M is defined by*

$$(3.12) \quad r(Z, \bar{Z}) = -\text{Im}z_n + \sum_{i=1}^{n-1} |z_i|^2 + \psi(Z, \bar{Z}), \quad \psi = O(|Z|^3)$$

near 0, and M' is defined by

$$(3.13) \quad \rho(W, \bar{W}) = -\text{Im}w_{n+1} + \sum_{j=1}^{n-1} |w_j|^2 + \phi(W, \bar{W}), \quad \phi = O(|W|^3)$$

near 0, where $Z = (z_1, \dots, z_n), W = (w_1, \dots, w_{n+1})$ are the coordinates of \mathbb{C}^n and \mathbb{C}^{n+1} , respectively. Furthermore, F satisfies:

$$(3.14) \quad \frac{\partial F_i}{\partial z_j}(0) = \delta_{ij}\sqrt{\lambda}, \quad 1 \leq i, j \leq n-1,$$

for some $\lambda > 0$, and moreover,

$$(3.15) \quad \frac{\partial F_n}{\partial z_j}(0) = 0, \quad 1 \leq j \leq n-1;$$

$$(3.16) \quad \frac{\partial F_{n+1}}{\partial z_j}(0) = 0, \quad 1 \leq j \leq n-1.$$

One can use the same arguments as given above to arrive at the following, more general conclusion, as pointed out by one of the anonymous referees; we record it here, even though we only need the version given above in this paper. In particular, the following formulation recovers as a special case ($m = n + \ell - 1$) a result of Berhanu and the third author [BX1, Lemma 4.2].

Proposition 3.4. *Let $M \subset \mathbb{C}^n (n \geq 2)$ be a strongly pseudoconvex real-analytic (resp. smooth) real hypersurface, $M' \subset \mathbb{C}^{n+\ell}$ a real-analytic (resp. smooth) real hypersurface. Assume that $F = (F_1, \dots, F_{n+\ell}) : M \mapsto M'$ is a CR-transversal CR mapping of class C^2 near $p_0 \in M$ with $F(p_0) = q_0$, and that the Levi-form of M' has, say m nonzero eigenvalues, all of the same sign. Then, after appropriate holomorphic changes of coordinates in \mathbb{C}^n and $\mathbb{C}^{n+\ell}$ respectively, we have $p_0 = 0, q_0 = 0$, and the following normalizations hold. M is defined by*

$$(3.17) \quad r(Z, \bar{Z}) = -\operatorname{Im}z_n + \sum_{i=1}^{n-1} |z_i|^2 + \psi(Z, \bar{Z}), \quad \psi = O(|Z|^3)$$

near 0, and M' is defined by

$$(3.18) \quad \rho(W, \bar{W}) = -\operatorname{Im}w_{n+\ell} + \sum_{j=1}^m |w_j|^2 + \phi(W, \bar{W}), \quad \phi = O(|W|^3)$$

near 0, where $Z = (z_1, \dots, z_n), W = (w_1, \dots, w_{n+\ell})$ are the coordinates of \mathbb{C}^n and $\mathbb{C}^{n+\ell}$, respectively. Furthermore, F satisfies:

$$(3.19) \quad \frac{\partial F_i}{\partial z_j}(0) = \delta_{ij}\sqrt{\lambda}, \quad 1 \leq i, j \leq n-1,$$

for some $\lambda > 0$, and moreover,

$$(3.20) \quad \frac{\partial F_{n+k}}{\partial z_j}(0) = 0, \quad 1 \leq j \leq n-1, \quad 1 \leq k \leq \ell.$$

In order to prove this more general assertion, one just has to follow the steps of the proof of Proposition 3.3 taking into account the more general codimension. This leads to a number of scalar quantities quantities to be vectors, but does not impact the main argument.

4. Proof of Theorem 1

In this section will make use of the normalization of M, M' in Proposition 3.3 and prove Theorem 1. We will see that in the setting of Theorem 1, the map is actually 2-nondegenerate. The result then follows in the smooth case by applying [L1, Theorem 2] and in the real-analytic case by applying [L2, Theorem 1.3]. Our first step therefore is the following computational Lemma for the uniformly 2–nondegenerate target hyperurface M' . For further results about normal forms along this line, see [E1].

Lemma 4.1. *Let $M \subset \mathbb{C}^n$ ($n \geq 2$) be a strongly pseudoconvex real-analytic (resp. smooth) hypersurface, and $M' \subset \mathbb{C}^{n+1}$ a uniformly 2–nondegenerate real-analytic (resp. smooth) hypersurface. Assume that $F = (F_1, \dots, F_{n+1}) : M \mapsto M'$ is a CR-transversal CR-mapping of class C^2 . Then F is 2–nondegenerate.*

Proof. We will write for $1 \leq k \leq n$,

$$(4.1) \quad \Lambda_k = 2i \left(\frac{\partial \rho}{\partial \bar{w}_{n+1}} \frac{\partial}{\partial \bar{w}_k} - \frac{\partial \rho}{\partial \bar{w}_k} \frac{\partial}{\partial \bar{w}_{n+1}} \right),$$

where $\{\Lambda_k\}_{1 \leq k \leq n}$ forms a basis for the CR vector fields along M' near 0. Note that

$$(4.2) \quad \begin{aligned} \Lambda_k &= (1 + 2i\phi_{n+1}) \frac{\partial}{\partial \bar{w}_k} - 2i(w_k + \phi_{\bar{k}}) \frac{\partial}{\partial \bar{w}_{n+1}}, \text{ if } 1 \leq k \leq n-1, \\ \Lambda_n &= (1 + 2i\phi_{n+1}) \frac{\partial}{\partial \bar{w}_n} - 2i(\phi_{\bar{n}}) \frac{\partial}{\partial \bar{w}_{n+1}}. \end{aligned}$$

Here and in the following, we write for $1 \leq i, j, k \leq n+1$, $\phi_{\bar{i}} = \phi_{\bar{w}_i} = \frac{\partial \phi}{\partial \bar{w}_i}$, $\phi_i = \phi_{w_i} = \frac{\partial \phi}{\partial w_i}$, $\phi_{i\bar{j}} = \phi_{w_i \bar{w}_j} = \frac{\partial^2 \phi}{\partial w_i \partial \bar{w}_j}$, $\phi_{i\bar{j}k} = \phi_{\bar{w}_i \bar{w}_j w_k} = \frac{\partial^3 \phi}{\partial \bar{w}_i \partial \bar{w}_j \partial w_k}$, etc.

Recall our notation $\rho_W := (\frac{\partial \rho}{\partial w_1}, \dots, \frac{\partial \rho}{\partial w_{n+1}})$. We compute

$$(4.3) \quad \rho_W(W, \overline{W}) = \left(\overline{w}_1 + \phi_1, \dots, \overline{w}_{n-1} + \phi_{n-1}, \phi_n, \frac{i}{2} + \phi_{n+1} \right)$$

We thus have

$$(4.4) \quad \Lambda_1 \rho_W(W, \overline{W}) = (h_{11}, \dots, h_{1(n+1)}),$$

where

$$(4.5) \quad \begin{aligned} h_{11} &= (1 + 2i\phi_{\overline{(n+1)}})(1 + \phi_{1\overline{1}}) - 2i(w_1 + \phi_{\overline{1}})\phi_{1\overline{(n+1)}}, \\ h_{12} &= (1 + 2i\phi_{\overline{(n+1)}})\phi_{2\overline{1}} - 2i(w_1 + \phi_{\overline{1}})\phi_{2\overline{(n+1)}}, \\ &\dots, \\ h_{1(n+1)} &= (1 + 2i\phi_{\overline{(n+1)}})\phi_{(n+1)\overline{1}} - 2i(w_1 + \phi_{\overline{1}})\phi_{(n+1)\overline{(n+1)}}. \end{aligned}$$

Hence

$$(4.6) \quad \Lambda_1 \rho_W(W, \overline{W}) = (1 + O(1), O(1), \dots, O(1)).$$

Here we write $O(m) = O(|W|^m)$ for any $m \geq 0$. Similarly, we have for $1 \leq k \leq n - 1$,

$$(4.7) \quad \Lambda_k \rho_W(W, \overline{W}) = (O(1), \dots, O(1), 1 + O(1), O(1), \dots, O(1)),$$

where the term $1 + O(1)$ is at the k^{th} position;

$$(4.8) \quad \Lambda_n \rho_W(W, \overline{W}) = (O(1), \dots, O(1), \phi_{n\overline{n}} + O(2), O(1)).$$

As a consequence, we have

$$(4.9) \quad \det \begin{pmatrix} \rho_W(W, \overline{W}) \\ \Lambda_1 \rho_W(W, \overline{W}) \\ \dots \\ \Lambda_n \rho_W(W, \overline{W}) \end{pmatrix} = \pm \frac{i}{2} \phi_{n\overline{n}} + O(2).$$

Recall that M' is uniformly 2–nondegenerate at 0, in particular, it is 1–degenerate at every point near 0. This implies (4.9) is identically zero near 0 along M' . Consequently, by applying $\Lambda_j, 1 \leq j \leq n$ to (4.9) and evaluating at 0, we obtain $\phi_{j\overline{nn}}(0) = 0$ for any $1 \leq j \leq n$.

Since M' is 2-nondegenerate, there exists a choice $j_0, k_0, 1 \leq j_0, k_0 \leq n$ such that $\Lambda_{j_0} \Lambda_{k_0} \rho_W(W, \bar{W})$ together with the $\Lambda_j \rho_W(W, \bar{W})$ and $\rho_W(W, \bar{W})$ spans \mathbb{C}^{n+1} for W close to 0. Hence, for that choice of j_0, k_0 , we have:

$$(4.10) \quad \phi_{\overline{j_0 k_0 n}}(0) \neq 0, \text{ for some } 1 \leq j_0, k_0 \leq n - 1.$$

Consequently, if we write

$$(4.11) \quad L_{j_0} L_{k_0} \rho_W(F, \bar{F})(0) := (\nu_1, \dots, \nu_{n-1}, \nu_n, \nu_{n+1}),$$

then ν_n is nonzero. Here L_j is as defined in (3.4). Indeed,

$$\begin{aligned} \nu_n &= \left. \frac{\partial^2 \phi_n(F, \bar{F})}{\partial \bar{z}_{j_0} \partial \bar{z}_{k_0}} \right|_0 = \sum_{i,j=1}^{n+1} \left. \frac{\partial^2 \phi_n}{\partial \bar{w}_i \partial \bar{w}_j} \right|_0 \left. \frac{\partial \bar{F}_i}{\partial z_{j_0}} \right|_0 \left. \frac{\partial \bar{F}_j}{\partial z_{k_0}} \right|_0 \\ &= \phi_{\overline{j_0 k_0 n}}(0) \left. \frac{\partial \bar{F}_{j_0}}{\partial z_{j_0}} \right|_0 \left. \frac{\partial \bar{F}_{k_0}}{\partial z_{k_0}} \right|_0 \neq 0. \end{aligned}$$

Moreover, it is easy to verify that

$$(4.12) \quad L_i \rho_W(F, \bar{F})(0) = (0, \dots, 0, \sqrt{\lambda}, 0, \dots, 0), \quad 1 \leq i \leq n - 1,$$

where $\sqrt{\lambda}$ is at the i^{th} position, and that

$$(4.13) \quad \rho_W(F, \bar{F})(0) = \left(0, \dots, 0, \frac{i}{2} \right).$$

Equations (4.13), (4.12) and (4.11) with $\nu_n \neq 0$ now imply that F is 2-nondegenerate at 0. □

Proof of Theorem 1. By the above-mentioned results of [L1, L2], F is real-analytic (resp. smooth) near 0, as required. □

5. Proof of Theorem 2

Let M' be as above and ρ as in (3.13). For any $1 \leq i_1 \leq \dots \leq i_l \leq n, q \in M'$, we define,

$$\Delta_{i_1 \dots i_l}(q) = \det \begin{pmatrix} \rho_W \\ \Lambda_1 \rho_W \\ \dots \\ \Lambda_{n-1} \rho_W \\ \Lambda_{i_1} \cdots \Lambda_{i_l} \rho_W \end{pmatrix} (q).$$

We first prove the following lemma.

Lemma 5.1. *Let M' be as above. Assume that M' is l -nondegenerate at 0 for some $l \geq 2$. Then there exist $1 \leq i_1 \leq \dots \leq i_l \leq n$, such that*

$$(5.1) \quad \Delta_{i_1 \dots i_l}(0) \neq 0.$$

Proof. We note that

$$(5.2) \quad \rho_W(0) = \left(0, \dots, 0, \frac{i}{2} \right),$$

$$(5.3) \quad \Lambda_j \rho_W(0) = (0, \dots, 0, 1, 0, \dots, 0), \quad 1 \leq j \leq n-1,$$

where 1 is at the j^{th} position. Thus $\rho_W(0), \Lambda_j \rho_W(0), 1 \leq j \leq n-1$, are linearly independent over \mathbb{C} . Then by the definition of l -nondegeneracy at 0, one easily sees that there exists $1 \leq i_1 \leq \dots \leq i_l \leq n$ such that

$$\det \begin{pmatrix} \rho_W \\ \Lambda_1 \rho_W \\ \dots \\ \Lambda_{n-1} \rho_W \\ \Lambda_{i_1} \cdots \Lambda_{i_l} \rho_W \end{pmatrix} (0) \neq 0.$$

□

Remark 5.2. In particular, when $l = 2$ in Lemma 5.1, there exist $1 \leq i_1 \leq i_2 \leq n$, such that $\Delta_{i_1 i_2}(0) \neq 0$. Note the n^{th} component of $\Lambda_{i_1} \Lambda_{i_2} \rho_W(0)$ is $\phi_{\frac{i_1 i_2 n}{2}}(0)$. By the form (5.2), (5.3) of $\rho_W(0)$ and $\Lambda_j \rho_W(0)$, we conclude that $\phi_{\frac{i_1 i_2 n}{2}}(0) \neq 0$.

We then prove the following proposition.

Proposition 5.3. *Let $M \subset \mathbb{C}^n (n \geq 2)$ be a strongly pseudoconvex real-analytic (resp. smooth) hypersurface, and $M' \subset \mathbb{C}^{n+1}$ be a real-analytic (resp. smooth) hypersurface. Assume that M' is either 1- or 2-nondegenerate*

at every point of it. Let $F = (F_1, \dots, F_{n+1}) : M \mapsto M'$ be a CR-transversal CR-mapping of class C^2 . Then F is real-analytic (resp. smooth) on a dense open subset of M .

Proof. We write Ω as the open subset of M where F is real-analytic (resp. smooth). Fix any $p_0 \in M$. Write $q_0 = F(p_0) \in M'$. We will need to prove $p_0 \in \bar{\Omega}$. We assume $p_0 = 0 \in M, q_0 = 0 \in M'$. By assumption, M' is either 1-nondegenerate or 2-nondegenerate at q_0 . We then split our argument in two cases.

Case I: M' is 1-nondegenerate at q_0 . That is, M' is Levi-nondegenerate near q_0 . Then it follows from Corollary 2.3 in [BX2] that $p_0 \in \bar{\Omega}$.

Case II: M' is 2-nondegenerate at q_0 . Let O be a small neighborhood of q_0 in \mathbb{C}^{n+1} . Let $V = O \cap M'$. We write V_1 as the set of 1-degeneracy of M' in V . More precisely,

$$V_1 = \{q \in V : M' \text{ is 1-degenerate at } q\}.$$

If there is a sequence $\{p_i\}_{i=1}^\infty \subset M$ converging to p_0 such that M' is 1-nondegenerate at each $F(p_i)$, i.e., $F(p_i) \in M \setminus V_1, i \geq 1$. Then by Case I, we have each $p_i \in \bar{\Omega}, i \geq 1$. Consequently, $p_0 \in \bar{\Omega}$. Thus we are only left with the case that there exists a neighborhood U of p on M such that $F(U) \subset V_1$. We apply then normalization to M, M' and the map F as in Proposition 3.3. Since M' is 2-nondegenerate at 0, we conclude by Lemma 5.1, $\Delta_{j_0 k_0}(0) = c \neq 0$, for some $1 \leq j_0 \leq k_0 \leq n$. We then further split into the following subcases.

Case II(a): There exist some $1 \leq j_0 \leq k_0 \leq n - 1$, such that, $\Delta_{j_0 k_0}(0) = c \neq 0$. Consequently, we have $\phi_{\overline{j_0 k_0 n}}(0) \neq 0$. Then similarly as in the proof of Theorem 1, we can show that F is finitely nondegenerate. Hence again by the results of [L1, L2], F is real-analytic (resp. smooth) at 0.

Case II(b): For any $1 \leq j \leq k \leq n - 1, \Delta_{jk}(0) = 0$. Moreover, there exists $1 \leq j_0 \leq n - 1$ such that, $\Delta_{j_0 n}(0) = c \neq 0$. Then by a similar argument as in Remark 5.2, we conclude $\phi_{\overline{jk n}}(0) = 0$, for any $1 \leq j \leq k \leq n - 1$, and $\phi_{\overline{j_0 n n}}(0) \neq 0$.

Note that $V_1 \subset \tilde{V}_1 \cap M'$, where \tilde{V}_1 is defined

$$(5.4) \quad \tilde{V}_1 := \{W \in O : \varphi(W, \overline{W}) = 0\},$$

with

$$(5.5) \quad \varphi(W, \overline{W}) = \det \begin{pmatrix} \rho_W \\ \Lambda_1 \rho_W \\ \dots \\ \Lambda_{n-1} \rho_W \\ \Lambda_n \rho_W \end{pmatrix} (W, \overline{W}).$$

Then we have

Claim 5.4. The \overline{w}_{j_0} -derivative of φ is nonzero at $q_0 = 0$.

Proof. It is equivalent to show that $\Lambda_{j_0} \varphi(0) \neq 0$. That is,

$$(5.6) \quad \Lambda_{j_0} \det \begin{pmatrix} \rho_W \\ \Lambda_1 \rho_W \\ \dots \\ \Lambda_{n-1} \rho_W \\ \Lambda_n \rho_W \end{pmatrix} (0) \neq 0.$$

Note that

$$\begin{aligned} \Lambda_{j_0} \det \begin{pmatrix} \rho_W \\ \Lambda_1 \rho_W \\ \dots \\ \Lambda_{n-1} \rho_W \\ \Lambda_n \rho_W \end{pmatrix} (0) &= \det \begin{pmatrix} \Lambda_{j_0} \rho_W \\ \Lambda_1 \rho_W \\ \dots \\ \Lambda_{n-1} \rho_W \\ \Lambda_n \rho_W \end{pmatrix} (0) + \det \begin{pmatrix} \rho_W \\ \Lambda_{j_0} \Lambda_1 \rho_W \\ \dots \\ \Lambda_{n-1} \rho_W \\ \Lambda_n \rho_W \end{pmatrix} (0) \\ &\quad + \dots + \det \begin{pmatrix} \rho_W \\ \Lambda_1 \rho_W \\ \dots \\ \Lambda_{j_0} \Lambda_{n-1} \rho_W \\ \Lambda_n \rho_W \end{pmatrix} (0) \\ &\quad + \det \begin{pmatrix} \rho_W \\ \Lambda_1 \rho_W \\ \dots \\ \Lambda_{n-1} \rho_W \\ \Lambda_{j_0} \Lambda_n \rho_W \end{pmatrix} (0). \end{aligned}$$

In the above equation, the first term is trivially zero. Then we note that in the row vector $\rho_W(0)$, or $\Lambda_i \rho_W(0)$, $1 \leq i \leq n$, the n^{th} component is zero. This is due to the fact that $\phi = O(|W|^3)$. Moreover, the n^{th} component in the row vector $\Lambda_{j_0} \Lambda_k \rho_W(0)$, $1 \leq k \leq n-1$, is $\phi_{\overline{j_0 k n}}(0)$, which is zero by the assumption. Consequently, the second term up to the n^{th} term in the above

equation are all zero. We also note the last term in the equation above is just equal to $\Delta_{j_0 n}(0)$, which is nonzero. Hence the lemma is established. \square

Recall that $F(U) \subset V_1 \subset \tilde{V}_1$. We have

$$(5.7) \quad \varphi(F(Z), \overline{F(Z)}) \equiv 0, \text{ for all } Z \in U \subset M.$$

Applying L_{j_0} to the above equation and evaluating at $Z = 0$, we have,

$$(5.8) \quad L_{j_0} \varphi(F, \overline{F})|_0 = \sum_{i=1}^{n+1} \varphi_{\bar{w}_i}(F, \overline{F})|_0 L_{j_0} \bar{F}_i|_0 = 0.$$

Note that by our normalization, $L_{j_0} \bar{F}_i(0) = 0$, if $i \neq j_0$. $L_{j_0} \bar{F}_{j_0}(0) \neq 0$. Moreover, by Claim 5.4, $\varphi_{\bar{w}_{j_0}}(0) \neq 0$. This is a contradiction to (5.8). Hence **Case II(b)** cannot happen in this setting.

Case II(c): $\Delta_{jk}(0) = \Delta_{jn}(0) = 0$, for all $1 \leq j, k \leq n - 1$, and $\Delta_{nn}(0) \neq 0$.

We let \tilde{V}_1 be defined by φ as above in (5.4), (5.5).

Claim 5.5. In the setting of this subcase, the \bar{w}_n -derivative of φ is nonzero at $q_0 = 0$.

Proof. Similar as Lemma Claim 5.4. \square

By Lemma Claim 5.5, we have $\varphi_{\bar{w}_n}(0) \neq 0$. Consequently, if we define $\tilde{\varphi}(W, \overline{W}) = \varphi(W, \overline{W})$, then

$$(5.9) \quad \tilde{\varphi}_{w_n}(0) \neq 0.$$

Note $F(U) \subset V_1 \subset \tilde{V}_1$. We have

$$\varphi(F(Z), \overline{F(Z)}) \equiv 0, \text{ for all } Z \in U.$$

Consequently,

$$(5.10) \quad \tilde{\varphi}(F(Z), \overline{F(Z)}) \equiv 0.$$

Recall that for all $Z \in U$,

$$(5.11) \quad \rho(F, \overline{F}) = 0,$$

$$(5.12) \quad L_i \rho(F, \bar{F}) = 0, \quad 1 \leq i \leq n - 1.$$

Combining (4.12), (4.13), (5.9), we conclude that

$$(5.13) \quad \det \begin{pmatrix} \rho_W(F, \bar{F}) \\ L_1 \rho_W(F, \bar{F}) \\ \dots \\ L_{n-1} \rho_W(F, \bar{F}) \\ \tilde{\varphi}_W(F, \bar{F}) \end{pmatrix} (0) \neq 0.$$

This implies that equations (5.10), (5.11), (5.12) form a nondegenerate system for F . Then it follows that F is real-analytic (resp. smooth) at 0 by a similar argument as in [L1, L2] or [BX1, BX2]. For the convenience of the readers, we sketch a proof here for the real-analytic category. The proof for the smooth category is similar (we remark on the differences). However, those readers unfamiliar with the strategy, we refer to the aforementioned papers for the necessary details.

We assume that M is defined near 0 by $\{(z, z_n) = (z, s + it) \in U_0 \times V : t = \phi(z, \bar{z}, s)\}$, where ϕ is a real-valued, real-analytic function with $\phi(0) = 0, d\phi(0) = 0$. Here $U_0 \subset \mathbb{C}^{n-1}$ and $V \subset \mathbb{R}$ are sufficiently small open subsets. In the local coordinates $(z, s) \in \mathbb{C}^{n-1} \times \mathbb{R}$, we may assume that,

$$L_j = \frac{\partial}{\partial \bar{z}_j} - i \frac{\phi_{\bar{z}_j}(z, \bar{z}, s)}{1 + i\phi_s(z, \bar{z}, s)} \frac{\partial}{\partial s}, \quad 1 \leq j \leq n - 1.$$

Since ϕ is real-analytic, we can complexify (resp. extend almost holomorphically, in the smooth case) in the s variable and write

$$M_j = \frac{\partial}{\partial \bar{z}_j} - i \frac{\phi_{\bar{z}_j}(z, \bar{z}, s + it)}{1 + i\phi_s(z, \bar{z}, s + it)} \frac{\partial}{\partial s}, \quad 1 \leq j \leq n - 1,$$

which are holomorphic in $s + it$ and extend the vector fields L_j .

Since ϕ and L_j are real-analytic now, equations (5.10), (5.11), (5.12) imply that there is real-analytic map $\Phi(W, \bar{W}, \Theta)$ defined in a neighborhood of $\{0\} \times \mathbb{C}^q$ in $\mathbb{C}^{n+1} \times \mathbb{C}^q$, polynomial in the last q variables for some integer q such that

$$\Phi(F, \bar{F}, (L^\alpha \bar{F})_{|\alpha|=1}) = 0$$

at $(z, s) \in U_0 \times V$. By (5.13) the matrix Φ_W is invertible at the central point 0, by the holomorphic version of the implicit function theorem. (In the

smooth category, we apply the “almost holomorphic” version of the implicit function theorem, cf. [L1]). We get a holomorphic map $\Psi = (\Psi_1, \dots, \Psi_{n+1})$ such that for (z, s) near the origin,

$$F_j = \Psi_j(\overline{F}, (L^\alpha \overline{F})_{|\alpha|=1}), \quad 1 \leq j \leq n + 1.$$

We now set for each $1 \leq j \leq n + 1$,

$$h_j(z, s, t) = \Psi_j(\overline{F}(z, s, -t), (M^\alpha \overline{F})_{|\alpha|=1}(z, s, -t)).$$

Since M is strongly pseudoconvex, the CR functions $F_j, 1 \leq j \leq n + 1$, all extend as holomorphic functions in $s + it$ to the side $t > 0$. Hence the conjugates $\overline{F}_j, 1 \leq j \leq n + 1$, extend holomorphically to the side $t < 0$. It now follows that $F_j, 1 \leq j \leq n + 1$, extend as holomorphic functions to a full neighborhood of the origin (See Lemma 9.2.9 in [BER]). (In the smooth category, we apply the edge-of-the-wedge theorem, as in [L1]). This establishes Proposition 5.3. \square

We then prove Theorem 2.

Proof of Theorem 2. We again write Ω for the open subset of M where F is smooth. Fix any $p_0 \in M$ and let $q_0 = F(p_0) \in M'$. We need to show that $p_0 \in \overline{\Omega}$ to establish the theorem. Assume that $p_0 = 0, q_0 = 0$. By assumption, M' is ℓ -nondegenerate at q_0 for some $\ell \geq 1$. We note that if $1 \leq \ell \leq 2$, it follows from Proposition 5.3 that $p_0 \in \overline{\Omega}$. We will prove $p_0 \in \overline{\Omega}$ for the general case $\ell \geq 3$ by induction on the order of nondegeneracy ℓ .

Suppose the statement $p_0 \in \overline{\Omega}$ holds when $\ell \leq k$ for some $k \geq 2$. We now consider the case $\ell = k + 1$, that is, we assume M' is $(k + 1)$ -nondegenerate at q_0 . Note that if there is a sequence $\{p_i\}_{i=1}^\infty \subset M$ converging to p_0 such that M' is at most k -nondegenerate at each $F(p_i)$, then it follows from the inductive hypothesis that each $p_i \in \overline{\Omega}$. Consequently, $p_0 \in \overline{\Omega}$ and hence the statement holds. Thus we are only left with the case that there exists a neighborhood U of p_0 on M such that $F(U) \subset V_k$. Here V_k is the set of k -degeneracy of M' near q_0 . More precisely,

$$V_k = \{q \in V : M' \text{ is } k\text{-degenerate at } q\},$$

for some small neighborhood $V = O \cap M'$ of q_0 . Here O is a small neighborhood of q_0 in \mathbb{C}^{n+1} . Since M' is $(k + 1)$ -nondegenerate at $q_0 = 0$, by Lemma

5.1, $\Delta_{i_0 i_1 \dots i_k}(0) \neq 0$, for some $1 \leq i_0 \leq i_1 \leq \dots \leq i_k \leq n$. We define

$$(5.14) \quad \varphi_k(W, \overline{W}) = \det \begin{pmatrix} \rho_W \\ \Lambda_1 \rho_W \\ \dots \\ \Lambda_{n-1} \rho_W \\ \Lambda_{i_1} \dots \Lambda_{i_k} \rho_W \end{pmatrix} (W, \overline{W}).$$

Set

$$\tilde{V}_k = \{W \in O : \varphi_k(W, \overline{W}) = 0\}.$$

We split our argument into two cases.

Case I: We first suppose that $i_0 \leq n-1$. Note that $F(U) \subset V_k \subset \tilde{V}_k$. We have

Claim 5.6. The \overline{w}_{i_0} -derivative $(\varphi_k)_{\overline{w}_{i_0}}$ of φ_k is nonzero at $q_0 = 0$.

Proof. It is equivalent to show that $\Lambda_{i_0} \varphi_k(0) \neq 0$. That is,

$$(5.15) \quad \Lambda_{i_0} \det \begin{pmatrix} \rho_W \\ \Lambda_1 \rho_W \\ \dots \\ \Lambda_{n-1} \rho_W \\ \Lambda_{i_1} \dots \Lambda_{i_k} \rho_W \end{pmatrix} (0) \neq 0$$

Note that

$$\begin{aligned} & \Lambda_{i_0} \det \begin{pmatrix} \rho_W \\ \Lambda_1 \rho_W \\ \dots \\ \Lambda_{n-1} \rho_W \\ \Lambda_{i_1} \dots \Lambda_{i_k} \rho_W \end{pmatrix} (0) \\ &= \det \begin{pmatrix} \Lambda_{i_0} \rho_W \\ \Lambda_1 \rho_W \\ \dots \\ \Lambda_{n-1} \rho_W \\ \Lambda_{i_1} \dots \Lambda_{i_k} \rho_W \end{pmatrix} (0) + \det \begin{pmatrix} \rho_W \\ \Lambda_{i_0} \Lambda_1 \rho_W \\ \dots \\ \Lambda_{n-1} \rho_W \\ \Lambda_{i_1} \dots \Lambda_{i_k} \rho_W \end{pmatrix} (0) + \dots \\ &+ \det \begin{pmatrix} \rho_W \\ \Lambda_1 \rho_W \\ \dots \\ \Lambda_{i_0} \Lambda_{n-1} \rho_W \\ \Lambda_{i_1} \dots \Lambda_{i_k} \rho_W \end{pmatrix} (0) + \det \begin{pmatrix} \rho_W \\ \Lambda_1 \rho_W \\ \dots \\ \Lambda_{n-1} \rho_W \\ \Lambda_{i_0} \Lambda_{i_1} \dots \Lambda_{i_k} \rho_W \end{pmatrix} (0) \end{aligned}$$

We claim that the first term up to the n^{th} term above are all zero. Indeed, otherwise, M' is at most k -nondegenerate at 0. This is a contradiction to our assumption.

We finally note the last term in the above equation just equals to $\Delta_{i_0 i_1 \dots i_k}(0)$, which is nonzero. This establishes the lemma. \square

Recall $F(U) \subset V_k \subset \tilde{V}_k$. We have

$$(5.16) \quad \varphi_k(F(Z), \overline{F(Z)}) \equiv 0, \text{ for all } Z \in U \subset M.$$

Applying L_{i_0} to the above equation and evaluating at $Z = 0$, we have,

$$(5.17) \quad L_{i_0} \varphi_k(F, \overline{F})|_0 = \sum_{i=1}^{n+1} (\varphi_k)_{\overline{w}_i}(F, \overline{F})|_0 L_{i_0} \overline{F}_i|_0 = 0.$$

Note that by our normalization, $L_{i_0} \overline{F}_i(0) = 0$, if $i \neq i_0$ and $L_{i_0} \overline{F}_{i_0}(0) \neq 0$. Moreover, by Claim 5.6, $(\varphi_k)_{\overline{w}_{i_0}}(0) \neq 0$. This is a contradiction to (5.17). Hence **Case I** cannot happen in this setting.

Case II: We are thus only left with the case if $i_0 = i_1 = \dots = i_k = n$. Let φ_k be as in (5.14) with $i_1 = \dots = i_k = n$. Again let $\tilde{V}_k = \{W \in O : \varphi_k(W, \overline{W}) = 0\}$.

By a similar argument as in the proof of Claim 5.6, we are able to prove the following lemma.

Claim 5.7. The \overline{w}_n -derivative $(\varphi_k)_{\overline{w}_n}$ of φ_k is nonzero at 0.

As a consequence of Lemma Claim 5.7, if we define $\tilde{\varphi}_k(W, \overline{W}) = \phi_k(W, \overline{W})$, then

$$(5.18) \quad (\tilde{\varphi}_k)_{w_n}(0) \neq 0.$$

Note $F(U) \subset V_k \subset \tilde{V}_k$. We have

$$\varphi_k(F(Z), \overline{F(Z)}) \equiv 0, \text{ for all } Z \in U.$$

Consequently,

$$(5.19) \quad \tilde{\varphi}_k(F(Z), \overline{F(Z)}) \equiv 0.$$

Recall that for all $Z \in U$,

$$(5.20) \quad \rho(F, \overline{F}) = 0,$$

$$(5.21) \quad L_i \rho(F, \overline{F}) = 0, 1 \leq i \leq n - 1.$$

Note that

$$(5.22) \quad \det \begin{pmatrix} \rho_W(F, \overline{F}) \\ L_1 \rho_W(F, \overline{F}) \\ \dots \\ L_{n-1} \rho_W(F, \overline{F}) \\ (\tilde{\varphi}_k)_W(F, \overline{F}) \end{pmatrix} (0) \neq 0$$

by equations (4.12), (4.13), (5.18). This implies that equations (5.19), (5.20), (5.21) forms a nondegenerate system for F . Then by a similar argument as in the proof of Proposition 5.3, it follows that F is smooth at 0. This proves $p_0 \in \overline{\Omega}$ when $\ell = k + 1$. Hence the statement holds for all $l \geq 1$ by mathematical induction. Theorem 2 is thus established. \square

6. Proof of Theorem 3

We are now going to prove Theorem 3. Fix $p_0 \in M$ and let $q_0 = F(p_0) \in M'$. We will show below that we can apply Theorem 2 for $q_0 \in M' \setminus X$ for some complex variety X in \mathbb{C}^{n+1} . We note that the transversality of F implies that the set $F^{-1}(M' \setminus X)$ is open and dense in M , and the statement of Theorem 3 follows. Indeed, to prove that last observation, suppose on the contrary that for a neighbourhood U of $p \in M$ we have that $F(M \cap U) \subset X$. Then $dF(\mathbb{C}T_p M) \subset \mathbb{C}T_{F(p)} X \subset T_{F(p)}^{(1,0)} M' + T_{F(p)}^{(0,1)} M'$, and so F is not transversal at p .

The following theorem gives the missing claim in the above argument. Let V be a small neighborhood of q_0 in \mathbb{C}^{n+1} . We first need to show that

Theorem 5. *M' is finitely nondegenerate near q_0 away from a complex analytic variety X in V .*

In order to do so, we shall first state and prove a useful general fact. For this, let $M \subset \mathbb{C}^N$ be a generic real-analytic submanifold of CR dimension n and real codimension d (i.e. $N = n + d$). We denote the set of germs at $p \in M$ of real-analytic functions on M with $\mathbb{C}\{M\}_p$. We say that an ideal $I \subset \mathbb{C}\{M\}_p$ is $\bar{\partial}_b$ -closed if for any CR vector field L on M and any $f \in I$ we have that $Lf \in I$. For any ideal $I \subset \mathbb{C}\{M\}_p$, we denote by $\mathcal{V}(I)$ the germ of the real-analytic subset of M given by the vanishing of all elements of I .

Proposition 6.1. *Let $I \subset \mathbb{C}\{M\}_p$ be an ideal which is $\bar{\partial}_b$ -closed. Then there exists a neighborhood U of p in \mathbb{C}^N and a complex subvariety $V \subset U$ such that, in the sense of germs at p , $V \cap M = \mathcal{V}(I)$.*

Proof. We choose normal coordinates $(z, w) \in \mathbb{C}^n \times \mathbb{C}^d$ for M at p ; in these coordinates, $p = 0$ and M is defined by

$$w = Q(z, \bar{z}, \bar{w}),$$

where $Q = (Q^1, \dots, Q^d)$ is a holomorphic map with values in \mathbb{C}^d , defined in a neighborhood of $(0, 0, 0) \in \mathbb{C}^n \times \mathbb{C}^n \times \mathbb{C}^d$, satisfying

$$(6.1) \quad Q(z, 0, \bar{w}) = Q(0, \bar{z}, \bar{w}) = \bar{w}, \quad Q(z, \bar{z}, \bar{Q}(\bar{z}, z, w)) = w.$$

A basis of the CR vector fields on M near 0 is given by

$$L_j = \frac{\partial}{\partial \bar{z}_j} + \sum_{k=1}^d \bar{Q}_{\bar{z}_j}^k(\bar{z}, z, w) \frac{\partial}{\partial \bar{w}_k}.$$

As usual, we use multiindex notation and for $\alpha = (\alpha_1, \dots, \alpha_n)$ we write $L^\alpha = L_1^{\alpha_1} \dots L_n^{\alpha_n}$.

Let $f \in \mathbb{C}\{M\}_p$. There exists a holomorphic function $F(z, w, \chi, \tau)$ defined in a neighborhood of $(0, 0, 0, 0) \in \mathbb{C}^n \times \mathbb{C}^d \times \mathbb{C}^n \times \mathbb{C}^d$ such that $f(z, w, \bar{z}, \bar{w}) = F(z, w, \bar{z}, \bar{Q}(\bar{z}, z, w))$ for $(z, w) \in M$. For any such f , we write $\varphi_f(z, w, \chi) = F(z, w, \bar{z}, \bar{Q}(\bar{z}, z, w))$. We note that $\frac{\partial^{|\alpha|} \varphi_f}{\partial \chi^\alpha}(z, w, \bar{z}) = \varphi_{L^\alpha f}(z, w, \bar{z})$. We also note that we can write

$$(6.2) \quad \begin{aligned} F(z, w, \chi, \bar{Q}(\chi, z, w)) &= \sum_{\alpha} \frac{1}{\alpha!} \frac{\partial^{|\alpha|}}{\partial \chi^\alpha} F(z, w, \chi, \bar{Q}(\chi, z, w)) \Big|_{\chi=0} \chi^\alpha \\ &= \sum_{\alpha} \frac{1}{\alpha!} L^\alpha F(z, w, 0, w) \chi^\alpha \\ &= \sum_{\alpha} \frac{1}{\alpha!} \frac{\partial^{|\alpha|} \varphi_f}{\partial \chi^\alpha}(z, w, 0) \chi^\alpha. \end{aligned}$$

So assume that we have chosen a small neighborhood of 0, such that inside this neighborhood, $\mathcal{V}(I)$ is defined by an ideal \tilde{I} of functions $f(z, w, \bar{z}, \bar{w})$ extending holomorphically to a common neighborhood of $(0, 0, 0, 0) \in \mathbb{C}^n \times \mathbb{C}^d \times \mathbb{C}^n \times \mathbb{C}^d$. We claim that $\mathcal{V}(\tilde{I}) = \{(z, w) : \varphi_f(z, w, 0) = 0, f \in \tilde{I}\}$.

Let $Z_0 = (z_0, w_0) \in \mathcal{V}(\tilde{I})$, and let $f \in \tilde{I}$. Then the holomorphic function $\chi \mapsto \varphi_f(z_0, w_0, \chi)$ vanishes to infinite order at $\chi = \bar{z}_0$; hence also

$\varphi_f(z_0, w_0, 0) = 0$. Assume now that $\varphi_g(z_0, w_0, 0) = 0$ for every $g \in \tilde{I}$. Then by (6.2), if $f \in \tilde{I}$ is arbitrary, then $f(z_0, w_0, \bar{z}_0, \bar{w}_0) = 0$. Hence, $\mathcal{V}(\tilde{I}) = \{(z, w) : \varphi_f(z, w, 0) = 0, f \in \tilde{I}\}$ as claimed. \square

The proof of Theorem 5 is a combination of Proposition 6.1 with the following fact.

Lemma 6.2. *Let $X \subset M$ be the set of points p in M at which M is not finitely nondegenerate of any order k . Then X can be defined, near every point $p \in M$, by an ideal which is $\bar{\partial}_b$ -closed.*

Proof. Let $p \in M$, and let $Z = (Z_1, \dots, Z_N)$ be coordinates near p . We note that M is k -nondegenerate if the space $E_k(p)$ has dimension N , where

$$E_0 = \Gamma(M, T^0M), \quad E_k = E_{k-1} + \{\mathcal{L}_L\omega : \omega \in E_{k-1}, L \text{ CR}\}.$$

Here $T^0M = \text{Re}(T^{(1,0)}M)^\perp \cap (T^{(0,1)}M)^\perp$ denotes the (real) characteristic bundle of M and \mathcal{L} the Lie derivative (of forms). It turns out that $E_k \subset \Gamma(M, T'M)$, where $T'M = (T^{(0,1)}M)^\perp$ is the bundle of holomorphic forms on M . We have that

$$T'M = \langle dZ_1, \dots, dZ_N \rangle.$$

We note that for

$$\omega = \sum_{j=1}^N \omega^j dZ_j,$$

it holds that

$$\mathcal{L}_L\omega = \sum_{j=1}^N (L\omega^j) dZ_j.$$

Choose a basis of characteristic forms $\theta_j = \sum_{k=1}^N \theta_j^k dZ_k$, where $j = 1, \dots, d$. The space E_k is therefore spanned by forms of the form

$$\mathcal{L}_{L^\alpha}\theta_j = \mathcal{L}_{L_1^{\alpha_1}} \cdots \mathcal{L}_{L_n^{\alpha_n}}\theta_j = \sum_{k=1}^N (L^\alpha\theta_j^k) dZ_k, \quad j = 1, \dots, d, \quad |\alpha| \leq k.$$

We therefore have that M is not ℓ -nondegenerate for some $\ell \leq k_0$ at p if and only if for every choice $r = (r_1, \dots, r_N)$ of integers $r_k \in \{1, \dots, d\}$ and for every choice of multiindices $A = (\alpha^1, \dots, \alpha^N)$, where $\alpha^j = (\alpha_1^j, \dots, \alpha_N^j)$

satisfies $|A| = \max\{|\alpha^j|: j = 1, \dots, N\} \leq k_0$, the determinant

$$D(A, r) = \det \begin{pmatrix} L^{\alpha^1} \theta_{r_1}^1 & \dots & L^{\alpha^1} \theta_{r_1}^N \\ \vdots & & \vdots \\ L^{\alpha^N} \theta_{r_N}^1 & \dots & L^{\alpha^N} \theta_{r_N}^N \end{pmatrix}$$

vanishes at p ; that is, if we denote by X_{k_0} the set of all points where p is not ℓ -nondegenerate where $\ell \leq k_0$, then X_{k_0} is defined by the ideal

$$I_{k_0} = (\{D(A, r): |A| \leq k_0\}).$$

Note that $LI_k \subset I_{k+1}$. The set $X = \bigcap_k X_k$ is now defined by $I = \bigcup_k I_k$, which is $\bar{\partial}_b$ -closed. □

By combining Proposition 6.1 and Lemma 6.2, we obtain the result in Theorem 5. Now the proof of Theorem 3 follows by combining Theorem 5 and the argument in the beginning of the section.

7. Proof of Theorem 4

Theorem 4 can be obtained very much in the same way as Theorem 3 from Theorem 2, noting that the statement of Theorem 3 is also valid in the real-analytic category. However, we are grateful to one of the anonymous referees, to point out another, conceptually very nice proof using the methods of the paper by Mir [Mi1], and using our normalization given in Proposition 3.3. We sketch that proof here, referring the reader for the details of the steps outlined in the cases below to [Mi1].

From Proposition 3.3, we have

$$(7.1) \quad \begin{aligned} \rho(F(Z), \overline{F(Z)}) &= -\text{Im}F_{n+1}(Z) + \sum_{j=1}^{n-1} |F_j(Z)|^2 \\ &+ \phi(F(Z), \overline{F(Z)}), \quad \phi = O(|F(Z)|^3). \end{aligned}$$

Applying a basis of CR vector fields L_1, \dots, L_{n-1} of M to that equation, and using the implicit function theorem in the resulting system of n equations, we obtain a holomorphic map Ψ , valued in \mathbb{C}^n , defined in a neighbourhood of $(0, L\bar{F}(0), 0) \in \mathbb{C}^{n+1} \times \mathbb{C}^{(n+1) \times (n-1)} \times \mathbb{C}$, such that with

$\tilde{F} = (F_1, \dots, F_{n-1}, F_{n+1})$ the following holds on M :

$$(7.2) \quad \tilde{F} = \Psi(\bar{F}, L\bar{F}, F_n) = (\Psi_1(\bar{F}, L\bar{F}, F_n), \dots, \Psi_n(\bar{F}, L\bar{F}, F_n)).$$

We now distinguish the following two cases:

Case 1. There exists a CR vector field X on M and a j_0 with $1 \leq j_0 \leq n$ such that $X\Psi_{j_0}(\bar{F}, L\bar{F}, T) \neq 0$ in a neighbourhood of $(0, 0) \in M \times \mathbb{C}_T$. In that case, one can apply the implicit function theorem to the system of equations formed by the equation $0 = X\Psi_{j_0}(\bar{F}, L\bar{F}, F_n)$ and the equations (7.2), at a generic point in M , to see that F is real-analytic there; hence we see that in Case 1, F is real-analytic on a dense, open subset of M .

Case 2. For all CR vector fields X on M and all $j, \leq j \leq n$, we have $X\Psi_j(\bar{F}, L\bar{F}, T) \equiv 0$. In that case, the reflection principle implies that $\Psi(\bar{F}(Z), L\bar{F}(Z), T) =: \Phi(Z, T)$ is holomorphic in a neighbourhood of $0 \in \mathbb{C}^n \times \mathbb{C}$. We write $\tilde{\Phi}(Z, T) = (\Phi_1(Z, T), \dots, \Phi_{N-1}(Z, T), T, \Phi_N(Z, T))$ and distinguish the following two alternatives:

Case 2A. $\varrho(\tilde{\Phi}(Z, T), \overline{\tilde{\Phi}(Z, T)}) \neq 0$ for $Z \in M$ and $T \in \mathbb{C}$: In that case, as in Case 1, it follows that actually F is real-analytic in a dense open subset of M .

Case 2B. $\varrho(\tilde{\Phi}(Z, T), \overline{\tilde{\Phi}(Z, T)}) \equiv 0$ for $Z \in M$ and $T \in \mathbb{C}$: By our normalization of F , we see that $\tilde{\Phi}$ is a *biholomorphism* from $M \times \mathbb{C}$ into M' . This is impossible if we assume M' to be holomorphically nondegenerate. Hence, Case 2B does not happen, and in any case, F is real-analytic on a dense open subset of M .

References

[BER] S. Baouendi, P. Ebenfelt, and L. Rothschild, *Real Submanifolds in Complex Space and Their Mappings*, Princeton University Press, Princeton, (1999).

[BHR] M. S. Baouendi, X. Huang, and L. P. Rothschild, *Regularity of CR-mappings between algebraic hypersurfaces*, *Invent. Math.* **125** (1996), 13–36.

[BK] V. K. Beloshapka and I. G. Kossovskiy, *The sphere in \mathbb{C}^2 as a model surface for degenerate hypersurfaces in \mathbb{C}^3* , *Russ. J. Math. Phys.* **22** (2015), 437–443.

[BX1] S. Berhanu and M. Xiao, *On the C^∞ version of reflection principle for CR-mappings between CR-manifolds*, *Amer. J. Math.* **137** (2015), 1365–1400.

- [BX2] S. Berhanu and M. Xiao, *On the regularity of CR-mappings between CR-manifolds of hypersurface type*, Trans. Amer. Math. Soc. **369** (2017), 6073–6086.
- [E1] P. Ebenfelt, *Normal forms and biholomorphic equivalence of real hypersurfaces in \mathbb{C}^3* , Indiana Univ. Math. J. **47** (1998), 311–366.
- [E2] P. Ebenfelt, *Uniformly Levi-degenerate CR-manifolds: the 5-dimensional case*, Duke Math. J. **110** (2001), 37–80. Correction in Duke Math. J. **131** (2006), 589–591.
- [EL] P. Ebenfelt and B. Lamel, *Finite jet determination of CR embeddings*, J. Geom. Anal. **14** (2004), 241–265.
- [Fa] J. Faran, *Maps from the two-ball to the three-ball*, Invent. Math. **68** (1982), 441–475.
- [Fe] C. Fefferman, *The Bergman kernel and biholomorphic mappings of pseudoconvex domains*, Invent. Math. **26** (1974), 1–65.
- [FK1] G. Fels and W. Kaup, *Classification of Levi-degenerate homogeneous CR-manifolds in dimension 5*, Acta Math. **201** (2008), 1–82.
- [FK2] G. Fels and W. Kaup, *CR-manifolds of dimension 5: a Lie algebra approach*, J. Reine Angew. Math. **604** (2007), 47–71.
- [Fr1] F. Forstnerič, *Extending proper holomorphic mappings of positive codimension*, Invent. Math. **95** (1989), 31–62.
- [Fr2] F. Forstnerič, *A Survey on Proper Holomorphic Mappings*, Proceeding of Year in SCVs at Mittag-Leffler Institute, Math. Notes **38**, Princeton, NJ: Princeton University Press, (1992).
- [Hu1] X. Huang, *On the mapping problem for algebraic real hypersurfaces in the complex spaces of different dimensions*, Ann. Inst. Fourier **44** (1994), 433–463.
- [Hu2] X. Huang, *Geometric analysis in several complex variables*, Ph.D. Thesis, Washington University in St. Louis (1994).
- [Hu3] X. Huang, *On a linearity problem for proper holomorphic maps between balls in complex spaces of different dimensions*, J. Differential Geom. **51** (1999), no. 1, 13–33.
- [IZ13] A. Isaev and D. Zaitsev, *Reduction of five-dimensional uniformly Levi-degenerate CR structures to absolute parallelisms*, J. Geom. Anal. **23** (2013), 1571–1605.

- [KaZa] W. Kaup and D. Zaitsev, *On local CR-transformations of Levi-degenerate group orbits in compact Hermitian symmetric spaces*, J. Eur. Math. Soc. **8** (2006), 465–490.
- [KiZa] S. Kim and D. Zaitsev, *Rigidity of CR-maps between Shilov boundaries of bounded symmetric domains*, Invent. Math. **193** (2013), 409–437.
- [KL1] I. Kossovskiy and B. Lamel, *On the analyticity of CR-diffeomorphisms*, Amer. J. Math. **140** (2018), no. 1, 139–188.
- [KL2] I. Kossovskiy and B. Lamel, *New extension phenomena for solutions of tangential Cauchy–Riemann equations*, Comm. Partial Differential Equations **41** (2016), 925–951.
- [L1] B. Lamel, *A C^∞ -regularity for nondegenerate CR-mappings*, Monatsh. Math. **142** (2004), 315–326.
- [L2] B. Lamel, *A reflection principle for real-analytic submanifolds of complex spaces*, J. Geom. Anal. **11** (2001), 625–631.
- [L3] B. Lamel, *Holomorphic maps of real submanifolds in complex spaces of different dimensions*, Pac. J. Math. **201** (2001), 357–387.
- [Le] H. Lewy, *On the boundary behavior of holomorphic mappings*, Acad. Naz. Lincei **3** (1977), 1–8.
- [MS] C. Medori and A. Spiro, *The equivalence problem for five-dimensional Levi-degenerate CR-manifolds*, Int. Math. Res. Not. IMRN. **2014** (2014), 5602–5647.
- [Mi1] N. Mir, *Holomorphic deformation of real-analytic CR-maps and analytic regularity of CR-mappings*, The Journal of Geometric Analysis **27** (2017), no. 3, 1920–1939.
- [Mo1] N. Mok, *Geometry of holomorphic isometries and related maps between bounded domains*, Geometry and Analysis. No. 2, pp. 225–270, Adv. Lect. Math. (ALM) **18**, Int. Press, Somerville, MA, (2011).
- [Mo2] N. Mok, *Nonexistence of proper holomorphic maps between certain classical bounded symmetric domains*, Chin. Ann. Math. Ser. B **29** (2008), 135–146.
- [Pi] S. I. Pinchuk, *On analytic continuation of biholomorphic mappings*, Mat. USSR Sb. **105** (1978), 574–594.

- [W] S. Webster, *On mapping an n -ball into an $(n + 1)$ -ball in complex space*, Pac. J. Math **81** (1979), 267–272.
- [XY] M. Xiao and Y. Yuan, *Holomorphic maps from the complex unit ball to type IV classical domains*, Journal de Mathématiques Pures et Appliquées **133** (2020), 139–166.

DEPARTMENT OF MATHEMATICS, MASARYK UNIVERSITY
611 37 BRNO, CZECH REPUBLIC

E-mail address: `ilya.kossovskiy@univie.ac.at`, `kossovskiyi@math.muni.cz`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF VIENNA
1090 WIEN, AUSTRIA

E-mail address: `bernhard.lamel@univie.ac.at`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA SAN DIEGO
LA JOLLA, CA 92093-0112, USA

E-mail address: `m3xiao@ucsd.edu`

RECEIVED MARCH 28, 2017

ACCEPTED APRIL 15, 2018

