

Algebraic virtual cycles for quantum singularity theories

HUAI-LIANG CHANG, YOUNG-HOON KIEM, AND JUN LI

We construct algebraic virtual cycles that give us the cohomological field theories of Fan-Jarvis-Ruan invariants by integral transformations.

1. Introduction

In this paper, we construct an algebraic virtual cycle that should provide us with the cohomological field theory of Fan-Jarvis-Ruan-Witten (FJRW for short) invariants in [7] by a Fourier-Mukai type integral transformation.

1.1. Background and motivation

Let $w : \mathbb{C}^N \rightarrow \mathbb{C}$ be a nondegenerate quasi-homogeneous polynomial (cf. §2.1) which defines a nonsingular hypersurface $Q_w = \mathbb{P}w^{-1}(0)$. Let \hat{G} be a subgroup of $(\mathbb{C}^*)^N$ and $\chi : \hat{G} \rightarrow \mathbb{C}^*$ be a homomorphism such that $w(g \cdot x) = \chi(g)w(x)$. The kernel of χ is a finite group denoted by G . If we let \hat{G} act on $\mathbb{C}^N \times \mathbb{C}$ by $g \cdot (x, t) = (g \cdot x, \chi(g)^{-1}t)$, the quotient stack

$$\mathfrak{X} = [(\mathbb{C}^N \times \mathbb{C})/\hat{G}]$$

admits a function $\mathbf{w}(x, t) = t \cdot w(x)$ and two GIT quotients

$$\mathfrak{X}_+ = ((\mathbb{C}^N - 0) \times \mathbb{C})/\hat{G}, \quad \mathfrak{X}_- = (\mathbb{C}^N \times (\mathbb{C} - 0))/\hat{G} = \mathbb{C}^N/G.$$

The former \mathfrak{X}_+ is an (orbi-)line bundle over the weighted projective stack \mathbb{P}^{N-1} and the critical locus of $\mathbf{w}|_{\mathfrak{X}_+}$ is Q_w , up to quotient by a finite group G/μ_d . On the other hand, $\mathbf{w}|_{\mathfrak{X}_-} = w$.

HLC was partially supported by Hong Kong GRF16301515 and GRF16301717; YHK was partially supported by Samsung Science and Technology Foundation SSTF-BA1601-01; JL was partially supported by NSF grants DMS-1564500 and DMS-1601211.

In [18], Witten conjectured that the Gromov-Witten invariants of Q_w should be computable by the Landau-Ginzburg (LG for short) model

$$w : \mathbb{C}^N/G \longrightarrow \mathbb{C}$$

whose curve counting invariants should be integrals on the solution space of Witten's equation on the moduli space X of G -spin curves (C, p_j, L_i, φ) (cf. §2.2) together with sections $(x_i) \in \prod_{i=1}^N H^0(L_i)$.

Through analysis, Fan, Jarvis and Ruan in [7] studied the solution space of Witten's equation and defined the FJRW invariants which were proved to satisfy nice properties like the splitting axioms, codified as cohomological field theories. Quantum singularity theories in the title refer to cohomological field theories arising from singularities like $w^{-1}(0)$.

Slightly later, Polishchuk and Vaintrob in [15, 16] provided a purely algebraic construction of cohomological field theories of the LG model $w : \mathbb{C}^N/G \rightarrow \mathbb{C}$ by matrix factorizations. They constructed a universal matrix factorization and their cohomological field theories are obtained by Fourier-Mukai type transformations on matrix factorizations and Hochschild homology. As the functors of matrix factorizations do not preserve the ordinary cohomology degrees, the algebraic theory in [15, 16] lacks in explicit interpretation in terms of cycles and basic properties like the homogeneity of dimension are not obvious.

An algebraic theory for FJRW invariants by algebraic cycles was provided in [5] for narrow sectors by constructing the virtual fundamental cycle for the moduli space X where Witten's equation is replaced by the cosection localization principle (cf. [12]). For the general case including broad sectors, the second and third named authors in [13] generalized the cosection localization of [12] to intersection homology and provided a direct construction of the cohomological field theories for both broad and narrow sectors. As the construction in [13] does not involve virtual cycles, one may wonder whether it is possible to construct the cohomological field theories by a Fourier-Mukai type integral transformation whose kernel is an algebraic virtual cycle.

The goal of this paper is to construct algebraic virtual cycles that give us the cohomological field theories of [13] by integral transformations.

1.2. Construction of virtual cycles by blowups

The moduli stack X of rigidified G -spin curves with sections can be written as the zero locus of a section s of a vector bundle E over a Deligne-Mumford stack Y (cf. §2.3). We also have a cosection $\sigma : E \rightarrow \mathcal{O}_Y$ satisfying $\sigma \circ s = 0$

and a smooth morphism $\mathbf{q} : Y \rightarrow Z = \underline{w}^{-1}(0)$, where \underline{w} is a nondegenerate quasi-homogeneous polynomial on a finite dimensional vector space. As X is usually not proper, the ordinary virtual fundamental class $[X]^{\text{vir}}$, as a Chow cycle supported in X , cannot be used for an integral transformation. On the other hand, the intersection S of X and the degeneracy locus \mathfrak{Z} (sometimes called the zero locus) of σ is the moduli space of rigidified G -spin curves; hence $S = X \cap \mathfrak{Z}$ is proper.

When Y is smooth, we have the canonical perfect obstruction theory $[T_Y|_X \xrightarrow{ds} E|_X]$ of X . Since $\sigma \circ s = 0$, $0 = d(\sigma \circ s) = \sigma \circ ds$ and hence the cosection σ descends to a cosection of the obstruction sheaf $Ob_X = \text{coker}(ds)$. In the narrow case, $Z = 0$ and one can apply the cosection localization principle in [12, Theorem 5.1] to obtain the cosection localized virtual cycle

$$[X]_{\text{loc}}^{\text{vir}} \in A_*(S)$$

which gives us the FJRW invariants in the narrow case (cf. [5]).

In the broad case, Y is singular and σ does not descend to Ob_X . In order to apply the cosection localization in [12], we will replace Z by its blowup Z' at the isolated singular point 0 , and pull back all the data above to Z' to get a smooth morphism $\mathbf{q}' : Y' \rightarrow Z'$, a vector bundle E' over Y' , a section s' of E' which defines X' , and a cosection σ' of E' whose zero locus \mathfrak{Z}' intersects with X' along the proper $S' = Z' \times_Z S$. Since Z' is smooth and $\sigma' \circ s' = 0$, Y' is smooth and the cosection σ' descends to a cosection of the obstruction sheaf $Ob_{X'} = \text{coker}(ds')$. Applying [12, Theorem 5.1], we obtain a cosection localized virtual cycle

$$[X']_{\text{loc}}^{\text{vir}} \in A_*(S').$$

The proper morphism p and the composite q below

$$p : S' \longrightarrow S, \quad q : S' \hookrightarrow Y' \xrightarrow{\mathbf{q}'} Z',$$

where the former is induced from the blowup morphism $Z' \rightarrow Z$, together with the virtual cycle $[X']_{\text{loc}}^{\text{vir}}$, give rise to an integral transformation

$$(1.1) \quad \Phi_{[X']_{\text{loc}}^{\text{vir}}} : H^*(Z') \longrightarrow H_*(S), \quad \alpha \mapsto p_*([X']_{\text{loc}}^{\text{vir}} \cap q^* \alpha).$$

The insertion space $\mathcal{H}^{\otimes n}$ of the FJRW theory (cf. §2.1) is contained in the direct sum of spaces of the form

$$IH_m(Z) \subset H^{m-2}(Z') \cong H_m(Z')$$

and we have a proper pushforward $H_*(S) \rightarrow H_*(\overline{M}_{g,n})$. Hence (1.1) enables us to define homomorphisms

$$(1.2) \quad \Omega'_{g,n} : \mathcal{H}^{\otimes n} \longrightarrow H_*(\overline{M}_{g,n}) \cong H^*(\overline{M}_{g,n}).$$

In [13, Theorem 3.2], the second and third named authors constructed the cosection localized Gysin maps for intersection homology

$$(1.3) \quad s_\sigma^! : IH_*(Y) \longrightarrow H_*(S).$$

Composed with the pullback $\mathbf{q}^* : IH_*(Z) \rightarrow IH_*(Y)$, (1.3) also enables us to define homomorphisms

$$(1.4) \quad \Omega_{g,n} : \mathcal{H}^{\otimes n} \longrightarrow H_*(\overline{M}_{g,n}) \cong H^*(\overline{M}_{g,n}).$$

In [13, Theorem 4.5], it was proved that (1.4) satisfies the axioms of the FJRW cohomological field theory.

The goal of this paper is to prove that

$$\Omega'_{g,n} = \Omega_{g,n},$$

and hence the integral transformation (1.1) by the algebraic virtual cycle $[X']_{\text{loc}}^{\text{vir}}$ provides us with the FJRW cohomological field theory (1.2) (cf. Theorem 5.1).

1.3. Layout

In §2, we recall the spin curves and their moduli space. In §3, we construct virtual cycles by blowup and define an integral transformation whose kernel is the virtual cycle. In §4, we recall the cohomological field theory construction by intersection homology in [13]. In §5, we prove the main theorem about the equality of the two cohomological field theories constructed in the previous sections.

1.4. Notation and convention

All varieties, schemes and stacks are defined over \mathbb{C} in this paper. We will use only the classical topology of algebraic varieties and schemes. All the topological spaces in this paper are locally compact Hausdorff countable CW complexes. Intersection homology in this paper refers to the middle perversity intersection homology unless stated otherwise. The Borel-Moore

homology groups are denoted by $H_*(-)$. We will not use the ordinary homology groups. All the cohomology groups in this paper have complex coefficients. The fundamental class of an irreducible closed substack V of a Deligne-Mumford stack Y in the Chow group $A_*(Y)$ is denoted by $[[V]]$ while the fundamental class of V in the Borel-Moore homology group $H_*(Y)$ is denoted by $[V]$.

2. Spin curves and FJRW invariants

In this section, we recall the Fan-Jarvis-Ruan-Witten theory from [7]. Our presentation follows [13, §4].

2.1. Hypersurface singularities

A polynomial $w : \mathbb{C}^N \rightarrow \mathbb{C}$ is quasi-homogeneous if for some $d_1, \dots, d_N, d \in \mathbb{Z}_{>0}$,

$$(2.1) \quad w(t^{d_1}x_1, \dots, t^{d_N}x_N) = t^d \cdot w(x_1, \dots, x_N).$$

Here we assume that $d > 0$ is the minimal possible. Let $q_i = d_i/d$. The quasi-homogeneous polynomial w is *nondegenerate* if the following are satisfied:

- 1) no monomial of w is of the form $x_i x_j$ for $i \neq j$;
- 2) the projective hypersurface Q_w defined by w is nonsingular:

$$Q_w = \mathbb{P}w^{-1}(0) \subset \mathbb{P}_{d_1, \dots, d_N}^{N-1} = (\mathbb{C}^N - 0)/\mathbb{C}^*$$

where \mathbb{C}^* acts on \mathbb{C}^N with weights d_1, \dots, d_N .

By (2), the hypersurface $w^{-1}(0) \subset \mathbb{C}^N$ has singularity only at the origin 0 and $q_i \leq \frac{1}{2}$.

We write $w = \sum_{k=1}^{\nu} c_k w_k$, where $c_k \in \mathbb{C}^*$ and w_k are distinct monomials. The kernel of the homomorphism

$$(2.2) \quad (w_1, \dots, w_{\nu}) : (\mathbb{C}^*)^N \longrightarrow (\mathbb{C}^*)^{\nu}$$

is the symmetry group

$$(2.3) \quad G_w = \{(\lambda_1 \cdots, \lambda_N) \in (\mathbb{C}^*)^N \mid w(\lambda_1 x_1, \dots, \lambda_N x_N) = w(x_1, \dots, x_N)\}$$

of w , which is finite by the nondegeneracy. Let

$$J_w = (e^{2\pi i q_1}, \dots, e^{2\pi i q_N}) \in G_w,$$

and fix a subgroup G of G_w containing J_w . The pair (w, G) is the input data for the FJRW theory in [7].

Consider the diagonal embedding $\mathbb{C}^* \rightarrow (\mathbb{C}^*)^\nu$ and the fiber product

$$\begin{array}{ccc} \hat{G}_w & \longrightarrow & \mathbb{C}^* \\ \downarrow & & \downarrow \\ (\mathbb{C}^*)^N & \longrightarrow & (\mathbb{C}^*)^\nu \end{array}$$

of (2.2). By the quasi-homogeneity (2.1), the homomorphism

$$\mathbb{C}^* \rightarrow (\mathbb{C}^*)^N, \quad t \mapsto (t^{d_1}, \dots, t^{d_N})$$

factors through \hat{G}_w , which together with the inclusion $G_w \rightarrow \hat{G}_w$ gives us a surjective homomorphism $G_w \times \mathbb{C}^* \rightarrow \hat{G}_w$ whose kernel is $\mu_d \leq \mathbb{C}^*$, the group of d -th roots of unity. The subgroup G of G_w thus determines

$$\hat{G} = G \times \mathbb{C}^* / \mu_d \subset \hat{G}_w$$

that fits into an exact sequence

$$(2.4) \quad 1 \longrightarrow G \longrightarrow \hat{G} \xrightarrow{\chi} \mathbb{C}^* \longrightarrow 1.$$

Since G acts trivially on w , for $\lambda \in \hat{G} \subset (\mathbb{C}^*)^N$,

$$(2.5) \quad w(\lambda \cdot x) = \chi(\lambda)w(x).$$

The *state space* for the singularity (w, G) in [7] is

$$(2.6) \quad \mathcal{H} = \bigoplus_{\gamma \in G} \mathcal{H}_\gamma, \quad \mathcal{H}_\gamma = H^{N_\gamma}(\mathbb{C}^{N_\gamma}, w_\gamma^\infty)^G.$$

Here \mathbb{C}^{N_γ} is the γ -fixed subspace of \mathbb{C}^N , and $w_\gamma^\infty = (\text{Re}(w_\gamma))^{-1}(a, \infty)$, where $w_\gamma = w|_{\mathbb{C}^{N_\gamma}}$ and $a \gg 0$.

If we let

$$Z'_\gamma \longrightarrow Z_\gamma = w_\gamma^{-1}(0)$$

be the weighted blowup at the origin, then Z'_γ is the line bundle $\mathcal{O}_{Q_{w_\gamma}}(-1)$ which is the restriction of $\mathcal{O}_{\mathbb{P}^{N_\gamma-1}}(-1)$ to $Q_{w_\gamma} = \mathbb{P}w_\gamma^{-1}(0) \subset \mathbb{P}^{N_\gamma-1}_{d_1, \dots, d_{N_\gamma}}$. As Q_{w_γ} is smooth by [7, Lemma 2.1.10], so is Z'_γ .

We consider the vanishing cohomology $H_{\text{van}}^*(\mathbb{P}w_\gamma^{-1}(0))$ and the primitive cohomology $H_{\text{prim}}^*(\mathbb{P}w_\gamma^{-1}(0))$ of $Q_{w_\gamma} = \mathbb{P}w_\gamma^{-1}(0)$. Complex Morse theory then provides us with isomorphisms

$$H^{N_\gamma}(\mathbb{C}^{N_\gamma}, w_\gamma^\infty) \cong H_{\text{van}}^{N_\gamma-2}(\mathbb{P}w_\gamma^{-1}(0)) \cong H_{\text{prim}}^{N_\gamma-2}(\mathbb{P}w_\gamma^{-1}(0)),$$

by [17, Proposition 2.27] since the weighted projective space has no primitive cohomology in non-zero degrees. On the other hand, the middle perversity intersection homology of $w_\gamma^{-1}(0)$ satisfies (cf. [3, p.20])

$$(2.7) \quad IH_i(w_\gamma^{-1}(0)) = \begin{cases} H_{\text{prim}}^{N_\gamma-2}(\mathbb{P}w_\gamma^{-1}(0)), & i = N_\gamma \\ \mathbb{C}, & i = 2N_\gamma - 2 \\ 0, & \text{otherwise.} \end{cases}$$

Hence we have

$$\begin{aligned} \mathcal{H}_\gamma &= IH_{N_\gamma}(w_\gamma^{-1}(0))^G \subset H_{\text{prim}}^{N_\gamma-2}(\mathbb{P}w_\gamma^{-1}(0)) \\ &\subset H^{N_\gamma-2}(\mathbb{P}w_\gamma^{-1}(0)) = H^{N_\gamma-2}(Z'_\gamma). \end{aligned}$$

For $\gamma_1, \gamma_2 \in G$, the Thom-Sebastiani sum

$$w_{\gamma_1} \boxplus w_{\gamma_2} : \mathbb{C}^{N_{\gamma_1}} \oplus \mathbb{C}^{N_{\gamma_2}} \rightarrow \mathbb{C}$$

is defined by $(x, y) \mapsto w_{\gamma_1}(x) + w_{\gamma_2}(y)$. By [14], we have canonical isomorphisms and a commutative square

$$\begin{array}{ccc} H_{\text{van}}^{N_{\gamma_1}-2}(\mathbb{P}w_{\gamma_1}^{-1}(0)) \otimes H_{\text{van}}^{N_{\gamma_2}-2}(\mathbb{P}w_{\gamma_2}^{-1}(0)) & \xrightarrow{\cong} & H_{\text{van}}^{N_{\gamma_1}+N_{\gamma_2}-2}(\mathbb{P}(w_{\gamma_1} \boxplus w_{\gamma_2})^{-1}(0)) \\ \downarrow \cong & & \downarrow \cong \\ IH_{N_{\gamma_1}}(w_{\gamma_1}^{-1}(0)) \otimes IH_{N_{\gamma_2}}(w_{\gamma_2}^{-1}(0)) & \xrightarrow{\cong} & IH_{N_{\gamma_1}+N_{\gamma_2}}((w_{\gamma_1} \boxplus w_{\gamma_2})^{-1}(0)). \end{array}$$

Therefore, $\mathcal{H}_{\gamma_1} \otimes \mathcal{H}_{\gamma_2}$ is canonically isomorphic to

$$(2.8) \quad \begin{aligned} &IH_{N_{\gamma_1}+N_{\gamma_2}}((w_{\gamma_1} \boxplus w_{\gamma_2})^{-1}(0))^{G \times G} \\ &\cong H_{\text{van}}^{N_{\gamma_1}+N_{\gamma_2}-2}(\mathbb{P}(w_{\gamma_1} \boxplus w_{\gamma_2})^{-1}(0))^{G \times G}. \end{aligned}$$

2.2. Moduli of spin curves

Given the input data (w, G) , we have the moduli stack of spin curves.

A pointed *twisted curve* refers to a proper Deligne-Mumford stack C with smooth substacks $p_1, \dots, p_n \subset C$, such that

- 1) denoting the coarse moduli space by $\rho : C \rightarrow |C|$, $|C|$ is a projective curve which has at worst nodal singularities and the markings $\rho(p_i) = |p_i|$ are smooth points of $|C|$;
- 2) ρ is an isomorphism away from special points (nodes or makings);
- 3) a marking is locally \mathbb{C}/μ_l for some $l > 0$, where μ_l is the group of l -th roots of unity;
- 4) a node is locally $\{xy = 0\}/\mu_l$ for some $l > 0$, where μ_l acts via $(x, y)^z = (zx, z^{-1}y)$.

The log dualizing sheaf of C is the pullback

$$\omega_C^{\log} = \rho^* \omega_{|C|}^{\log} = \rho^* \omega_{|C|}(|p_1| + \dots + |p_n|).$$

A G -spin curve is a principal \hat{G} -bundle P on a pointed twisted curve (C, p_1, \dots, p_n) equipped with an isomorphism

$$\varphi : \chi_* P \cong P(\omega_C^{\log})$$

of principal \mathbb{C}^* -bundles. Here $P(\omega_C^{\log})$ is the principal \mathbb{C}^* -bundle associated to the line bundle ω_C^{\log} ; χ is as in (2.4) and $\chi_* P$ is the principal \mathbb{C}^* -bundle obtained by applying χ to the fibers of P . Applying the inclusion map $\hat{G} \rightarrow (\mathbb{C}^*)^N$ to P , we obtain a principal $(\mathbb{C}^*)^N$ -bundle $P \times_{\hat{G}} (\mathbb{C}^*)^N$ over C which gives us line bundles (L_1, \dots, L_N) . The stabilizer group G_{p_j} of a marking p_j acts on the fiber $\oplus_i L_i|_{p_j}$ by $\gamma_j = (\gamma_{ij})_{1 \leq i \leq N} \in G$. We let $\underline{\gamma} = (\gamma_1, \dots, \gamma_n) \in G^n$ and call it the type of the G -spin curve. The stabilizer group G_p of a node p acts on $\oplus_i L_i|_p$ by a $\gamma_p \in G$.

The spin curve (C, p_j, L_i, φ) is *stable* if $(|C|, |p_1|, \dots, |p_n|)$ is a stable curve, and the homomorphism from the stabilizer group of a marking p_j (resp. a node p) into G that sends the generator to γ_j (resp. γ_p) is injective.

Theorem 2.1. *[7, Theorem 2.2.6] [16, Proposition 3.2.6] The stack $S_{g,n}$ of stable G -spin curves is a smooth proper Deligne-Mumford stack with projective coarse moduli. The forgetful morphism $S_{g,n} \rightarrow \overline{M}_{g,n}$ sending a G -spin*

curve (C, p_j, L_i, φ) to the underlying stable curve $(|C|, |p_j|)$ is flat proper and quasi-finite.

A rigidification of a G -spin curve at a marking p_j is an isomorphism

$$\psi_j : L_1|_{p_j} \oplus \cdots \oplus L_N|_{p_j} \xrightarrow{\cong} [\mathbb{C}^N / \langle \gamma_j \rangle],$$

where $\langle \gamma_j \rangle \leq G$ is the subgroup generated by γ_j , such that $w_k \circ \psi_j = \text{res}_{p_j} \circ \varphi_k|_{p_j}$ for every monomial w_k of w . Here $\varphi_k : L_1 \oplus \cdots \oplus L_N \rightarrow \omega_C^{\text{log}}$ is the homomorphism induced by w_k and φ . The moduli stack $S_{g,n}^{\text{rig}}$ of stable G -spin curves with rigidification is an étale cover over $S_{g,n}$ and hence $S_{g,n}^{\text{rig}}$ is a proper smooth Deligne-Mumford stack. The moduli stack of G -spin curves of type $\underline{\gamma}$ with rigidification is denoted by $S_{g,\underline{\gamma}}^{\text{rig}}$. So we have the disjoint union

$$S_{g,n}^{\text{rig}} = \sqcup_{\underline{\gamma}} S_{g,\underline{\gamma}}^{\text{rig}}.$$

2.3. Moduli of spin curves with sections

To simplify the notation, let $S = S_{g,\underline{\gamma}}^{\text{rig}}$. Let \mathcal{L}_i be the universal line bundle over the universal curve $\pi : \mathcal{C} \rightarrow S$ over S . By [16, §4.2], there are a locally free resolution

$$(2.9) \quad R\pi_*(\oplus_{i=1}^N \mathcal{L}_i) \cong [M \xrightarrow{\beta} F]$$

and a smooth morphism

$$\mathbf{q}_M : M \longrightarrow B = B_{\underline{\gamma}} = \prod_{j=1}^n \mathbb{C}^{N_{\gamma_j}}.$$

The Thom-Sebastiani sum $w_{\underline{\gamma}} = w_{\gamma_1} \boxplus \cdots \boxplus w_{\gamma_n}$ is a polynomial function on $B = B_{\underline{\gamma}}$ whose zero locus is denoted by $Z = Z_{\underline{\gamma}}$.

Let $E_M = \mathbf{p}_M^* F$ and s_M be the section of E_M defined by β where $\mathbf{p}_M : M \rightarrow S$ is the bundle projection. Then the zero locus $s_M^{-1}(0)$ of the section is the moduli space

$$X = X_{g,\underline{\gamma}}^{\text{rig}}$$

of stable G -spin curves (C, p_j, L_i, φ) of type $\underline{\gamma}$ together with rigidification ψ and sections $(x_1, \cdots, x_N) \in \oplus_i H^0(L_i)$ of the line bundles L_i . By [16, §4.2], E_M admits a cosection (i.e. a homomorphism to the structure sheaf) $\sigma_M :$

$E_M \rightarrow \mathcal{O}_M$, which satisfies

$$\sigma_M \circ s_M = w_\gamma \circ \mathbf{q}_M \quad \text{and} \quad X \cap \sigma_M^{-1}(0)_{\text{red}} = S.$$

Since the sum of residues of any meromorphic 1-form over a curve is zero,

$$(2.10) \quad X = X_{g,\gamma}^{\text{rig}} \subset Y = Y_{g,\gamma} := Z \times_B M \subset M.$$

In summary we have the following diagram:

$$(2.11) \quad \begin{array}{ccccccc} & & E_M & \xrightarrow{\sigma_M} & \mathcal{O}_M & \xlongequal{\quad} & M \times \mathbb{C} & \longrightarrow & \mathbb{C} \\ & & \downarrow & & \downarrow & & \downarrow & \nearrow & \\ X & \xlongequal{\quad} & s_M^{-1}(0) & \hookrightarrow & M & & & & \\ & & \downarrow \mathbf{p}_X & & \downarrow \mathbf{p}_M & & \downarrow \mathbf{q}_M & & \\ & & S & & B & \xrightarrow{w_\gamma} & \mathbb{C} & & \end{array}$$

By (2.10), we have a fiber diagram

$$(2.12) \quad \begin{array}{ccccc} X & \xrightarrow{i} & Y & \longrightarrow & M \\ & \searrow \mathbf{q}_X & \downarrow \mathbf{q}_Y & & \downarrow \mathbf{q}_M \\ & & Z & \hookrightarrow & B. \end{array}$$

Here \mathbf{q}_Y is smooth as \mathbf{q}_M is smooth. The restriction of E_M (resp. σ_M , resp. s_M) to Y is denoted by E (resp. σ , resp. s). By (2.10), $X = s^{-1}(0)$.

Because Z has at most an isolated hypersurface singularity by our assumption on the quasi-homogeneous polynomial w , for $\dim_{\mathbb{C}} Z = m - 1 \geq 2$, the affine variety Z is normal and hence Y is a normal as well. When $m \leq 2$, we may replace Z by its normalization. The intersection homology remains the same under normalization and all the arguments in this paper go through. Therefore for the FJRW theory, it suffices to work under the following.

Assumption 2.2. *Let Y be a normal Deligne-Mumford stack over \mathbb{C} . Let $s \in H^0(E)$ for a vector bundle E of rank r over Y and let $X = s^{-1}(0)$. Let $\sigma \in H^0(E^\vee) = \text{Hom}_Y(E, \mathcal{O}_Y)$ be a cosection of E satisfying*

$$(2.13) \quad \sigma \circ s = 0.$$

Let $\mathfrak{Z} = \sigma^{-1}(0) = \text{zero}(\sigma)$ be the degeneracy locus where σ is zero (i.e. not surjective). We assume that $S = X \cap \mathfrak{Z}$ is proper and there is a smooth

morphisms $\mathbf{q} = \mathbf{q}_Y : Y \rightarrow Z$ where $Z = w_\gamma^{-1}(0) \subset B = \mathbb{C}^m$ is the hypersurface defined by a nondegenerate quasi-homogeneous polynomial w_γ . Let $g : Z' \rightarrow Z$ be the blowup of Z at the origin, so that Z' is smooth. We let $f : Y' = Y \times_Z Z' \rightarrow Y$ denote the pullback of g by $\mathbf{q} : Y \rightarrow Z$.

More precisely, X (resp. \mathfrak{J}) is the closed substack defined by the image of $s^\vee : E^\vee \rightarrow \mathcal{O}_Y$ (resp. $\sigma : E \rightarrow \mathcal{O}_Y$) in \mathcal{O}_Y .

3. Algebraic virtual cycles

In this section, we construct algebraic virtual cycles for quantum singularity theories and define Fourier-Mukai type integral transformations which will give us cohomological field theories in the subsequent sections.

We use the notation in §2. Let E, Y, s, σ be as in Assumption 2.2. In particular, the moduli space $X = X_{g,\gamma}^{\text{rig}}$ of rigidified G -spin curves with sections

$$(C, p_j, L_i, \varphi, \psi, x_i),$$

where $(C, p_j, L_i, \varphi, \psi) \in S = S_{g,\gamma}^{\text{rig}}$ and $x_i \in H^0(L_i)$, is the zero locus of $s \in H^0(E)$ in Y . The restriction of the smooth morphism $\mathbf{q} = \mathbf{q}_Y$ to X is denoted by $\mathbf{q}_X : X \rightarrow Z = Z_\gamma$.

3.1. Cosection localized virtual cycle by blowup

By [1, §6], there is a relative perfect obstruction theory

$$(3.1) \quad \phi_{X/Z} : \mathbb{E}_{X/Z} = [E|_X^\vee \xrightarrow{ds} \Omega_{Y/Z}|_X] \longrightarrow \mathbb{L}_{X/Z}$$

where $\mathbb{L}_{X/Z} = \tau^{\geq -1} L_{X/Z}$ is the truncated relative cotangent complex of \mathbf{q}_X . In other words, $h^0(\phi_{X/Z})$ is an isomorphism and $h^{-1}(\phi_{X/Z})$ is surjective. Since $\sigma \circ s = w_\gamma \circ \mathbf{q} = 0$ (cf. (2.11)), $\sigma \circ ds|_{T_{Y/Z}} = 0$ and hence $\sigma : E \rightarrow \mathcal{O}_Y$ induces a cosection

$$(3.2) \quad \sigma_{X/Z} : \text{Ob}_{X/Z} = \text{coker}(T_{Y/Z}|_X \xrightarrow{ds} E|_X) \longrightarrow \mathcal{O}_X$$

of the relative obstruction sheaf. Let $X^\circ \subset X$ be the preimage of the smooth part $Z_{\text{sm}} \subset Z$. Then an easy argument shows that the cosection (3.2) descends to a cosection of the obstruction sheaf $\text{Ob}_X|_{X^\circ}$ of the induced absolute perfect obstruction theory of X° . The desired descent fails over $X - X^\circ$.

In order to obtain an absolute perfect obstruction theory with a cosection of its obstruction sheaf, we consider the blowup $g : Z' \rightarrow Z = Z_\gamma$ at the

origin and the fiber product

$$(3.3) \quad \begin{array}{ccccccc} S' & \xrightarrow{i'} & X' & \longrightarrow & Y' & \xrightarrow{q'} & Z' \\ p \downarrow & & \downarrow & & \downarrow f & & \downarrow g \\ S & \xrightarrow{i} & X & \longrightarrow & Y & \xrightarrow{q} & Z \end{array}$$

so that Y' (resp. Z') is a smooth model of Y (resp. Z). We denote the pullbacks of $E, s, \sigma, \mathfrak{Z}$ to Y' by $E', s', \sigma', \mathfrak{Z}'$ so that $X' = s'^{-1}(0)$ and $\mathfrak{Z}' = \sigma'^{-1}(0)$ while $S' = X' \cap \mathfrak{Z}'$.

The pullback of (3.1) to Y' is a relative perfect obstruction theory

$$(3.4) \quad \phi_{X'/Z'} : \mathbb{E}_{X'/Z'} = [E'|_{X'} \xrightarrow{ds'} \Omega_{Y'/Z'}|_{X'}] \longrightarrow \mathbb{L}_{X'/Z'}.$$

We also have the absolute perfect obstruction theory

$$(3.5) \quad \phi_{X'} : \mathbb{E}_{X'} = [E|_{X'} \xrightarrow{ds'} \Omega_{Y'}|_{X'}] \longrightarrow \mathbb{L}_{X'}.$$

As $\sigma' \circ s' = 0$ on Y' , σ' desends to a cosection

$$(3.6) \quad \sigma_{X'} : Ob_{X'} = \text{coker}(ds' : T_{Y'}|_{X'} \rightarrow E'|_{X'}) \longrightarrow \mathcal{O}_{X'}.$$

Therefore we can apply the cosection localization principle.

Theorem 3.1. [12, Theorem 5.1] *Let X' be a Deligne-Mumford stack equipped with a perfect obstruction theory $\phi_{X'}$ and a cosection $\sigma_{X'} : Ob_{X'} \rightarrow \mathcal{O}_{X'}$. Then X' admits a localized virtual cycle*

$$[[X']]_{\text{loc}}^{\text{vir}} \in A_*(S')$$

where S' is the zero locus of $\sigma_{X'}$. Its image in $A_*(X')$ by the inclusion $S' \subset X'$ is the ordinary virtual fundamental class $[[X']]^{\text{vir}}$ and $[[X']]_{\text{loc}}^{\text{vir}}$ is deformation invariant in the sense of intersection theory (cf. [1]).

Under Assumption 2.2, the construction of $[[X']]_{\text{loc}}^{\text{vir}}$ goes as follows: By [12, Proposition 4.3], the normal cone $C_{X'/Y'} \subset E'|_{X'}$ has support in

$$E'|_{X'}(\sigma_{X'}) = E'|_{S'} \cup \ker(\sigma_{X'} : E'|_{X'-S'} \rightarrow \mathcal{O}_{X'-S'}).$$

Then we apply the cosection localized Gysin map

$$(3.7) \quad 0^!_{E'|_{X'}, \sigma_{X'}} : A_*(E'|_{X'}(\sigma_{X'})) \longrightarrow A_*(S')$$

to the cycle $[[C_{X'/Y'}]]$ to obtain

$$(3.8) \quad [[X']]_{\text{loc}}^{\text{vir}} = 0_{E|_{X'}, \sigma_{X'}}^1 [[C_{X'/Y'}]] \in A_*(S').$$

Since $\dim C_{X'/Y'} = \dim Y = \dim S + \text{rank}(M) - 1$, the dimension of $[[X']]_{\text{loc}}^{\text{vir}}$ is

$$(3.9) \quad \dim S + \text{rank}(M) - 1 - \text{rank}(E) = 3g - 3 + n - 1 + \sum_i \chi(L_i).$$

This class $[[X']]_{\text{loc}}^{\text{vir}}$ depends only on the perfect obstruction theory $\phi_{X'}$ and the cosection $\sigma_{X'}$. In particular, the virtual cycle $[[X']]_{\text{loc}}^{\text{vir}} \in A_*(S')$ is independent of a choice of the resolution (2.9), and hence the choices of Y , E , etc in §2.3.

The construction of (3.7) in [12, §2] under Assumption 2.2 goes as follows: Let $\rho : \tilde{Y}' \rightarrow Y'$ be the blowup of Y' along $\tilde{\mathfrak{Z}}'$ (equivalently along the ideal $\sigma'(E')$) so that the pullback of σ' is a surjection $\tilde{E}' = \rho^* E' \rightarrow \mathcal{O}_{\tilde{Y}'}(-\tilde{\mathfrak{Z}}')$ where $\tilde{\mathfrak{Z}}'$ denotes the exceptional divisor of ρ . Restricting these to $\tilde{X}' = X' \times_{Y'} \tilde{Y}'$, we have a short exact sequence

$$0 \longrightarrow F' \longrightarrow \tilde{E}'|_{\tilde{X}'} \longrightarrow \mathcal{O}_{\tilde{X}'}(-\tilde{S}') \longrightarrow 0$$

of locally free sheaves where $\tilde{S}' = S' \times_{Y'} \tilde{Y}'$. For $\xi' \in A_*(E'|_{X'}(\sigma_{X'}))$, we pick $\zeta' \in A_*(F')$ and $\eta' \in A_*(E'|_{S'})$ such that

$$(3.10) \quad \xi' = \rho_* \zeta' + \iota'_* \eta', \quad \text{and} \quad \xi'|_{X'-S'} = \zeta'|_{\tilde{X}'-\tilde{S}'},$$

where ι'_* is the pushforward induced by the inclusion $S' \subset X'$. Then (3.7) is defined by

$$(3.11) \quad 0_{E'|_{X'}, \sigma_{X'}}^1(\xi') = -\rho_{S*} \left(\tilde{S}' \cdot 0_{F'}^1(\zeta') \right) + 0_{E'|_{S'}}^1(\eta') \in A_*(S')$$

where $\rho_S : \tilde{S}' \rightarrow S'$ is the restriction of ρ , $0_{F'}^1$ and $0_{E'|_{S'}}^1$ denote the ordinary Gysin maps and $\tilde{S}' \cdot$ denotes the intersection with the Cartier divisor \tilde{S}' (cf. [9]). By [12, §2], (3.11) is independent of all the choices. Moreover, instead of ρ , we may use any $\sigma_{X'}$ -regularizing morphism (Definition 4.2) and (3.7) is independent of this choice as well.

3.2. Integral transformations by virtual cycles

In this subsection, we define integral transformations by the virtual cycles constructed in Theorem 3.1. We will see in the subsequent section that these transformations form cohomological field theories.

Under Assumption 2.2, by Theorem 3.1, we have the virtual cycle $[[X']_{\text{loc}}^{\text{vir}} \in A_*(S')$ where $S' = S \times_X X'$. From (3.3), we have morphisms

$$(3.12) \quad Z' \xleftarrow{q} S' \xrightarrow{p} S$$

where p is obtained from g by base change and q is the restriction of \mathbf{q}' to S' . Since g is the blowup at the origin, p is proper. As $S = S_{g,\underline{\gamma}}^{\text{rig}}$ is proper, so is S' .

As the blowup Z' is a line bundle over $\mathbb{P}Z = \mathbb{P}w_{\underline{\gamma}}^{-1}(0)$, the cohomology $H^*(Z')$ of Z' is isomorphic to $H^*(\mathbb{P}w_{\underline{\gamma}}^{-1}(0)) \supset H_{\text{prim}}^*(\mathbb{P}w_{\underline{\gamma}}^{-1}(0))$. By the Thom-Sebastiani isomorphism (2.8), we have

$$(3.13) \quad \begin{aligned} \mathcal{H}_{\underline{\gamma}} &= \bigotimes_{j=1}^n \mathcal{H}_{\gamma_j} \cong H_{\text{prim}}^{\sum_j N_{\gamma_j} - 2}(\mathbb{P}w_{\underline{\gamma}}^{-1}(0))^{G^n} \\ &\subset H^{\sum_j N_{\gamma_j} - 2}(\mathbb{P}w_{\underline{\gamma}}^{-1}(0)) = H^{\sum_j N_{\gamma_j} - 2}(Z'). \end{aligned}$$

Since N_{γ_j} are often an odd number, a class in $\mathcal{H}_{\underline{\gamma}}$ is not algebraic in general. Hence for a Fourier-Mukai type integral transformation with our state space $\mathcal{H}_{\underline{\gamma}}$, we cannot use the Chow groups.

For an irreducible variety V , we can associate the Borel-Moore homology class of V after choosing a suitable triangulation. See [4, 11] for Borel-Moore homology. We thus have the cycle class map (cf. [9, Chapter 19])

$$(3.14) \quad h_{S'} : A_*(S') \longrightarrow H_*(S')$$

and the homological virtual cycle

$$(3.15) \quad [X']_{\text{loc}}^{\text{vir}} = h_{S'}[[X']_{\text{loc}}^{\text{vir}}] \in H_*(S').$$

By (3.12) and (3.15), we define our integral transformations as

$$(3.16) \quad \Phi_{[X']_{\text{loc}}^{\text{vir}}} : H^*(Z') \longrightarrow H^*(S), \quad \Phi_{[X']_{\text{loc}}^{\text{vir}}}(v) = p_*([X']_{\text{loc}}^{\text{vir}} \cap q^*(v)).$$

We have a forgetful morphism

$$\text{st} : S = S_{g,\underline{\gamma}}^{\text{rig}} \longrightarrow \overline{M}_{g,n}, \quad (C, p_j, L_i, \varphi, \psi) \mapsto (|C|, |p_j|)$$

whose pushforward is denoted by $st_* : H_*(S) \rightarrow H_*(\overline{M}_{g,n}) \cong H^*(\overline{M}_{g,n})$. Composing (3.16) and (3.13) with

$$(3.17) \quad \frac{(-1)^D}{\deg st} st_* : H_*(S) \longrightarrow H_*(\overline{M}_{g,n})$$

where $D = -\sum_i \chi(L_i)$, we obtain the composite

$$(3.18) \quad \Omega'_{g,n,\underline{\gamma}} : \mathcal{H}_{\underline{\gamma}} \longrightarrow H^*(Z') \longrightarrow H_*(S) \longrightarrow H_*(\overline{M}_{g,n}) \cong H^*(\overline{M}_{g,n}).$$

Summing up for $\underline{\gamma} \in G^n$, we obtain

Definition 3.2. For $g \geq 0$ and $n \geq 0$ with $2g - 2 + n > 0$, we have homomorphisms

$$(3.19) \quad \Omega'_{g,n} : \mathcal{H}^{\otimes n} = \bigoplus_{\underline{\gamma}} \mathcal{H}_{\underline{\gamma}} \longrightarrow H_*(\overline{M}_{g,n}) \cong H^*(\overline{M}_{g,n}).$$

By (3.9), the image of $\Omega'_{g,n}|_{\mathcal{H}_{\underline{\gamma}}}$ lies in degree

$$(3.20) \quad \begin{aligned} &2\left(3g - 3 + n - 1 + \sum_i \chi(L_i)\right) - \left(\sum_j N_j - 2\right) \\ &= 6g - 6 + 2n + 2 \sum_i \chi(L_i) - \sum_j N_{\gamma_j} \end{aligned}$$

which matches the computation in [7, Theorem 4.1.1].

In §5, we will see that the homomorphisms in Definition 3.2 form a cohomological field theory by comparing them with the cohomological field theory constructed in [13].

4. Quantum singularity theories via intersection homology

In this section, we recall the construction of cohomological field theories by intersection homology in [13].

We refer to [13, §2] for useful facts about Borel-Moore homology and intersection homology. For instance, we will use the natural map

$$\epsilon_Y : IH_i(Y) \longrightarrow H_i(Y)$$

that sends the middle perversity intersection homology cycles to itself in the Borel-Moore homology. Also, we will use proper pushforwards and placid (flat) pullbacks of Borel-Moore homology groups.

We first recall the cosection localized Gysin maps for intersection homology groups. Let E be a vector bundle of rank r over a Deligne-Mumford stack Y . Let s be a section and $X = s^{-1}(0)$. The canonical orientation on the fibers by the complex structure gives us the Thom class $\tau_{Y/E} \in H^{2r}(E, E - 0_E) = H_Y^{2r}(E)$ where 0_E denotes the zero section of E . The section s induces a map $(Y, X) \rightarrow (E, 0_E)$ and $e(E, s) = s^*\tau_{Y/E} \in H^{2r}(Y, Y - X) = H_X^{2r}(Y)$. The (ordinary) Gysin map is now defined as

$$s^! : H_i(Y) \longrightarrow H_{i-2r}(X), \quad \xi \mapsto \xi \cap e(E, s).$$

When E is equipped with a cosection $\sigma : E \rightarrow \mathcal{O}_Y$, $s^!$ further localizes to $S = X \cap \mathfrak{Z}$ where \mathfrak{Z} is the locus where σ is not surjective. Let $\iota : S \rightarrow X$ be the inclusion map. The following cosection localized Gysin map is the main machinery in this section.

Theorem 4.1. *[13, Theorem 3.2] Under Assumption 2.2, we have a homomorphism*

$$(4.1) \quad s_\sigma^! : IH_i(Y) \longrightarrow H_{i-2r}(S)$$

whose composition with $\iota_* : H_{i-2r}(S) \rightarrow H_{i-2r}(X)$ equals $s^! \circ \epsilon_Y$.

Here is an outline of the construction: For $s_\sigma^!$, we have to resolve the degeneracy of σ .

Definition 4.2. *Under Assumption 2.2, a proper morphism $\rho : \tilde{Y} \rightarrow Y$ is called σ -regularizing if it is an isomorphism over $Y - \mathfrak{Z}$ and the pullback $\tilde{E} = \rho^*E \rightarrow \mathcal{O}_{\tilde{Y}}$ of $\sigma : E \rightarrow \mathcal{O}_Y$ factors through a surjective homomorphism $\tilde{\sigma} : \tilde{E} \rightarrow \mathcal{O}_{\tilde{Y}}(-\tilde{\mathfrak{Z}})$ for an effective Cartier divisor $\tilde{\mathfrak{Z}}$ lying over \mathfrak{Z} .*

For instance, the blowup of Y along the ideal $I = \sigma(E) \subset \mathcal{O}_Y$ is σ -regularizing.

Let $\rho : \tilde{Y} \rightarrow Y$ be a σ -regularizing morphism and F be the kernel of the surjection $\tilde{\sigma} : \tilde{E} \rightarrow \mathcal{O}_{\tilde{Y}}(-\tilde{\mathfrak{Z}})$. By the decomposition theorem [2], for any $\xi \in H_i(Y)$, we can always find $\zeta \in H_i(\tilde{Y})$ and $\eta \in H_i(\mathfrak{Z})$ such that

$$\epsilon_Y(\xi) = \rho_*(\zeta) + j_*(\eta), \quad \epsilon_Y(\xi)|_{Y-\mathfrak{Z}} = \zeta|_{\tilde{Y}-\tilde{\mathfrak{Z}}}$$

where $j : \mathfrak{Z} \rightarrow Y$ denotes the inclusion (cf. [13, Lemma 2.2]). Then $s_\sigma^!$ is defined by

$$(4.2) \quad s_\sigma^!(\xi) = -\rho_{S*} \left(\zeta \cap e(F, \tilde{s}) \cap e(\mathcal{O}_{\tilde{X}}(\tilde{S}), t_{\tilde{S}}) \right) + \eta \cap e(E|_{\mathfrak{Z}}, s|_{\mathfrak{Z}})$$

where $t_{\tilde{S}}$ is the section of $\mathcal{O}_{\tilde{X}}(\tilde{S})$ whose zero locus is \tilde{S} . It was proved in [13, §3] that $s_{\sigma}^!(\xi)$ is independent of the choices of ζ, η and ρ .

In the FJRW theory, with $S = S_{g,\underline{\gamma}}^{\text{rig}}, X = X_{g,\underline{\gamma}}^{\text{rig}}, B_{\underline{\gamma}} = \prod_{j=1}^n \mathbb{C}^{N_{\gamma_j}}$ and $\underline{w} = w_{\underline{\gamma}}$, Assumption 2.2 is satisfied and hence we have the cosection localized Gysin map

$$(4.3) \quad s_{\sigma}^! : IH_i(Y) \longrightarrow H_{i-2\text{rank}(E)}(S)$$

by Theorem 4.1. Moreover since \mathbf{q}_Y is smooth, we have the pullback homomorphism

$$(4.4) \quad \mathbf{q}_Y^* : IH_i(Z) \longrightarrow IH_{i+\dim_{\mathbb{R}} M - \dim_{\mathbb{R}} B}(Y).$$

Composing (4.3) and (4.4) with the Thom-Sebastiani isomorphism

$$\mathcal{H}_{\underline{\gamma}} = \bigotimes_j \mathcal{H}_{\gamma_j} \cong IH_{\sum_j N_{\gamma_j}}(w_{\underline{\gamma}}^{-1}(0))^{G^n},$$

we obtain

$$(4.5) \quad \mathcal{H}_{\underline{\gamma}} \cong IH_{\sum_j N_{\gamma_j}}(w_{\underline{\gamma}}^{-1}(0))^{G^n} \xrightarrow{\mathbf{q}_Y^*} IH_*(Y) \xrightarrow{s_{\sigma}^!} H_{2\text{vd}(X) - \sum N_{\gamma_j}}(S_{g,\underline{\gamma}}^{\text{rig}})$$

where $\text{vd}(X) = 3g - 3 + n + \sum_i \chi(L_i)$ is the virtual dimension of X .

Composing (4.5) with (3.17) and summing over $\underline{\gamma}$, we obtain

$$(4.6) \quad \Omega_{g,n} : \mathcal{H}^{\otimes n} \longrightarrow H^*(\overline{M}_{g,n}).$$

Theorem 4.3. [13, Theorem 4.5] *The homomorphisms $\{\Omega_{g,n}\}_{2g-2+n>0}$ in (4.6) define a cohomological field theory with a unit for the state space \mathcal{H} . Moreover this cohomological field theory coincides with that in [7, Theorem 4.2.2] when we restrict \mathcal{H} to the narrow sector $\bigoplus_{\underline{\gamma}: N_{\gamma_j}=0, \forall j} \bigotimes_j \mathcal{H}_{\gamma_j}$.*

Here a cohomological field theory is a term that codifies nice properties expected from curve counting invariants as follows.

Definition 4.4. *Let \mathcal{H} be a vector space equipped with a basis $\{e_1, \dots, e_m\}$ and a perfect pairing $\langle e_k, e_l \rangle = c_{kl}$. A cohomological field theory with a unit $\mathbf{1}$ for the state space \mathcal{H} consists of homomorphisms*

$$(4.7) \quad \Omega_{g,n} : \mathcal{H}^{\otimes n} \longrightarrow H^*(\overline{M}_{g,n}), \quad \text{for } 2g - 2 + n > 0$$

satisfying the following:

- 1) if we let the symmetric group S_n act on $\overline{M}_{g,n}$ by permuting the markings and on $\mathcal{H}^{\otimes n}$ by permuting the factors, $\Omega_{g,n}$ is S_n -equivariant;
- 2) if we let $u : \overline{M}_{g-1,n+2} \rightarrow \overline{M}_{g,n}$ denote the gluing of the last two markings, then we have

$$(4.8) \quad u^* \Omega_{g,n}(v_1, \dots, v_n) = \sum_{k,l} c^{kl} \Omega_{g-1,n+2}(v_1, \dots, v_n, e_k, e_l)$$

in $H^*(\overline{M}_{g-1,n+2})$ for all $v_i \in \mathcal{H}$ where $(c^{kl}) = (c_{kl})^{-1}$;

- 3) if we let $u : \overline{M}_{g_1,n_1+1} \times \overline{M}_{g_2,n_2+1} \rightarrow \overline{M}_{g,n}$ with $g = g_1 + g_2$ and $n = n_1 + n_2$ denote the gluing of the last markings, then

$$(4.9) \quad u^* \Omega_{g,n}(v_1, \dots, v_n) = \sum_{k,l} c^{kl} \Omega_{g_1,n_1+1}(v_1, \dots, v_{n_1}, e_k) \otimes \Omega_{g_2,n_2+1}(v_{n_1+1}, \dots, v_n, e_l)$$

in $H^*(\overline{M}_{g_1,n_1+1}) \otimes H^*(\overline{M}_{g_2,n_2+1})$ for all $v_i \in \mathcal{H}$;

- 4) if we let $\theta : \overline{M}_{g,n+1} \rightarrow \overline{M}_{g,n}$ denote the morphism forgetting the last marking, we have

$$(4.10) \quad \Omega_{g,n+1}(v_1, \dots, v_n, \mathbf{1}) = \theta^* \Omega_{g,n}(v_1, \dots, v_n), \quad \forall v_i \in \mathcal{H};$$

- 5) $\Omega_{0,3}(v_1, v_2, \mathbf{1}) = \langle v_1, v_2 \rangle$ for $v_i \in \mathcal{H}$.

For a smooth projective variety Q , letting $\overline{M}_{g,n}(Q, d)$ denote the moduli stack of stable maps to Q of genus g and degree d , it is well known that

$$H^*(Q)^{\otimes n} \longrightarrow H^*(\overline{M}_{g,n}), \quad (v_j) \mapsto p_* \left([\overline{M}_{g,n}(Q, d)]^{\text{vir}} \cap \prod_{j=1}^n ev_j^*(v_j) \right)$$

form a cohomological field theory where $p : \overline{M}_{g,n}(Q, d) \rightarrow \overline{M}_{g,n}$ is the forgetful morphism and $ev_j : \overline{M}_{g,n}(Q, d) \rightarrow Q$ is the evaluation map at the j -th marking.

The homomorphisms (4.6) are suitable for proving the axioms of cohomological field theories while those in Definition 3.2 are defined by Fourier-Mukai type integral operators with algebraic kernels $[[X']_{\text{loc}}^{\text{vir}}$. In the subsequent section, we will prove that actually they are the same.

We end this section with the following.

Proposition 4.5. *Let E be an algebraic vector bundle of rank r over a smooth Deligne-Mumford stack Y . Let $s \in H^0(E)$ and $\sigma \in H^0(E^\vee)$ satisfy $\sigma \circ s = 0$. Let $X = \text{zero}(s)$, $\mathfrak{Z} = \text{zero}(\sigma)$ and $S = X \cap \mathfrak{Z}$. Then*

$$s_\sigma^! [Y] = [X]_{\text{loc}}^{\text{vir}} := h_S \llbracket X \rrbracket_{\text{loc}}^{\text{vir}} \in H_{2 \dim Y - 2r}(S),$$

where $s_\sigma^!$ is from Theorem 4.1 and $\llbracket X \rrbracket_{\text{loc}}^{\text{vir}}$ is from Theorem 3.1 (with primes removed).

Proof. Suppose $\sigma = 0$. Let $\Gamma \subset E \times \mathbb{C}^*$ be the graph of the section $(y, t) \mapsto t^{-1}s(y)$. Let $\bar{\Gamma}$ be the closure of Γ in $E \times \mathbb{C}$. Then for $t \neq 0$, the fiber $\bar{\Gamma}_t$ over t is isomorphic to Y and the fiber $\bar{\Gamma}_0$ over $t = 0$ is the normal cone $C_{X/Y}$ (cf. [9]). As $Y = \bar{\Gamma}_1$ is homologous to $\bar{\Gamma}_0 = C_{X/Y}$, $s^! [Y] = s^! [C_{X/Y}]$. The proposition now follows from the fact that $s^! [C_{X/Y}] = h_X \circ 0_E^! \llbracket C_{X/Y} \rrbracket$ by [9, Chapter 19] where $0_E^!$ denotes the algebraic Gysin map of Chow groups.

When σ is not necessarily zero, we let $\rho : \tilde{Y} \rightarrow Y$ be the blowup of Y along \mathfrak{Z} so that we have an exact sequence

$$0 \longrightarrow F \longrightarrow \tilde{E} \xrightarrow{\tilde{\sigma}} \mathcal{O}_{\tilde{Y}}(-\tilde{\mathfrak{Z}}) \longrightarrow 0$$

where $\tilde{\mathfrak{Z}}$ is the exceptional divisor, \tilde{E} , \tilde{s} and $\tilde{\sigma}$ denote the pullbacks of E , s and σ to \tilde{Y} respectively. Then

$$\llbracket Y \rrbracket = \rho_* \llbracket \tilde{Y} \rrbracket.$$

From the commutative diagram

$$\begin{array}{ccc} \tilde{Y} \times \mathbb{C}^* & \xrightarrow{t^{-1}\tilde{s}} & \tilde{E} \times \mathbb{C}^* \\ \downarrow & & \downarrow \\ Y \times \mathbb{C}^* & \xrightarrow{t^{-1}s} & E \times \mathbb{C}^*, \end{array}$$

we find that $\llbracket C_{X/Y} \rrbracket = \hat{\rho}_* \llbracket C_{\tilde{X}/\tilde{Y}} \rrbracket$ as $C_{X/Y}$ (resp. $C_{\tilde{X}/\tilde{Y}}$) is rationally equivalent to Y in E (resp. \tilde{Y} in \tilde{E}). Here $\hat{\rho} : \tilde{E}|_{\tilde{X}} \rightarrow E|_X$ is the induced map from ρ . By (3.11),

$$\llbracket X \rrbracket_{\text{loc}}^{\text{vir}} = 0_{E|_X, \sigma}^! \llbracket C_{X/Y} \rrbracket = -\rho_{S*}(\tilde{S} \cdot 0_{F|\tilde{X}}^! \llbracket C_{\tilde{X}/\tilde{Y}} \rrbracket).$$

By [9, Chapter 19], (4.2) and the case for trivial cosection, if we apply the cycle class map h_S , we obtain

$$h_S \llbracket X \rrbracket_{\text{loc}}^{\text{vir}} = -\rho_{S*}(t_{\tilde{S}}^! \tilde{s}^! \llbracket \tilde{Y} \rrbracket) = s_\sigma^! [Y]$$

as desired. □

A direct consequence of Proposition 4.5 is the following.

Corollary 4.6. *Under Assumption 2.2, letting $Z' \rightarrow Z = Z_{\underline{\gamma}}$ be the blowup at the origin and using the notation of §3, we have*

$$(4.11) \quad [X']_{\text{loc}}^{\text{vir}} = s'^!_{\sigma'}[Y'], \quad \text{and} \quad \Phi_{[X']_{\text{loc}}^{\text{vir}}}(\alpha) = p_*(s'^!_{\sigma'}[Y'] \cap q^*(\alpha))$$

for $\alpha \in H^*(Z')$.

5. Comparison

The goal of this section is to prove the following.

Theorem 5.1. *The homomorphism $\Omega_{g,n}$ in (4.6) equals $\Omega'_{g,n}$ in (3.19). Hence, the integral transformations $\Phi_{[X']_{\text{loc}}^{\text{vir}}}$ with algebraic kernels $[X']_{\text{loc}}^{\text{vir}}$ give rise to cohomological field theories $\{\Omega'_{g,n}\}$.*

Proof. Recall that the maps (4.6) (resp. (3.19)) are obtained by composing $s^!_{\sigma} \mathbf{q}^*$ (resp. $\Phi_{[X']_{\text{loc}}^{\text{vir}}}$) with the stabilization (3.17). Therefore the theorem follows once we show that whenever

$$(5.1) \quad \epsilon_Z(v) = g_*([Z'] \cap \alpha) \quad \text{for } \alpha \in H^*(Z'), v \in IH_*(Z),$$

$$(5.2) \quad \Phi_{[X']_{\text{loc}}^{\text{vir}}}(\alpha) = p_*([X']_{\text{loc}}^{\text{vir}} \cap q^* \alpha) = s^!_{\sigma} \mathbf{q}^*(v),$$

which is Theorem 5.4 below. Note that since $Z = w_{\underline{\gamma}}^{-1}(0)$ is a nondegenerate hypersurface singularity, if we let $m - 1 = \dim Z$, $IH_m(Z) \cong H_{\text{prim}}^{m-2}(\mathbb{P}Z) \subset H^{m-2}(\mathbb{P}Z) \cong H^{m-2}(Z') \cong H_m(Z')$ and we have a commutative diagram

$$(5.3) \quad \begin{array}{ccccc} H_m(Z') & \xleftarrow[\cong]{[Z'] \cap} & H^{m-2}(Z') & \xleftarrow{\cong} & H^{m-2}(\mathbb{P}Z) \\ g_* \downarrow & & & & \uparrow \\ H_m(Z) & \xleftarrow{\epsilon_Z} & IH_m(Z) & \xrightarrow{\cong} & H_{\text{prim}}^{m-2}(\mathbb{P}Z). \end{array}$$

By the isomorphism $IH_m(Z) \cong H_{\text{prim}}^{m-2}(\mathbb{P}Z)$, $\alpha \in H_{\text{prim}}^{m-2}(\mathbb{P}Z)$ and $v \in IH_m(Z)$ determine each other uniquely. □

For a proof of (5.2), we need a couple of propositions on the cosection localized Gysin map $s_\sigma^!$.

By [10, 5.2], for a Deligne-Mumford stack Y , we have a cap product

$$(5.4) \quad IH_i(Y) \times H^j(Y) \xrightarrow{\cap} IH_{i-j}(Y), \quad (\xi, \alpha) \mapsto \xi \cap \alpha$$

which fits into a commutative diagram

$$(5.5) \quad \begin{array}{ccc} IH_i(Y) \times H^j(Y) & \xrightarrow{\cap} & IH_{i-j}(Y) \\ \epsilon_Y \times 1 \downarrow & & \downarrow \epsilon_Y \\ H_i(Y) \times H^j(Y) & \xrightarrow{\cap} & H_{i-j}(Y) \end{array}$$

where the bottom arrow is the usual cap product [11, IX.3]. The cap product satisfies the projection formula (cf. [11, IX.3.7])

$$(5.6) \quad g_*(\xi \cap f^*\alpha) = f_*\xi \cap \alpha, \quad \xi \in H_i(X), \alpha \in H_W^j(Y) = H^j(Y, Y - W)$$

where $f : X \rightarrow Y$ is proper and $g : f^{-1}(W) \rightarrow W$ is the restriction of f to $f^{-1}(W) = W \times_Y X$ for closed $W \subset Y$. For closed A and B in Y ,

$$(5.7) \quad (\xi \cap \alpha) \cap \beta|_A = \xi \cap (\alpha \cup \beta) = (\xi \cap \beta) \cap \alpha|_B$$

for $\xi \in H_i(Y)$, $\alpha \in H_A^j(Y)$ and $\beta \in H_B^k(Y)$ when jk is even.

Proposition 5.2. *Let $X = \text{zero}(s)$, $\mathfrak{Z} = \text{zero}(\sigma)$ and $S = X \cap \mathfrak{Z}$. For $\alpha \in H^j(Y)$ and $\xi \in IH_i(Y)$, we have*

$$s_\sigma^!(\xi \cap \alpha) = s_\sigma^!(\xi) \cap \alpha|_S \in H_{i-j-2r}(S).$$

Proof. Let $\rho : \tilde{Y} \rightarrow Y$ be a σ -regularizing birational morphism such that the exceptional divisor $\tilde{\mathfrak{Z}}$ lies over $\mathfrak{Z} = \text{zero}(\sigma)$. The cosection $\sigma : E \rightarrow \mathcal{O}_Y$ lifts to a surjective homomorphism $\tilde{\sigma} : \tilde{E} = \rho^*E \rightarrow \mathcal{O}_{\tilde{Y}}(-\tilde{\mathfrak{Z}})$ whose kernel is denoted by F . For $s_\sigma^!(\xi)$, we pick $\zeta \in H_i(\tilde{Y})$ and $\eta \in H_i(\mathfrak{Z})$ such that

$$\epsilon_Y(\xi) = \rho_*\zeta + j_*\eta, \quad \epsilon_Y(\xi)|_{Y-\mathfrak{Z}} = \zeta|_{\tilde{Y}-\tilde{\mathfrak{Z}}}$$

where $j : \mathfrak{Z} \rightarrow Y$ denotes the inclusion. By the definition of $s_\sigma^!$, we have

$$(5.8) \quad s_\sigma^!(\xi) = \eta \cap e(E|_{\mathfrak{Z}}, s|_{\mathfrak{Z}}) - \rho_{S*}(\zeta \cap e(F, \tilde{s}) \cap e(\mathcal{O}_{\tilde{X}}(\tilde{S}), t_{\tilde{S}}))$$

where $\rho_S : \tilde{S} \rightarrow S$ is the restriction of ρ to $\tilde{S} = \tilde{Y} \times_Y S$ and $t_{\tilde{S}}$ is the section of $\mathcal{O}_{\tilde{X}}(\tilde{S})$ whose vanishing locus is the divisor \tilde{S} .

By (5.5) and (5.6), we have

$$\begin{aligned} \epsilon_Y(\xi \cap \alpha) &= \epsilon_Y(\xi) \cap \alpha = \rho_*(\zeta) \cap \alpha + j_*(\eta) \cap \alpha \\ &= \rho_*(\zeta \cap \rho^* \alpha) + j_*(\eta \cap j^* \alpha). \end{aligned}$$

Hence we have

$$\begin{aligned} s_\sigma^!(\xi \cap \alpha) &= (\eta \cap j^* \alpha) \cap e(E|_{\mathfrak{Z}}, s|_{\mathfrak{Z}}) \\ &\quad - \rho_{S*}((\zeta \cap \rho^* \alpha) \cap e(F, \tilde{s}) \cap e(\mathcal{O}_{\tilde{X}}(\tilde{S}), t_{\tilde{S}})). \end{aligned}$$

By (5.7), the above line equals

$$(5.9) \quad (\eta \cap e(E|_{\mathfrak{Z}}, s|_{\mathfrak{Z}})) \cap \alpha|_S - \rho_{S*}(\zeta \cap e(F, \tilde{s}) \cap e(\mathcal{O}_{\tilde{X}}(\tilde{S}), t_{\tilde{S}}) \cap \rho_{S^*}^* \alpha|_S).$$

By the projection formula again and (5.8), (5.9) equals

$$\begin{aligned} &(\eta \cap e(E|_{\mathfrak{Z}}, s|_{\mathfrak{Z}})) \cap \alpha|_S - \rho_{S*}(\zeta \cap e(F, \tilde{s}) \cap e(\mathcal{O}_{\tilde{X}}(\tilde{S}), t_{\tilde{S}})) \cap \alpha|_S \\ &= s_\sigma^!(\xi) \cap \alpha|_S. \end{aligned}$$

This proves the proposition. □

Proposition 5.3. *Let $f : Y' \rightarrow Y$ be a proper morphism of normal Deligne-Mumford stacks. Let $E', s', \sigma', X', \mathfrak{Z}', S'$ etc be the pullbacks of $E, s, \sigma, X, \mathfrak{Z}, S$ etc by f . Let $p : S' \rightarrow S$ denote the restriction of f to S' . Let $\xi \in IH_i(Y)$. Suppose there exists $\xi' \in IH_i(Y')$ such that $f_* \epsilon_{Y'}(\xi') = \epsilon_Y(\xi)$. Then we have*

$$p_* s'^!_{\sigma'}(\xi') = s^!_{\sigma}(\xi) \in H_{i-2r}(S).$$

Proof. Let $\rho : \tilde{Y} \rightarrow Y$ be a σ -regularizing birational morphism so that σ lifts to the surjective homomorphism $\tilde{\sigma} : \tilde{E} = \rho^* E \rightarrow \mathcal{O}_{\tilde{Y}}(-\tilde{\mathfrak{Z}})$ over \tilde{Y} with kernel F . Consider the fiber product

$$\begin{array}{ccc} \tilde{Y}' & \xrightarrow{\tilde{f}} & \tilde{Y} \\ \rho' \downarrow & & \downarrow \rho \\ Y' & \xrightarrow{f} & Y \end{array}$$

Then $\tilde{\sigma}$ lifts to a surjective homomorphism $\tilde{\sigma}' : \tilde{E}' = \rho'^* E' \rightarrow \mathcal{O}_{\tilde{Y}'}(-\tilde{\mathfrak{Z}}')$ where $\tilde{\mathfrak{Z}}' = \tilde{\mathfrak{Z}} \times_{\tilde{Y}} \tilde{Y}'$.

For $s'_{\sigma'}(\xi')$, we pick $\zeta' \in H_i(\tilde{Y}')$ and $\eta' \in H_i(\mathfrak{Z}')$ such that

$$(5.10) \quad \epsilon_{Y'}(\xi') = \rho'_* \zeta' + j'_* \eta', \quad \epsilon_{Y'}(\xi')|_{Y'-\mathfrak{Z}'} = \zeta'|_{\tilde{Y}'-\tilde{\mathfrak{Z}'}}$$

where $j' : \mathfrak{Z}' \rightarrow Y'$ denotes the inclusion. By the definition of $s'_{\sigma'}$, we have

$$(5.11) \quad s'_{\sigma'}(\xi') = \eta' \cap e(E'|_{\mathfrak{Z}'}, s'|_{\mathfrak{Z}'}) - \rho_{S' *} (\zeta' \cap e(F', \tilde{s}') \cap e(\mathcal{O}_{\tilde{X}'}(\tilde{S}'), t_{\tilde{S}'}))$$

By applying f_* to (5.10), we have

$$\epsilon_Y(\xi) = f_* \epsilon_{Y'}(\xi') = f_* \rho'_* \zeta' + f_* j'_* \eta' = \rho_*(\tilde{f}_*(\zeta')) + j_*(f_{\mathfrak{Z}*}(\eta'))$$

where $f_{\mathfrak{Z}} : \mathfrak{Z}' \rightarrow \mathfrak{Z}$ denotes the restriction of f to $\mathfrak{Z}' = \mathfrak{Z} \times_Y Y'$. Moreover, $\tilde{f}_*(\zeta')|_{\tilde{Y}-\tilde{\mathfrak{Z}}} = \tilde{f}_* \epsilon_{Y'}(\xi')|_{Y-\mathfrak{Z}} = \epsilon_Y(\xi)|_{Y-\mathfrak{Z}}$ by (5.10). Hence, we have

$$(5.12) \quad s_{\sigma}(\xi) = f_{\mathfrak{Z}*}(\eta') \cap e(E|_{\mathfrak{Z}}, s|_{\mathfrak{Z}}) - \rho_{S*}(\tilde{f}_*(\zeta') \cap e(F, \tilde{s}) \cap e(\mathcal{O}_{\tilde{X}}(\tilde{S}), t_{\tilde{S}})).$$

By the projection formula (5.6), $f_{\mathfrak{Z}*}(\eta') \cap e(E|_{\mathfrak{Z}}, s|_{\mathfrak{Z}}) = p_*(\eta' \cap e(E'|_{\mathfrak{Z}'}, s'|_{\mathfrak{Z}'}))$ and

$$\begin{aligned} & \rho_{S*}(\tilde{f}_*(\zeta') \cap e(F, \tilde{s}) \cap e(\mathcal{O}_{\tilde{X}}(\tilde{S}), t_{\tilde{S}})) \\ &= \rho_{S*} \tilde{p}_*(\zeta' \cap e(F', \tilde{s}') \cap e(\mathcal{O}_{\tilde{X}'}(\tilde{S}'), t_{\tilde{S}'})) \\ &= p_* \rho_{S' *} (\zeta' \cap e(F', \tilde{s}') \cap e(\mathcal{O}_{\tilde{X}'}(\tilde{S}'), t_{\tilde{S}'})), \end{aligned}$$

where \tilde{p} comes from the fiber diagram

$$\begin{array}{ccc} \tilde{S}' & \xrightarrow{\tilde{p}} & \tilde{S} \\ \rho_{S'} \downarrow & & \downarrow \rho_S \\ S' & \xrightarrow{p} & S. \end{array}$$

Hence by (5.11), (5.12) equals

$$p_* \left(\eta' \cap e(E'|_{\mathfrak{Z}'}, s'|_{\mathfrak{Z}'}) - \rho_{S' *} (\zeta' \cap e(F', \tilde{s}') \cap e(\mathcal{O}_{\tilde{X}'}(\tilde{S}'), t_{\tilde{S}'})) \right) = p_* s'_{\sigma'}(\xi').$$

This proves the proposition. □

Theorem 5.4. *Under Assumption 2.2, we let $g : Z' \rightarrow Z$ be a birational proper morphism with Z' smooth. Consider the fiber product*

$$(5.13) \quad \begin{array}{ccc} Y' & \xrightarrow{f} & Y \\ \mathbf{q}' \downarrow & & \downarrow \mathbf{q} \\ Z' & \xrightarrow{g} & Z \end{array}$$

so that Y' is smooth. Let E', s', σ', X', S' etc be the pullbacks of E, s, σ, X, S etc by f . Let $p : S' \rightarrow S$ denote the restriction of f to S' and $q : S' \rightarrow Z'$ be the restriction of \mathbf{q}' to S' . Then (5.2) holds and we have a commutative diagram

$$(5.14) \quad \begin{array}{ccccccc} H_*(Z') & \xleftarrow{\cong} & H^*(Z') & \xrightarrow{q^*} & H^*(S') & \xrightarrow{[X']_{\text{loc}}^{\text{vir}} \cap} & H_*(S') \\ \downarrow g_* & & & & & & \downarrow p_* \\ H_*(Z) & \xleftarrow{\epsilon_Z} & IH_*(Z) & \xrightarrow{\mathbf{q}^*} & IH_*(Y) & \xrightarrow{s_{\sigma'}^!} & H_*(S) \end{array}$$

where by Corollary 4.6, the virtual cycle for X' is

$$(5.15) \quad [X']_{\text{loc}}^{\text{vir}} = s_{\sigma'}^! [Y'] = [Y'] \cap e_{\sigma'}(E', s') \in H_*(S').$$

Proof. By (5.15) and Proposition 5.2, we have

$$(5.16) \quad \begin{aligned} p_*([X']_{\text{loc}}^{\text{vir}} \cap q^* \alpha) &= p_*(s_{\sigma'}^! [Y'] \cap q^* \alpha) \\ &= p_* s_{\sigma'}^! ([Y'] \cap \mathbf{q}'^* \alpha) = p_* s_{\sigma'}^! \mathbf{q}'^* ([Z'] \cap \alpha) \end{aligned}$$

since Y', Z', \mathbf{q}' are smooth. From the fiber diagram (5.13), we have

$$f_* \mathbf{q}'^* ([Z'] \cap \alpha) = \mathbf{q}^* g_* ([Z'] \cap \alpha) = \mathbf{q}^* \epsilon_Z(v) = \epsilon_Y(\mathbf{q}^* v)$$

since \mathbf{q} is smooth. Therefore, by Proposition 5.3, (5.16) equals $s_{\sigma'}^! \mathbf{q}^* v$ as desired. □

References

- [1] K. Behrend and B. Fantechi. *The intrinsic normal cone*. Invent. Math. **128** (1997), 45–88.
- [2] A. Beilinson, J. Bernstein and P. Deligne. *Faisceaux pervers*. Analysis and topology on singular spaces, I (Luminy, 1981), 5–171, Astérisque, **100**, Soc. Math. France, Paris, 1982.

- [3] A. Borel et al. *Intersection cohomology*. Notes on the seminar held at the University of Bern, Bern, 1983.
- [4] Bredon. *Sheaf theory*. Graduate Texts in Mathematics, **170**. Springer-Verlag, New York, 1997.
- [5] H.-L. Chang, J. Li and W.-P. Li. *Witten's top Chern class via cosection localization*. *Invent. Math.* **200** (2015), no. 3, 1015–1063.
- [6] A. Chiodo. The Witten top Chern class via K-theory. *J. Algebraic Geom.* **15** (2006), no. 4, 681–707.
- [7] H. Fan, T. Jarvis and Y. Ruan. *The Witten equation, mirror symmetry, and quantum singularity theory*. *Ann. of Math. (2)* **178** (2013), no. 1, 1–106.
- [8] H. Fan, T. Jarvis and Y. Ruan. *A mathematical theory of the gauged linear sigma model*. *Geom. Topol.* **22**, No. 1 (2018), 235–303.
- [9] W. Fulton. *Intersection theory*. Second edition. *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics*, 2. Springer-Verlag, Berlin, 1998.
- [10] M. Goresky and R. MacPherson. *Intersection cohomology. II*. *Invent. Math.* **72** (1983), no. 1, 77–129.
- [11] B. Iverson. *Cohomology of sheaves*. Universitext. Springer-Verlag, Berlin, 1986.
- [12] Y.-H. Kiem and J. Li. *Localizing virtual cycles by cosections*. *J. Amer. Math. Soc.* **26** (2013), no. 4, 1025–1050.
- [13] Y.-H. Kiem and J. Li. *Quantum singularity theories via cosection localization*. *J. Reine Angew. Math.* **766** (2020), 73–107.
- [14] D. Massey. *The Sebastiani-Thom isomorphism in the derived category*. *Compositio Math.* **125** (2001), no. 3, 353–362.
- [15] A. Polishchuk and A. Vaintrob. *Algebraic construction of Witten's top Chern class*. *Advances in algebraic geometry motivated by physics (Lowell, MA, 2000)*, 229–249, *Contemp. Math.*, 276, Amer. Math. Soc., Providence, RI, 2001.
- [16] A. Polishchuk and A. Vaintrob. *Matrix factorizations and cohomological field theories*. *J. Reine Angew. Math.* **714** (2016), 1–122.

- [17] C. Voisin. *Hodge theory and complex algebraic geometry. II*. Cambridge Studies in Advanced Mathematics, 77. Cambridge University Press, Cambridge, 2007.
- [18] E. Witten. *Algebraic geometry associated with matrix models of two-dimensional gravity*. Topological methods in modern mathematics (Stony Brook, NY, 1991), 235–269, Publish or Perish, Houston, TX, 1993.

DEPARTMENT OF MATHEMATICS
HONG KONG UNIVERSITY OF SCIENCE AND TECHNOLOGY
CLEARWATER BAY, KOWLOON, HONG KONG
E-mail address: mahlchang@ust.hk

DEPARTMENT OF MATHEMATICS AND RESEARCH INSTITUTE OF MATHEMATICS
SEOUL NATIONAL UNIVERSITY, SEOUL 08826, KOREA
E-mail address: kiem@snu.ac.kr

SHANGHAI CENTER FOR MATHEMATICAL SCIENCES
FUDAN UNIVERSITY, SHANGHAI, CHINA
E-mail address: lijun2210@fudan.edu.cn

RECEIVED JANUARY 30, 2019

ACCEPTED JULY 13, 2019