

## ENERGY CONSERVATION FOR 2D EULER WITH VORTICITY IN $L(\log L)^{\alpha*}$

GENNARO CIAMPA<sup>†</sup>

**Abstract.** In these notes we discuss the conservation of the energy for weak solutions of the two-dimensional incompressible Euler equations. Weak solutions with vorticity in  $L_t^\infty L_x^p$  with  $p \geq 3/2$  are always conservative, while for less integrable vorticity the conservation of the energy may depend on the approximation method used to construct the solution. Here we prove that the canonical approximations introduced by DiPerna and Majda provide conservative solutions when the initial vorticity is in the class  $L(\log L)^\alpha$  with  $\alpha > 1/2$ .

**Keywords.** 2D Euler equations; vanishing viscosity; vortex methods; conservation of energy.

**AMS subject classifications.** 35Q35; 35Q31.

### 1. Introduction

The motion of an incompressible, homogeneous, planar fluid is described by the system of the 2D Euler equations

$$\begin{cases} \partial_t u + (u \cdot \nabla) u + \nabla p = 0, \\ \operatorname{div} u = 0, \\ u(0, \cdot) = u_0, \end{cases} \quad (1.1)$$

where  $u: [0, T] \times \mathbb{R}^2 \rightarrow \mathbb{R}^2$  is the velocity of the fluid,  $p: [0, T] \times \mathbb{R}^2 \rightarrow \mathbb{R}$  is the pressure and  $u_0: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  is a given initial configuration. The first set of equations are derived from Newton's second law while the divergence-free condition expresses the conservation of mass. A peculiar fact of the 2D case is that the vorticity  $\omega$ , defined as

$$\omega = \partial_{x_1} u_2 - \partial_{x_2} u_1,$$

is a scalar quantity which is advected by the velocity  $u$ . In fact, the Equations (1.1) can be rewritten in the vorticity formulation

$$\begin{cases} \partial_t \omega + u \cdot \nabla \omega = 0, \\ u = K * \omega, \\ \omega(0, \cdot) = \omega_0, \end{cases} \quad (1.2)$$

where  $K(x) = x^\perp / (2\pi|x|^2)$  is the 2D Biot-Savart kernel. Note that the Equation (1.2) is a non-linear and non-local transport equation.

The well-posedness of (1.1) is an old and outstanding problem. For smooth initial data, the existence and uniqueness of classical solutions was proved in [18, 28]. The existence of weak solutions has been proved by DiPerna and Majda in [16] by assuming that the initial datum  $\omega_0 \in L^1 \cap L^p(\mathbb{R}^2)$  with  $1 < p \leq \infty$ . Besides this result, the goal of [16] was to develop a rigorous framework for the study of *approximate solution sequences* of the two-dimensional Euler equations. In particular, the authors proved a general

---

\*Received: March 03, 2021; Accepted (in revised form): October 06, 2021. Communicated by Alexander Kiselev.

<sup>†</sup>BCAM - Basque Center for Applied Mathematics, Alameda de Mazarredo 14, E48009 Bilbao, Basque Country - Spain ([gciampa@bcamath.org](mailto:gciampa@bcamath.org)).

compactness theorem towards measure-valued solutions by assuming that  $\omega_0$  is a *vortex-sheet*, i.e.  $\omega_0 \in \mathcal{M} \cap H_{\text{loc}}^{-1}(\mathbb{R}^2)$ . They described three different methods to construct approximate solution sequences:

- (ES) approximation by exact smooth solutions of (1.1);
- (VV) vanishing viscosity from the two-dimensional Navier-Stokes equations;
- (VB) vortex-blob approximation.

In [16] DiPerna and Majda showed the existence of weak solutions via a compactness argument based on the methods (ES) and (VV). The counterpart for the vortex-blob method was proved by Beale in [2]. In these results, the  $L^p$ -integrability with  $1 < p \leq \infty$  of  $\omega_0$  is crucial in order to use Sobolev embeddings which guarantee the strong compactness in  $L^2$  of a sequence of approximating velocity fields. This is enough to deal with the non-linear term in the equations. However, in the case  $\omega_0$  is just  $L^1$  or a measure with distinguished sign, it turns out that the limit vector field is a solution of (1.1) even though concentrations may occur in the non-linearity. This is a purely 2D phenomenon known as *concentration-cancellation*, and it was studied first in [15, 25].

The uniqueness of weak solutions in the class considered in [16] is still an open problem, contrary to the case  $p = \infty$  which has been proved by Yudovich [29]. There exist several partial results towards the non-uniqueness in the case of unbounded initial vorticity, see [5, 6, 21, 26, 27].

Smooth solutions of (1.1) are known to be *conservative*, which means that  $\|u(t)\|_{L^2} = \|u_0\|_{L^2}$  for all times, while this property is not trivial when we consider weak solutions. The problem of the energy conservation, assuming only integrability conditions on the vorticity, has been addressed in [9]: The authors consider the 2D Euler equations on the two-dimensional flat torus  $\mathbb{T}^2$  and they prove that *all* weak solutions satisfy the energy conservation if the vorticity  $\omega \in L^\infty((0, T); L^p(\mathbb{T}^2))$  with  $p \geq 3/2$ . The proof is based on a mollification argument and the exponent  $p = 3/2$  is required in order to have weak continuity of a commutator term in the energy balance. The authors also give an example of the sharpness of the exponent  $p = 3/2$  in their argument, but that still leaves open the question of the existence of non-conservative solutions below this integrability threshold. Moreover, they show that if  $\omega \in L^\infty((0, T); L^p(\mathbb{T}^2))$ , with  $1 < p < 3/2$ , solutions constructed via (ES) and (VV) conserve the kinetic energy.

Here we discuss the conservation of the energy for solutions of the 2D Euler equations when the initial vorticity is slightly more than integrable, namely  $\omega_0 \in L^1 \cap L(\log L)^\alpha(\mathbb{R}^2)$  with  $\alpha > 1/2$ . Existence of weak solutions of (1.1) in this setting was proved by Chae, first in [7] in the case  $\omega_0 \in L^1 \cap L \log L(\mathbb{R}^2)$ , and then extended to the case  $\omega_0 \in L^1 \cap L(\log L)^{1/2}(\mathbb{R}^2)$  in [8]. In these results, the strategy of the proof is based on the properties of Calderón-Zygmund singular integral operators and compact embeddings of Orlicz-Sobolev spaces into  $L_{\text{loc}}^2(\mathbb{R}^2)$ .

In a similar fashion to the framework of DiPerna and Majda, in [19] the authors introduce the definition of  $H_{\text{loc}}^{-1}$ -stability for a sequence of approximating vorticity  $\omega^\varepsilon$ , showing that it is a sharp criterion for the strong  $L_{\text{loc}}^2$ -convergence of an approximate solution sequence  $u^\varepsilon$ . With their approach they are able to recover previous existence results, expanding the set of possible initial data to much more general *rearrangement invariant* spaces, such as the Orlicz spaces  $L(\log L)^\alpha$ , with  $\alpha \geq 1/2$ , and the Lorentz spaces  $L^{(1, q)}$  with  $1 < q \leq 2$ .

Finally, in [17] it has been proven that the strong  $L^2$ -compactness of a sequence of velocity fields constructed via (VV) is equivalent to the energy conservation property. In virtue of this result, by posing the problem in the two-dimensional torus, the authors

obtained as a corollary that the vanishing viscosity limit produce conservative weak solutions for initial vorticity in the rearrangement invariant spaces considered in [19], including  $L(\log L)^\alpha$  with  $\alpha > 1/2$ .

The contribution of these notes in the theory of conservative weak solutions of (1.1) is the following: We consider an initial datum  $u_0 \in L^2(\mathbb{R}^2)$  such that  $\omega_0 \in L(\log L)^\alpha(\mathbb{R}^2)$  with compact support and we prove that the canonical approximations introduced in [16] produce approximate solution sequences such that the velocity converges *globally* in  $L^2$  if  $\alpha > 1/2$ . This allows us to prove that the vortex-blob method yields conservative weak solutions and, in this setting, we extend the results of [9, 17] concerning (ES) and (VV) to the case in which the domain is the whole plane  $\mathbb{R}^2$ . In order to get the strong convergence in  $C([0, T]; L^2(\mathbb{R}^2))$  of the approximating velocity, we will exploit the techniques of [11, 24] by adapting the Serfati identity [1, 23] to this less integrable setting. In particular, it would be crucial that the approximating vorticity converge strongly in  $C([0, T]; L^1(\mathbb{R}^2))$ , as shown recently in [10, 11], obtained as a consequence of the Lagrangian property of the limit vorticity. See also [4, 13, 14] for a deeper understanding of Lagrangian solutions of the 2D Euler equations.

The main theorem of this work can be resumed in the following.

**THEOREM 1.1.** *Let  $\omega_0 \in L(\log L)_c^\alpha(\mathbb{R}^2)$ , with  $\alpha > 1/2$ , with zero total mass, i.e.  $\int_{\mathbb{R}^2} \omega(x) dx = 0$ . Let  $u$  be a weak solution of (1.1), with  $\text{curl} u_0 = \omega_0$ , that can be obtained as a limit of a sequence  $u^n$  constructed via one of the methods (ES), (VV), (VB). Then,  $u^n$  satisfies the following convergence*

$$u^n \rightarrow u \quad \text{in } C([0, T]; L^2(\mathbb{R}^2)), \tag{1.3}$$

and  $u$  is conservative.

**2. The two-dimensional Euler equations**

The goal of this section is to provide some preliminary results on weak solutions of the 2D Euler equations. First, we introduce the notations used in the paper. Then, we will pay particular attention to the theory developed by DiPerna and Majda in [16]. Finally, we will summarize some more recent results concerning conservative weak solutions.

**2.1. Notations.** We will denote by  $L^p(\mathbb{R}^d)$  the standard Lebesgue spaces and with  $\|\cdot\|_{L^p}$  their norm. Moreover,  $L_c^p(\mathbb{R}^d)$  denotes the space of  $L^p$  functions defined on  $\mathbb{R}^d$  with compact support. The Sobolev space of  $L^p$  functions with distributional derivatives of first order in  $L^p$  is denoted by  $W^{1,p}(\mathbb{R}^d)$ . The spaces  $L_{\text{loc}}^p(\mathbb{R}^d), W_{\text{loc}}^{1,p}(\mathbb{R}^d)$  denote the space of functions which are locally in  $L^p(\mathbb{R}^d), W^{1,p}(\mathbb{R}^d)$  respectively. We will denote by  $H^1(\mathbb{R}^d)$  the space  $W^{1,2}(\mathbb{R}^d)$  and by  $H^{-1}(\mathbb{R}^d)$  its dual space. Moreover, we will say that a function  $u$  is in  $H_{\text{loc}}^{-1}(\mathbb{R}^d)$  if  $\rho u \in H^{-1}(\mathbb{R}^d)$  for every function  $\rho \in C_c^\infty(\mathbb{R}^d)$ . We denote with  $L(\log L)^\alpha(\mathbb{R}^d)$  the space of functions  $f$  such that the following quantity is finite

$$\int_{\mathbb{R}^d} |f(x)| (\log^+(|f(x)|))^\alpha dx. \tag{2.1}$$

The space  $L(\log L)^\alpha(\mathbb{R}^d)$  is a Banach space if endowed with the Luxemburg norm

$$\|f\|_{L(\log L)^\alpha} = \inf \left\{ k > 0 : \int_{\mathbb{R}^d} \frac{|f|}{k} \left( \log^+ \left( \frac{|f|}{k} \right) \right)^\alpha dx \leq 1 \right\}, \tag{2.2}$$

where the function  $\log^+$  is defined as

$$\log^+(t) = \begin{cases} \log(t) & \text{if } t \geq 1, \\ 0 & \text{otherwise,} \end{cases}$$

and  $L(\log L)_c^\alpha(\mathbb{R}^d)$  will be the space of functions in  $L(\log L)^\alpha(\mathbb{R}^d)$  with compact support. We remark that the quantity in (2.1) does not define a norm therefore the necessity of the definition (2.2). We refer to the classical book of Bennett and Sharpley [3] for more details.

We denote by  $L^p((0, T); L^q(\mathbb{R}^d))$  the space of all measurable functions  $u$  defined on  $[0, T] \times \mathbb{R}^d$  such that

$$\|u\|_{L^p((0, T); L^q(\mathbb{R}^d))} := \left( \int_0^T \|u(t, \cdot)\|_{L^q}^p dt \right)^{\frac{1}{p}} < \infty,$$

for all  $1 \leq p < \infty$ , and

$$\|u\|_{L^\infty((0, T); L^q(\mathbb{R}^d))} := \operatorname{ess\,sup}_{t \in [0, T]} \|u(t, \cdot)\|_{L^q} < \infty,$$

and analogously for the spaces  $L^p((0, T); W^{1, q}(\mathbb{R}^d))$ . The space of continuous functions on  $[0, T]$  with values in  $L^p(\mathbb{R}^d)$  is denoted by  $C([0, T]; L^p(\mathbb{R}^d))$  and it is endowed with the norm  $\|\cdot\|_{L^\infty((0, T); L^p(\mathbb{R}^d))}$ . We denote by  $B_R$  the ball of radius  $R > 0$  centered in the origin of  $\mathbb{R}^d$ . In the estimates we will denote with  $C$  a positive constant which may change from line to line. Moreover, for  $\alpha \in \mathbb{R}$  we define the function

$$\beta(s) := s(\log(e + s))^\alpha, \tag{2.3}$$

which will be used extensively in what follows. Finally, it is useful to denote with  $\star$  the following variant of the convolution

$$v \star w = \sum_{i=1}^2 v_i \star w_i \quad \text{if } v, w \text{ are vector fields in } \mathbb{R}^2,$$

$$A \star B = \sum_{i, j=1}^2 A_{ij} \star B_{ij} \quad \text{if } A, B \text{ are matrix-valued functions in } \mathbb{R}^2.$$

With the notations above, it is easy to check that if  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  is a scalar function and  $v: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  is a vector field, then

$$f \star \operatorname{curl} v = \nabla^\perp f \star v,$$

$$\nabla^\perp f \star \operatorname{div}(v \otimes v) = \nabla \nabla^\perp f \star (v \otimes v).$$

**2.2. Weak solutions.** We recall the definition of weak solution of the Euler equations as in [16].

**DEFINITION 2.1.** *A vector valued function  $u \in L^\infty((0, T); L^2_{\operatorname{loc}}(\mathbb{R}^2))$  is a weak solution of (1.1) if it satisfies:*

(1) for all test functions  $\Phi \in C_0^\infty((0, T) \times \mathbb{R}^2)$  with  $\operatorname{div} \Phi = 0$ ,

$$\int_0^T \int_{\mathbb{R}^2} (\partial_t \Phi \cdot u + \nabla \Phi : u \otimes u) \, dx \, dt = 0; \tag{2.4}$$

(2)  $\operatorname{div} u = 0$  in the sense of distributions;

(3)  $u \in \operatorname{Lip}([0, T]; H_{\operatorname{loc}}^{-L}(\mathbb{R}^2))$  for some  $L > 0$  and  $u(0, x) = u_0(x)$ .

REMARK 2.1. The choice of divergence-free test functions removes the pressure in the weak formulation (2.4). However, it can be formally recovered by the formula

$$-\Delta p = \operatorname{div} \operatorname{div}(u \otimes u),$$

which is obtained by applying the divergence in the momentum equation in (1.1).

In [16], DiPerna and Majda introduced the following definition of an *approximate solution sequence* of the 2D Euler equations.

DEFINITION 2.2. A sequence of smooth velocity fields  $u^n$  with vorticity  $\operatorname{curl} u^n = \omega^n \in C([0, T]; L^1(\mathbb{R}^2))$  is an approximate solution sequence for the 2D Euler equations provided that

(i)  $u^n$  has uniformly bounded local kinetic energy and  $u^n$  is incompressible, i.e., for each  $R > 0$  and  $T > 0$ , there exists  $C(R) > 0$  such that

$$\max_{t \in [0, T]} \int_{B_R} |u^n(t, x)|^2 \, dx \leq C(R), \quad \operatorname{div} u^n = 0;$$

(ii) the vorticity  $\omega^n$  is uniformly bounded in  $L^1$ , i.e., for every  $T > 0$ ,

$$\max_{t \in [0, T]} \int_{\mathbb{R}^2} |\omega^n(t, x)| \, dx \leq C;$$

(iii) for some  $L > 0$ , the sequence  $u^n$  is uniformly bounded in  $\operatorname{Lip}([0, T]; H_{\operatorname{loc}}^{-L}(\mathbb{R}^2))$ ;

(iv)  $u^n$  is weakly consistent with the 2D Euler equations, i.e.

$$\lim_{n \rightarrow \infty} \int_0^T \int_{\mathbb{R}^2} (\partial_t \Phi \cdot u^n + \nabla \Phi : u^n \otimes u^n) \, dx \, dt = 0, \tag{2.5}$$

for every  $\Phi \in C_c^\infty((0, T) \times \mathbb{R}^2)$  with  $\operatorname{div} \Phi = 0$ .

Besides the very general definition, in [16] the authors give three different examples of approximate solutions sequences, which are important for physical or numerical reasons. They are the following.

**(ES) Approximation by exact smooth solutions of (1.1).** We consider a smooth approximation of the initial datum  $u_0^\delta$  such that  $u_0^\delta \rightarrow u_0$  in  $L_{\operatorname{loc}}^2$  and we define  $u^\delta$  as the unique solution of the approximating problem

$$\begin{cases} \partial_t u^\delta + (u^\delta \cdot \nabla) u^\delta + \nabla p^\delta = 0, \\ \operatorname{div} u^\delta = 0, \\ u^\delta(0, \cdot) = u_0^\delta. \end{cases} \tag{2.6}$$

Then, a solution  $u$  of (1.1) is constructed analyzing the limit of the sequence  $u^\delta$  as  $\delta \rightarrow 0$ .

**(VV) Vanishing viscosity from the two-dimensional Navier-Stokes equations.** We consider the two-dimensional incompressible Navier-Stokes equations

$$\begin{cases} \partial_t u^\nu + (u^\nu \cdot \nabla)u^\nu + \nabla p^\nu = \nu \Delta u^\nu, \\ \operatorname{div} u^\nu = 0, \\ u^\nu(0, \cdot) = u_0^\nu, \end{cases} \tag{2.7}$$

where  $\nu > 0$  is the viscosity of the fluid and  $u_0^\nu$  is smooth and converges in  $L^2_{\text{loc}}$  towards  $u_0$  as  $\nu \rightarrow 0$ . Then, a solution  $u$  of (1.1) is constructed analyzing the vanishing viscosity limit of the sequence  $u^\nu$ .

**(VB) Vortex-blob approximation.** It is a numerical method which is the prototype of several important numerical schemes. It is based on the idea of approximating the vorticity with a finite number of cores which evolve according to the velocity of the fluid. Without going into details, the approximating velocity  $u^\varepsilon$  solves the system

$$\begin{cases} \partial_t u^\varepsilon + (u^\varepsilon \cdot \nabla)u^\varepsilon + \nabla p^\varepsilon = K * E_\varepsilon, \\ \operatorname{div} u^\varepsilon = 0, \\ u^\varepsilon(0, \cdot) = u_0^\varepsilon, \end{cases} \tag{2.8}$$

where  $u_0^\varepsilon$  is a suitable smooth approximation of the initial datum and  $E_\varepsilon$  is an error term which comes from the fact that, roughly speaking, each blob is rigidly translated by the flow. We give the precise construction together with its main properties in the Appendix.

By assuming only integrability hypothesis on the initial vorticity  $\omega_0$ , the existence of weak solutions constructed with the methods above has been proven in [2, 16]. For simplicity of exposition, for the remainder of this subsection we will use  $n$  as an approximation parameter for all the three methods.

**THEOREM 2.1.** *Let  $u_0 \in L^2_{\text{loc}}(\mathbb{R}^2)$  be a divergence-free vector field and let  $\omega_0 = \operatorname{curl} u_0 \in L^p_c(\mathbb{R}^2)$  for some  $p > 1$ . Let  $u^n$  be an approximate solution sequence constructed via one of the methods (ES), (VV), (VB), where the associated initial datum  $u^n \rightarrow u_0$  in  $L^2_{\text{loc}}(\mathbb{R}^2)$ . Then, there exists a subsequence of  $u^n$  and a vector field  $u \in L^\infty((0, T); L^2_{\text{loc}}(\mathbb{R}^2)) \cap \operatorname{Lip}([0, T]; H^{-L}_{\text{loc}}(\mathbb{R}^2))$  with the following properties:*

- $u(0, \cdot) = u_0$ ,
- $u^n \rightarrow u$  in  $L^2((0, T); L^2_{\text{loc}}(\mathbb{R}^2))$ ,
- $\omega^n \xrightarrow{*} \omega$  in  $L^\infty((0, T); L^p(\mathbb{R}^2))$ ,
- $\omega^n \rightarrow \omega$  in  $C([0, T]; H^{-L-1}_{\text{loc}}(\mathbb{R}^2))$ .

**REMARK 2.2.** Note that the setting of the previous theorem is for a regime where the uniqueness of solutions of (1.1) is not known. Therefore, the three methods could have multiple limit points which may also change depending on the approximation.

As already mentioned in the Introduction, the previous theorem has been generalized by Chae [7, 8]:

**THEOREM 2.2.** *Let  $u_0 \in L^2_{\text{loc}}(\mathbb{R}^2)$  be a divergence-free vector field such that  $\operatorname{curl} u_0 = \omega_0 \in L(\log L)^\alpha_c(\mathbb{R}^2)$  with  $\alpha \geq 1/2$ . Then, there exists a weak solution  $u$  of (1.1) with initial datum  $u_0$  satisfying*

$$u \in C([0, T]; L^2_{\text{loc}}(\mathbb{R}^2)). \tag{2.9}$$

The proof of Theorem 2.2 strongly relies on the fact that the operator

$$T : f \in L(\log L)_c^\alpha(\mathbb{R}^2) \rightarrow K * f \in L_{loc}^2(\mathbb{R}^2), \tag{2.10}$$

is compact for  $\alpha > 1/2$ , where  $K$  is the two dimensional Biot-Savart kernel. It is worth to note that the solutions are constructed analyzing the vanishing viscosity limit of the corresponding Navier-Stokes equations with the same initial data. We now recall an intriguing example from [24].

EXAMPLE 2.1. *Let us consider the function  $\omega_0$  given by*

$$\omega_0(x) := \begin{cases} \frac{1}{|x|^2 \left(\log\left(\frac{1}{|x|}\right)\right)^{1+\gamma}}, & \text{if } |x| < e^{-1}, \\ 0, & \text{otherwise.} \end{cases} \tag{2.11}$$

*It is a classical fact that, since  $\omega_0$  depends only on  $|x|$ , the corresponding velocity field is given by*

$$u_0(x) = \frac{x^\perp}{|x|^2} \int_0^{|x|} s \omega_0(s) ds, \tag{2.12}$$

*see [16]. By using that*

$$\log(e + \omega_0) \leq C \log(1/|x|),$$

*when  $\omega_0 > 0$ , a direct computation shows that*

- *for  $\alpha \in (0, 1]$ , choosing  $\gamma > \alpha$ , the function  $\omega_0 \in L_c(\log L)^\alpha(\mathbb{R}^2)$  but  $\omega_0 \notin L^p(\mathbb{R}^2)$  for every  $p > 1$ ;*
- *for  $\alpha < \gamma < 1/2$ ,  $u_0$  is not locally square integrable;*
- *for  $1/2 < \alpha < \gamma$ ,  $u_0$  is locally square integrable.*

*Moreover, as remarked in [24], an argument based on [8], shows that  $u_0 \in L_{loc}^2(\mathbb{R}^2)$  when  $\gamma > \alpha = 1/2$ .*

We conclude this subsection by summarizing some known results about the strong convergence in  $C([0, T]; L^p(\mathbb{R}^2))$  of the approximating vorticity  $\omega^n$ . This problem has been addressed by several authors in different settings, especially with regard to the inviscid limit of the Navier-Stokes equations, see for example [10, 12, 22]. We collect the results we need in the following theorem, see [4, 10, 11].

THEOREM 2.3. *Let  $\omega_0 \in L_c^p(\mathbb{R}^2)$  with  $p \geq 1$  and let  $\omega^n$  be a sequence of approximating vorticity constructed via one of the three methods (ES), (VV), or (VB). Then, there exists  $\omega \in C([0, T]; L^1 \cap L^p(\mathbb{R}^2))$  such that*

$$\omega^n \rightarrow \omega \quad \text{in } C([0, T]; L^1 \cap L^p(\mathbb{R}^2)). \tag{2.13}$$

REMARK 2.3. Being  $L(\log L)_c^\alpha \subset L_c^1$ , assuming  $\omega_0 \in L(\log L)_c^\alpha$  by Theorem 2.3 if  $\omega^n$  is a sequence constructed via one of the aforementioned methods, then there exists  $\omega \in C([0, T]; L^1(\mathbb{R}^2))$  such that

$$\omega^n \rightarrow \omega \quad \text{in } C([0, T]; L^1(\mathbb{R}^2)). \tag{2.14}$$

**2.3. Conservative solutions.** In this subsection we discuss the conservation of the energy for the 2D Euler equations. We recall the following definition.

**DEFINITION 2.3.** *Let  $u \in C([0, T]; L^2(\mathbb{R}^2))$  be a weak solution of (1.1) with initial datum  $u_0 \in L^2(\mathbb{R}^2)$ . We say that  $u$  is a conservative weak solution if*

$$\|u(t, \cdot)\|_{L^2} = \|u_0\|_{L^2} \quad \forall t \in [0, T].$$

It is well-known that in the two-dimensional case, even if we assume that the vorticity is bounded, the velocity field is in general not square integrable. In order to define the kinetic energy, we need to require that the vorticity has zero total mass, see [20, Proposition 3.3] for more details.

**PROPOSITION 2.1.** *Let  $u \in L^2_{loc}(\mathbb{R}^2)$  be an incompressible velocity field in  $\mathbb{R}^2$  such that its vorticity  $\omega = \operatorname{curl} u \in L^1_c(\mathbb{R}^2)$  and it has zero total mass, i.e.*

$$\int_{\mathbb{R}^2} \omega(x) \, dx = 0. \tag{2.15}$$

*Then,  $u$  is globally square integrable, i.e.  $u \in L^2(\mathbb{R}^2)$ .*

As already explained in the Introduction, the problem of the conservation of the energy in this low regularity setting has been addressed in [9]: they showed that every weak solution is conservative if  $\operatorname{curl} u \in L^\infty((0, T); L^p(\mathbb{T}^2))$  with  $p \geq 3/2$ , while for less integrable vorticities the conservation of the energy may depend on the approximation procedure. In particular, by collecting the results of [9–11] we have the following theorem.

**THEOREM 2.4.** *Let  $u \in C([0, T]; L^2(\mathbb{R}^2))$  be a weak solution of (1.1) with vorticity  $\omega = \operatorname{curl} u$  of zero total mass. Then,*

- *if  $\omega \in L^\infty((0, T); L^1 \cap L^{\frac{3}{2}}(\mathbb{R}^2))$ , then  $u$  is conservative;*
- *if  $\omega \in L^\infty((0, T); L^1 \cap L^p(\mathbb{R}^2))$  with  $p > 1$ , and  $u$  is constructed as limit of one of the approximations (ES), (VV), or (VB), then  $u$  is conservative.*

We finish this subsection by recalling a theorem that has been proved in [17]. It characterizes the compactness of (VV) and the energy conservation in terms of the classical structure function

$$S_2^T(u; r) := \left( \int_0^T \int_{\mathbb{T}^2} \int_{B_r} |u(t, x+h) - u(t, x)|^2 \, dh \, dx \, dt \right)^{1/2}.$$

The main statement from [17] is the following.

**THEOREM 2.5.** *Let  $u^\nu$  be the unique solution of (2.7) with a smooth initial datum  $u_0^\nu$  such that*

$$u_0^\nu \rightarrow u_0 \quad \text{in } L^2(\mathbb{T}^2).$$

*Let  $u \in L^\infty((0, T); L^2(\mathbb{T}^2))$  be a solution of (1.1) with initial datum  $u_0$  such that, up to a sub-sequence,*

$$u^\nu \rightharpoonup u \quad \text{in } L^2(\mathbb{T}^2).$$

*Then the following are equivalent:*

- (i)  $u^\nu \rightarrow u$  strongly in  $L^p((0, T); L^2(\mathbb{T}^2))$  for some  $1 \leq p < \infty$ ,
- (ii) there exists a bounded modulus of continuity  $\phi(r)$  such that, uniformly in  $\nu$ ,

$$S_2^T(u^\nu; r) \leq \phi(r) \quad \forall r \geq 0,$$

- (iii)  $u$  is a conservative weak solution.

It is important to note that, by using the result in [19], the previous theorem implies that solutions constructed via (VV) are conservative if  $\omega_0 \in L(\log L)^\alpha(\mathbb{T}^2)$ . Then, our Theorem 4.2 will extend the aforementioned result to the class  $\omega_0 \in L(\log L)_c^\alpha(\mathbb{R}^2)$ .

REMARK 2.4. The Theorem 2.5 holds even if we replace the two-dimensional torus  $\mathbb{T}^2$  with the whole plane  $\mathbb{R}^2$ , taking into account the appropriate technical considerations.

### 3. A priori estimates

In this section we summarize some a priori estimates for the approximating vorticity constructed via the approximation methods introduced in Section 2.2. We will always assume that  $\omega_0 \in L(\log L)_c^\alpha(\mathbb{R}^2)$  with  $\alpha > 1/2$ . As already stressed in the Introduction, these estimates will be crucial in order to address the strong convergence of the velocity field in  $C([0, T]; L^2(\mathbb{R}^2))$ , which will be the topic of the next section. It is worth pointing out that we will not estimate quantities like (2.2), as the reader might expect, but we will provide bounds (uniform with respect to the parameter in question) on the following quantity

$$\int_{\mathbb{R}^d} |f(x)|(\log(e + |f(x)|))^\alpha dx. \tag{3.1}$$

This is the type of bound that we will need in Section 4. However, note that if  $f$  is supported on a compact subset of  $\mathbb{R}^d$  (as in the case of  $\omega_0$ ), the following equivalence holds

$$(2.1) < \infty \iff (2.2) < \infty \iff (3.1) < \infty.$$

The same equivalence holds if the support of  $f$  is not compact but in addition  $f \in L^1(\mathbb{R}^d)$ . This latter will be the case of the approximating vorticity.

**3.1. Limit of exact smooth solutions.** Let  $\rho_\delta$  be a standard smooth mollifier and consider the following Cauchy problem

$$\begin{cases} \partial_t \omega^\delta + u^\delta \cdot \nabla \omega^\delta = 0, \\ u^\delta = K * \omega^\delta, \\ \omega^\delta(0, \cdot) = \omega_0^\delta, \end{cases} \tag{3.2}$$

where  $\omega_0^\delta = \omega_0 * \rho_\delta$ . We have the following.

LEMMA 3.1. *Let  $\omega^\delta$  be the unique smooth solution of (3.2). Then,*

$$\sup_{t \in (0, T)} \int_{\mathbb{R}^2} |\omega^\delta(t, x)|(\log(e + |\omega^\delta(t, x)|))^\alpha dx \leq C, \tag{3.3}$$

where  $C$  is a positive constant which does not depend on  $\delta$ .

*Proof.* Let  $\beta$  be as in (2.3) and multiply the equations in (3.2) by  $\beta'(|\omega^\delta|)$ . Then, by integrating in space and time we get that

$$\frac{d}{dt} \int_{\mathbb{R}^2} \beta(|\omega^\delta(t, x)|) dx = 0. \tag{3.4}$$

By using the convexity of  $\beta$  and Jensen’s inequality, it follows that

$$\int_{\mathbb{R}^2} \beta(|\omega_0^\delta|) dx \leq \int_{\mathbb{R}^2} \beta(|\omega_0|) dx \leq C \left( \|\omega_0\|_{L^1} + \int_{\mathbb{R}^2} |\omega_0| (\log^+(|\omega_0|))^\alpha dx \right) < \infty,$$

and then, by integrating in time in (3.4) we have the result. □

**3.2. The vanishing viscosity limit.** We now deal with the vanishing viscosity limit of the Navier-Stokes equations. Let  $\rho_\nu$  a standard smooth mollifier and let  $\omega^\nu$  be the solution of

$$\begin{cases} \partial_t \omega^\nu + u^\nu \cdot \nabla \omega^\nu = \nu \Delta \omega^\nu, \\ u^\nu = K * \omega^\nu, \\ \omega^\nu(0, \cdot) = \omega_0^\nu, \end{cases} \tag{3.5}$$

where  $\omega_0^\nu = \omega_0 * \rho_\nu$ . We have the following.

LEMMA 3.2. *Let  $\omega^\nu$  be the unique smooth solution of (3.5). Then,*

$$\sup_{t \in (0, T)} \int_{\mathbb{R}^2} |\omega^\nu(t, x)| (\log(e + |\omega^\nu(t, x)|))^\alpha dx \leq C, \tag{3.6}$$

where  $C$  is a positive constant which does not depend on  $\nu$ .

*Proof.* We just sketch the proof since it is very similar to the one of Lemma 3.1. Let  $\beta$  be as in (2.3) and multiply the equations in (3.5) by  $\beta'(|\omega^\nu|)$ . Then, by integrating in space and time we get that

$$\frac{d}{dt} \int_{\mathbb{R}^2} \beta(|\omega^\nu(t, x)|) dx = -\nu \int_0^T \int_{\mathbb{R}^2} |\nabla \omega^\nu(t, x)|^2 \beta''(|\omega^\nu(t, x)|) dx dt \leq 0, \tag{3.7}$$

since  $\beta$  is convex. Then, integrating in time (3.7) we have the result. □

**3.3. The vortex-blob method.** We finally deal with the vortex-blob method. The reader can find the precise definition of the vortex-blob method and some of its properties in the Appendix at the end of this note.

LEMMA 3.3. *Let  $\omega^\varepsilon$  be the approximating vorticity constructed via the vortex-blob method. Then,*

$$\sup_{t \in (0, T)} \int_{\mathbb{R}^2} |\omega^\varepsilon(t, x)| (\log(e + |\omega^\varepsilon(t, x)|))^\alpha dx \leq C, \tag{3.8}$$

where  $C$  is a positive constant which does not depend on  $\varepsilon$ .

*Proof.* We start by proving that  $\omega^\varepsilon(0, \cdot)$  satisfies the bound (3.8), see (A.5). Let  $\beta$  be as in (2.3),  $\rho_{\delta(\varepsilon)}$  be a standard mollifier, and  $\omega_0^\varepsilon = \omega_0 * \rho_{\delta(\varepsilon)}$ . By Jensen’s inequality we have that

$$\int_{\mathbb{R}^2} \beta(|\omega_0 * \rho_{\delta(\varepsilon)}|) dx \leq \int_{\mathbb{R}^2} \beta(|\omega_0|) dx.$$

We consider now  $\omega_0^\varepsilon * \varphi_\varepsilon$  where  $\varphi_\varepsilon$  is the mollifier which appears in the definition of the approximate vorticity (A.2). Again by Jensen’s inequality we have

$$\int_{\mathbb{R}^2} \beta(|\omega_0^\varepsilon * \varphi_\varepsilon|) dx \leq \int_{\mathbb{R}^2} \beta(|\omega_0|) dx.$$

Then, by the properties of the function  $\beta$  we have that

$$\int_{\mathbb{R}^2} \beta(|\omega^\varepsilon(0,x)|) dx \leq C \int_{\mathbb{R}^2} \beta(|\omega^\varepsilon(0,x) - \omega_0^\varepsilon * \varphi_\varepsilon(x)|) dx + C \int_{\mathbb{R}^2} \beta(|\omega_0^\varepsilon * \varphi_\varepsilon|) dx.$$

We know that the second term on the right-hand side is uniformly bounded in  $\varepsilon$ , while for the first term we have that

$$\begin{aligned} \int_{\mathbb{R}^2} |\omega^\varepsilon(0,x) - \omega_0^\varepsilon * \varphi_\varepsilon(x)| \log(e + |\omega^\varepsilon(0,x) - \omega_0^\varepsilon * \varphi_\varepsilon(x)|) dx \\ \leq \int_{\mathbb{R}^2} |\omega^\varepsilon(0,x) - \omega_0^\varepsilon * \varphi_\varepsilon(x)| \log(e + C\varepsilon) dx \\ \leq \log(e + C\varepsilon) \|\omega^\varepsilon(0, \cdot) - \omega_0^\varepsilon * \varphi_\varepsilon\|_{L^1} \\ \leq C\varepsilon^3 \log(e + C\varepsilon) \leq C, \end{aligned}$$

where we have used Lemma A.1 with  $p = \infty$  in the second line and with  $p = 1$  in the fourth line. As a consequence of the previous estimate we obtain

$$\int_{\mathbb{R}^2} |\omega^\varepsilon(0,x)| (\log(e + |\omega^\varepsilon(0,x)|))^\alpha dx \leq C.$$

Let  $u^\varepsilon$  be the velocity field constructed with the vortex-blob method and consider the linear problem

$$\begin{cases} \partial_t \bar{\omega}^\varepsilon + u^\varepsilon \cdot \nabla \bar{\omega}^\varepsilon = 0, \\ \bar{\omega}^\varepsilon(0, \cdot) = \omega^\varepsilon(0, \cdot). \end{cases} \tag{3.9}$$

By arguing as in the proof of Lemma 3.1 we have that

$$\int_{\mathbb{R}^2} |\bar{\omega}^\varepsilon| (\log(e + |\bar{\omega}^\varepsilon|))^\alpha dx = \int_{\mathbb{R}^2} |\omega^\varepsilon(0,x)| (\log(e + |\omega^\varepsilon(0,x)|))^\alpha dx \leq C,$$

from which it follows that

$$\int_{\mathbb{R}^2} \beta(|\bar{\omega}^\varepsilon * \varphi_\varepsilon|(t,x)) dx \leq C.$$

So, in the end we get that

$$\begin{aligned} \int_{\mathbb{R}^2} \beta(|\omega^\varepsilon|(t,x)) dx &\leq C \int_{\mathbb{R}^2} \beta(|\bar{\omega}^\varepsilon * \varphi_\varepsilon|(t,x)) dx + C \int_{\mathbb{R}^2} \beta(|\omega^\varepsilon - \bar{\omega}^\varepsilon * \varphi_\varepsilon|(t,x)) dx \\ &\leq C + C\varepsilon^3 (\log(e + C\varepsilon))^\alpha \leq C, \end{aligned}$$

which concludes the proof. □

#### 4. Strong convergence of the velocity field and conservation of the energy

In this section we will prove our main result, namely Theorem 1.1. In particular, the uniform bound proved in Section 3 will be crucial in order to prove the global strong convergence in  $C([0, T]; L^2(\mathbb{R}^2))$ . To make the presentation easier to follow, we split Theorem 1.1 into three theorems, one for each approximation. We start by proving the result for (ES), then with the appropriate modifications we will describe how to prove such a result also for (VV) and (VB).

**THEOREM 4.1.** *Let  $\omega_0 \in L(\log L)_c^\alpha(\mathbb{R}^2)$ , with  $\alpha > 1/2$ , with zero total mass. Let  $u$  be a weak solution of (1.1), with  $\text{curl} u_0 = \omega_0$ , that can be obtained as a limit of a sequence  $u^\delta$  constructed via (ES). Then,  $u^\delta$  satisfies the following convergence*

$$u^\delta \rightarrow u \quad \text{in } C([0, T]; L^2(\mathbb{R}^2)), \tag{4.1}$$

and  $u$  is conservative.

*Proof.* In order to prove the convergence stated in (4.1), we will prove that  $u^\delta$  is a Cauchy sequence in  $C([0, T]; L^2(\mathbb{R}^2))$ . We recall that the parameter  $\delta$  is always supposed to vary over a countable set, therefore given the sequence  $\delta_n \rightarrow 0$ , we denote with  $u^n$  and  $\omega^n$  the sequences  $u^{\delta_n}$  and  $\omega^{\delta_n}$ . We divide the proof in several steps.

Step 1 *A Serfati identity with fixed vorticity.*

In this step we derive a formula for the approximate velocity  $u^n$ , in the same spirit of the Serfati identity derived in [1, 23].

Let  $a \in C_c^\infty(\mathbb{R}^2)$  be a smooth function such that  $a(x) = 1$  if  $|x| < 1$  and  $a(x) = 0$  for  $|x| > 2$ . Differentiating in time the Biot-Savart formula we obtain that for  $i = 1, 2$

$$\begin{aligned} \partial_s u_i^n(s, x) &= K_i * (\partial_s \omega^n)(s, x) \\ &= (aK_i) * (\partial_s \omega^n)(s, x) + [(1-a)K_i] * (\partial_s \omega^n)(s, x). \end{aligned} \tag{4.2}$$

Now we use the equation for  $\omega^n$  obtaining

$$\partial_s \omega^n = -u^n \cdot \nabla \omega^n,$$

and substituting in (4.2) we get

$$\partial_s u_i^n = (aK_i) * (\partial_s \omega^n) - [(1-a)K_i] * (u^n \cdot \nabla \omega^n). \tag{4.3}$$

By using the identity

$$u^n \cdot \nabla \omega^n = \text{curl}(u^n \cdot \nabla u^n) = \text{curl} \text{div}(u^n \otimes u^n),$$

we obtain that

$$[(1-a)K_i] * (u^n \cdot \nabla \omega^n) = (\nabla \nabla^\perp [(1-a)K_i]) \star (u^n \otimes u^n). \tag{4.4}$$

Substituting the expressions (4.4) in (4.2) and integrating in time we have that  $u^n$  satisfies the following formula, known as *Serfati identity*:

$$\begin{aligned} u_i^n(t, x) &= u_i^n(0, x) + (aK_i) * (\omega^n(t, \cdot) - \omega^n(0, \cdot))(x) \\ &\quad - \int_0^t (\nabla \nabla^\perp [(1-a)K_i]) \star (u^n(s, \cdot) \otimes u^n(s, \cdot))(x) ds. \end{aligned} \tag{4.5}$$

We modify the Serfati identity (4.5) by introducing a new cut-off function  $a_\varepsilon$ : let  $\varepsilon \in (0, 1)$  and define  $a_\varepsilon$  to be equal to 1 on  $B_\varepsilon$  and 0 outside  $B_{2\varepsilon}$ . In this way we rewrite (4.5) as

$$\begin{aligned} u_i^n(t, x) &= u_i^n(0, x) + (a_\varepsilon K_i) * (\omega^n(t, \cdot) - \omega^n(0, \cdot))(x) + [(a - a_\varepsilon)K_i] * (\omega^n(t, \cdot) - \omega^n(0, \cdot))(x) \\ &\quad - \int_0^t (\nabla \nabla^\perp [(1-a)K_i]) \star (u^n(s, \cdot) \otimes u^n(s, \cdot))(x) ds. \end{aligned}$$

We can prove that  $u^n$  is a Cauchy sequence using the previous formula. We consider  $u^n, u^m$  with  $n, m \in \mathbb{N}$ . By linearity of the convolution we have that  $u^n - u^m$  satisfies the following

$$\begin{aligned}
 u_i^n(t, x) - u_i^m(t, x) &= \underbrace{u_i^n(0, x) - u_i^m(0, x)}_{(I)} \\
 &+ \underbrace{(a_\varepsilon K_i) * (\omega^n(t, \cdot) - \omega^m(t, \cdot))}_{(II)}(x) + \underbrace{(a_\varepsilon K_i) * (\omega_0^m - \omega_0^n)}_{(III)}(x) \\
 &+ \underbrace{((a - a_\varepsilon)K_i) * (\omega^n(t, \cdot) - \omega^m(t, \cdot))}_{(IV)}(x) + \underbrace{((a - a_\varepsilon)K_i) * (\omega_0^m - \omega_0^n)}_{(V)}(x) \\
 &- \int_0^t \underbrace{(\nabla \nabla^\perp [(1 - a)K_i]) \star (u^n(s, \cdot) \otimes u^n(s, \cdot) - u^m(s, \cdot) \otimes u^m(s, \cdot))}_{(VI)}(x) \, ds. \tag{4.6}
 \end{aligned}$$

In order to prove that  $u^n$  is a Cauchy sequence, we fix a parameter  $\eta > 0$  and we will estimate all the terms in (4.6). First of all, since the initial datum  $u_0^n$  converges strongly in  $L^2$ , it is obvious that there exists  $N_1$  such that  $\forall n, m > N_1$

$$\|u_i^n(0, \cdot) - u_i^m(0, \cdot)\|_{L^2} < \eta.$$

Step 2 *Estimate on (VI).*

By Young’s convolution inequality we have that

$$\begin{aligned}
 &\|\nabla \nabla^\perp [(1 - a)K] \star (u^n(s, \cdot) \otimes u^n(s, \cdot) - u^m(s, \cdot) \otimes u^m(s, \cdot))\|_{L^2} \\
 &\leq \|\nabla \nabla^\perp [(1 - a)K]\|_{L^2} \underbrace{\|u^n(s, \cdot) \otimes u^n(s, \cdot) - u^m(s, \cdot) \otimes u^m(s, \cdot)\|_{L^1}}_{(VI^*)}. \tag{4.7}
 \end{aligned}$$

We add and subtract  $u^n(s, \cdot) \otimes u^m(s, \cdot)$  in (VI\*) and by Hölder inequality we have

$$\begin{aligned}
 &\|u^n(s, \cdot) \otimes u^n(s, \cdot) - u^n(s, \cdot) \otimes u^m(s, \cdot)\|_{L^1} \\
 &\leq (\|u^n(t, \cdot)\|_{L^2} + \|u^m(t, \cdot)\|_{L^2}) \|u^n(s, \cdot) - u^m(s, \cdot)\|_{L^2}.
 \end{aligned}$$

For the first factor in (4.7) we have that

$$\nabla \nabla^\perp [(1 - a)K_i] = -(\nabla \nabla^\perp a)K_i - \nabla^\perp a \nabla K_i - \nabla a \nabla^\perp K_i + (1 - a) \nabla \nabla^\perp K_i,$$

and it is easy to see that each term on the right-hand side has uniformly bounded  $L^2$  norm. Then we have that

$$\begin{aligned}
 &\int_0^t \|\nabla \nabla^\perp [(1 - a)K] \star (u^n(s, \cdot) \otimes u^n(s, \cdot) - u^m(s, \cdot) \otimes u^m(s, \cdot))\|_{L^2} \, ds \\
 &\leq C \|u_0\|_{L^2} \int_0^t \|u^n(s, \cdot) - u^m(s, \cdot)\|_{L^2} \, ds. \tag{4.8}
 \end{aligned}$$

Step 3 *Estimate on (II) and (III).*

For simplicity we will estimate (III), but it will be clear from the proof that by using the uniform estimates proved in Section 3 the same estimate holds true for (II). We compute

$$\|(a_\varepsilon K_i) * (\omega_0^n - \omega_0^m)\|_{L^2}^2 = \int_{\mathbb{R}^2} \left| \int_{B_{2\varepsilon}(x)} a_\varepsilon(x - y) K_i(x - y) (\omega_0^n(y) - \omega_0^m(y)) \, dy \right|^2 \, dx$$

$$\begin{aligned}
 &\leq \int_{\mathbb{R}^2} \left( \int_{B_{2\varepsilon}(x)} \frac{1}{|x-y|} |\omega_0^n(y) - \omega_0^m(y)| dy \right)^2 dx \\
 &= \int_{\mathbb{R}^2} \left( \int_{B_{2\varepsilon}(x)} \frac{1}{|x-y|(\log(1/|x-y|))^\alpha} \sqrt{|\omega_0^n(y) - \omega_0^m(y)|(\log(e + |\omega_0^n(y) - \omega_0^m(y)|))^\alpha} \right. \\
 &\quad \left. \times \left( \log \left( \frac{1}{|x-y|} \right) \right)^\alpha \sqrt{\frac{|\omega_0^n(y) - \omega_0^m(y)|}{(\log(e + |\omega_0^n(y) - \omega_0^m(y)|))^\alpha}} dy \right)^2 dx \\
 &\leq \int_{\mathbb{R}^2} \int_{B_{2\varepsilon}(x)} \frac{1}{|x-y|^2(\log(1/|x-y|))^{2\alpha}} |\omega_0^n(y) - \omega_0^m(y)|(\log(e + |\omega_0^n(y) - \omega_0^m(y)|))^\alpha dy \\
 &\quad \times \int_{B_{2\varepsilon}(x)} \left( \log \left( \frac{1}{|x-y|} \right) \right)^{2\alpha} \frac{|\omega_0^n(y) - \omega_0^m(y)|}{(\log(e + |\omega_0^n(y) - \omega_0^m(y)|))^\alpha} dy dx \\
 &\leq \underbrace{\sup_x \int_{B_{2\varepsilon}(x)} \left( \log \left( \frac{1}{|x-y|} \right) \right)^{2\alpha} \frac{|\omega_0^n(y) - \omega_0^m(y)|}{(\log(e + |\omega_0^n(y) - \omega_0^m(y)|))^\alpha} dy}_{(I^*)} \\
 &\quad \times \underbrace{\int_{\mathbb{R}^2} \int_{B_{2\varepsilon}(x)} \frac{1}{|x-y|^2(\log(1/|x-y|))^{2\alpha}} |\omega_0^n(y) - \omega_0^m(y)|(\log(e + |\omega_0^n(y) - \omega_0^m(y)|))^\alpha dy dx}_{(II^*)}.
 \end{aligned}$$

We estimate  $(I^*)$  and  $(II^*)$  separately. Let  $\beta$  be as in (2.3) and

$$g_\varepsilon(x) = \chi_{B_{2\varepsilon}}(x) \frac{1}{|x|^2(\log(1/|x|))^{2\alpha}},$$

we have that

$$(II^*) = \|g_\varepsilon * \beta(|\omega_0^n - \omega_0^m|)\|_{L^1} \leq \|g_\varepsilon\|_{L^1} \|\beta(|\omega_0^n - \omega_0^m|)\|_{L^1}. \tag{4.9}$$

By using the convexity of  $\beta$  and Lemma 3.1, we have that

$$\|\beta(|\omega_0^n - \omega_0^m|)\|_{L^1} \leq C,$$

where  $C$  is independent from  $n, m$ , while for  $\alpha > 1/2$

$$\|g_\varepsilon\|_{L^1} = \frac{C}{(\log(1/\varepsilon))^{2\alpha-1}}, \tag{4.10}$$

which can be made as small as we want by choosing  $\varepsilon$  properly. For  $(I^*)$  we use the following facts on the Legendre transform. The maximum of  $st - \beta(t)$  occurs at a point  $t$  where  $s \geq (\log(e+t))^{2\alpha}$ , that is, where  $t_*(s) \leq e^{s/(2\alpha)}$ , so that, by denoting with  $\beta^*$  the Legendre transform of  $\beta$ , we have that  $\beta^*(s) \leq se^{s/(2\alpha)}$ . We apply the inequality

$$st \leq \beta(t) + \beta^*(s) \leq se^{s/(2\alpha)} + t(\log(e+t))^{2\alpha},$$

to  $s = \left(\log\left(\frac{1}{|x-y|}\right)\right)^{2\alpha}$  and  $t = \frac{|\omega_0^n - \omega_0^m|}{(\log(e + |\omega_0^n - \omega_0^m|))^\alpha}$  and we find that  $(I^*)$  is bounded by

$$\begin{aligned}
 (I^*) &\leq \sup_x \left\{ \int_{B_{2\varepsilon}} \frac{(\log(1/|z|))^{2\alpha}}{|z|} dz \right. \\
 &\quad \left. + \int_{B_{2\varepsilon}(x)} \underbrace{\frac{|\omega_0^n(y) - \omega_0^m(y)|}{(\log(e + |\omega_0^n(y) - \omega_0^m(y)|))^\alpha} \log^2 \left( e + \frac{|\omega_0^n(y) - \omega_0^m(y)|}{(\log(e + |\omega_0^n(y) - \omega_0^m(y)|))^\alpha} \right)}_{(I^{**})} dy \right\},
 \end{aligned}$$

and we can estimate  $(I^{**})$  by

$$(I^{**}) \leq |\omega_0^n - \omega_0^m| (\log(e + |\omega_0^n - \omega_0^m|))^\alpha,$$

so that  $(I^*)$  is finite using the properties of the function  $t \mapsto t(\log(e+t))^\alpha$  together with Lemma 3.1. So, by decreasing  $\varepsilon$  if necessary we get that

$$(II) + (III) \leq C\eta.$$

Step 4 *Estimates on (IV) and (V).*

In the previous step we have fixed the constant  $\varepsilon$ , so by applying Young’s inequality to

$$\|[(a - a_\varepsilon)K_i] * (\omega_0^n - \omega_0^m)\|_{L^2} \leq \|(a - a_\varepsilon)K_i\|_{L^2} \|\omega_0^n - \omega_0^m\|_{L^1} \leq C(\varepsilon) \|\omega_0^n - \omega_0^m\|_{L^1},$$

$$\begin{aligned} \|[(a - a_\varepsilon)K_i] * (\omega^n(t, \cdot) - \omega^m(t, \cdot))\|_{L^2} &\leq \|(a - a_\varepsilon)K_i\|_{L^2} \|\omega^n(t, \cdot) - \omega^m(t, \cdot)\|_{L^1} \\ &\leq C(\varepsilon) \|\omega^n(t, \cdot) - \omega^m(t, \cdot)\|_{L^1}, \end{aligned}$$

where  $C(\varepsilon)$  blows up as  $\varepsilon \rightarrow 0$ . Now,  $\varepsilon = \varepsilon(\eta)$  has been fixed in the previous step and by Remark 2.3 the vorticity converges strongly in  $C([0, T], L^1(\mathbb{R}^2))$ . Then, we have that there exists  $N_2$  such that  $\forall n, m > N_2$

$$\|\omega_0^n - \omega_0^m\|_{L^1}, \sup_{t \in (0, T)} \|\omega^n(t, \cdot) - \omega^m(t, \cdot)\|_{L^1} < \eta / C(\varepsilon).$$

Step 5  *$u^n$  is a Cauchy sequence in  $C([0, T]; L^2(\mathbb{R}^2))$ .*

By collecting all the estimates obtained in the previous steps we get that for all  $n, m > N := \max\{N_1, N_2\}$

$$\|u^n(t, \cdot) - u^m(t, \cdot)\|_{L^2} \leq C \left( \eta + \int_0^t \|u^n(s, \cdot) - u^m(s, \cdot)\|_{L^2} ds \right), \tag{4.11}$$

and by Gronwall’s lemma

$$\|u^n(t, \cdot) - u^m(t, \cdot)\|_{L^2} \leq C(T)\eta. \tag{4.12}$$

Taking the supremum in time in (4.12) we have the result.

Step 6 *Conservation of the energy.*

Since  $u^n$  is an exact smooth solution and smooth solutions are conservative, we have that

$$\|u^n(t, \cdot)\|_{L^2} = \|u_0^n\|_{L^2}. \tag{4.13}$$

Then, since  $u^n$  converges strongly to  $u$  in  $C([0, T]; L^2(\mathbb{R}^2))$ , by letting  $n \rightarrow \infty$  in (4.13) we have the result. □

REMARK 4.1. The  $L(\log L)^\alpha$  integrability of the vorticity plays a role in the proof of the Theorem 4.1 only in Step 3, allowing us to make the quantities (II) and (III) as small as we want. For  $\alpha < 1/2$  this cannot be achieved: Let  $\omega_0$  be the vorticity in Example 2.1 and decompose the corresponding velocity field  $u_0$  as

$$u_0 = (a_\varepsilon K) * \omega_0 + [(1 - a_\varepsilon)K] * \omega_0.$$

Since  $(1 - a_\varepsilon)K \in L^\infty(\mathbb{R}^2)$  and  $\omega_0 \in L^1(\mathbb{R}^2)$ , it follows that  $[(1 - a_\varepsilon)K] * \omega_0$  is bounded, thus in  $L^2_{\text{loc}}(\mathbb{R}^2)$ . But  $u_0$  is not locally square integrable and then necessarily  $(a_\varepsilon K) * \omega_0 \notin L^2_{\text{loc}}(\mathbb{R}^2)$  as well.

Now we deal with the vanishing viscosity method.

**THEOREM 4.2.** *Let  $\omega_0 \in L(\log L)_c^\alpha(\mathbb{R}^2)$ , with  $\alpha > 1/2$ , with zero total mass. Let  $u$  be a weak solution of (1.1), with  $\text{curl} u_0 = \omega_0$ , that can be obtained as a limit of a sequence  $u^\nu$  constructed via (VV). Then,  $u^\nu$  satisfies the following convergence*

$$u^\nu \rightarrow u \quad \text{in } C([0, T]; L^2(\mathbb{R}^2)), \tag{4.14}$$

and  $u$  is conservative.

*Proof.* Since the parameter  $\nu$  is supposed to vary over a countable set, given the sequence  $\nu_n \rightarrow 0$ , we denote with  $u^n$  and  $\omega^n$  the sequences  $u^{\nu_n}$  and  $\omega^{\nu_n}$ . Thanks to Remark 2.4, it is enough to prove that  $u^n$  is a Cauchy sequence in  $C([0, T]; L^2(\mathbb{R}^2))$ . We proceed as in the proof of Theorem 4.1. The only difference is that an error term appears in the Serfati identity, which is

$$\int_0^t (\Delta[(1 - a)K_i]) * (\nu_n \omega^n(s, \cdot) - \nu_m \omega^m(s, \cdot)) \, ds. \tag{4.15}$$

By Young’s inequality we have that

$$\begin{aligned} & \|(\Delta[(1 - a)K_i]) * (\nu_n \omega^n(s, \cdot) - \nu_m \omega^m(s, \cdot))\|_{L^2} \\ & \leq \nu_n \|\Delta[(1 - a)K_i]\|_{L^2} \|\omega^n(s, \cdot) - \omega^m(s, \cdot)\|_{L^1} \\ & \quad + |\nu_m - \nu_n| \|\Delta[(1 - a)K_i]\|_{L^2} \|\omega^m(s, \cdot)\|_{L^1}. \end{aligned}$$

Since  $\Delta K_i$  is in  $L^2(B_1^c)$ , a straightforward computation shows that  $\Delta[(1 - a)K]$  is bounded in  $L^2$ . So, because of Remark 2.3, there exists  $N_3$  such that for all  $n, m > N_3$  we have that

$$\|(\Delta[(1 - a)K]) * (\nu_n \omega^n(s, \cdot) - \nu_m \omega^m(s, \cdot))\|_{L^2} \leq C\eta,$$

and this concludes the proof. □

Finally we deal with the vortex-blob method. The theorem is the following.

**THEOREM 4.3.** *Let  $\omega_0 \in L(\log L)_c^\alpha(\mathbb{R}^2)$ , with  $\alpha > 1/2$ , with zero total mass. Let  $u$  be a weak solution of (1.1), with  $\text{curl} u_0 = \omega_0$ , that can be obtained as the limit of a sequence  $u^\varepsilon$  constructed via (VB). Then,  $u^\varepsilon$  satisfies the following convergence*

$$u^\varepsilon \rightarrow u \quad \text{in } C([0, T]; L^2(\mathbb{R}^2)), \tag{4.16}$$

and  $u$  is conservative.

*Proof.* Since the parameter  $\varepsilon$  is supposed to vary over a countable set, given the sequence  $\varepsilon_n \rightarrow 0$ , we denote with  $u^n$  and  $\omega^n$  the sequences  $u^{\varepsilon_n}$  and  $\omega^{\varepsilon_n}$ . We divide the proof in several steps.

Step 1  $u^n$  is a Cauchy sequence in  $C([0, T]; L^2(\mathbb{R}^2))$ .

We proceed as in the proof of Theorem 4.1. The only difference is that an error term appears in the Serfati identity, which is

$$\int_0^t ((\nabla[(1 - a)K_i]) \star (F_n(s, \cdot) - F_m(s, \cdot)))(x) \, ds. \tag{4.17}$$

Since  $\nabla[(1-a)K_i] \in L^2(\mathbb{R}^2)$ , by using Young’s inequality we get that

$$\|((\nabla[(1-a)K_i]) \star (F_n(s, \cdot) - F_m(s, \cdot)))\|_{L^2} \leq \|\nabla[(1-a)K_i]\|_{L^2} \|F_n(s, \cdot) - F_m(s, \cdot)\|_{L^1},$$

which can be made as small as we want because of Lemma A.2.

Step 2 *Conservation of the energy.*

We prove now that  $u$  is a conservative weak solution. With our notations, multiplying (2.8) by  $u^n$  and integrating in space and time we have that

$$\int_{\mathbb{R}^2} |u^n|^2(t, x) dx = \int_{\mathbb{R}^2} |u^n|^2(0, x) dx - \int_0^t \int_{\mathbb{R}^2} (\nabla K \star F_n) \cdot u^n dx. \tag{4.18}$$

For the second term on the right-hand side, by Lemma A.2 we have that

$$\begin{aligned} \left| \int_0^t \int_{\mathbb{R}^2} (\nabla K \star F_n) \cdot u^n dx \right| &\leq \|\nabla K \star F_n(s, \cdot)\|_{L^2} \|u^n(s, \cdot)\|_{L^2} \\ &\leq \|F_n(s, \cdot)\|_{L^2} \|u^n(s, \cdot)\|_{L^2} \\ &\leq C(\delta_n)^{-\frac{7}{3}} (\varepsilon_n)^{\frac{1}{3}}, \end{aligned}$$

which goes to 0 as  $\varepsilon_n \rightarrow 0$ . Then, by the convergence (4.16) letting  $\varepsilon_n \rightarrow 0$  in (4.18) we have that

$$\int_{\mathbb{R}^2} |u|^2(t, x) dx = \int_{\mathbb{R}^2} |u_0|^2(x) dx,$$

which gives the result. □

**REMARK 4.2.** In the case of a vorticity  $\omega_0 \in L_c(\log L)^{1/2}(\mathbb{R}^2)$ , the  $L^2$ -compactness of a sequence of approximating velocities depends on the approximation method. This is the case of the methods examined in this work and it follows from the a priori estimates of Section 3 and the compactness arguments of [8]. However, it does not seem possible to find explicit quantitative rates of convergence by resorting to the Serfati identity, because of the quantities (II) and (III) in Step 3 of Theorem 4.1. Even if that were the case, those rates would implicitly depend on the specific choice of  $\omega_0$  and not on its norm.

**Acknowledgments.** The author gratefully acknowledges useful discussions with Gianluca Crippa and Stefano Spirito. The author thanks the anonymous referees for several helpful comments. This work was started while the author was a PostDoc at the Departement Mathematik und Informatik of the Universität Basel. This research has been partially supported by the ERC Starting Grant 676675 FLIRT, by the Basque Government under program BCAM-BERC 2018-2021, and by the Spanish Ministry of Science, Innovation and Universities under the BCAM Severo Ochoa accreditation SEV-2017-0718.

**Appendix. The vortex-blob method.** In this appendix we describe the vortex-blob approximation and some of its properties. Let us consider an initial vorticity  $\omega_0 \in L_c^p(\mathbb{R}^2)$  with  $1 \leq p \leq \infty$ . Let  $\varepsilon \in (0, 1)$ , we consider two small parameters in  $(0, 1)$ , which later will be chosen as functions of  $\varepsilon$ , denoted by  $\delta(\varepsilon)$  and  $h(\varepsilon)$ .

First of all, we consider the lattice

$$\Lambda_h := \{\alpha_i \in h\mathbb{Z} \times h\mathbb{Z} : \alpha_i = h(i_1, i_2), \text{ where } i_1, i_2 \in \mathbb{Z}\},$$

and define  $R_i$  as the square with sides of length  $h$  parallel to the coordinate axis and centered at  $\alpha_i \in \Lambda_h$ . Let  $\rho_\delta$  be a standard mollifier and define

$$\omega_0^\varepsilon := \omega_0 * \rho_{\delta(\varepsilon)}. \tag{A.1}$$

For any  $\delta \in (0,1)$  the support of  $\omega_0^\varepsilon$  is contained in a fixed compact set in  $\mathbb{R}^2$ , then it can be tiled by a finite number  $N(\varepsilon)$  of squares  $R_i$ . Define the quantities

$$\Gamma_i^\varepsilon = \int_{R_i} \omega_0^\varepsilon(x) \, dx, \quad \text{for } i = 1, \dots, N(\varepsilon).$$

Let  $\varphi_\varepsilon$  be another mollifier, we define the approximate vorticity to be

$$\omega^\varepsilon(t, x) = \sum_{i=1}^{N(\varepsilon)} \Gamma_i^\varepsilon \varphi_\varepsilon(x - X_i^\varepsilon(t)), \tag{A.2}$$

where  $\{X_i^\varepsilon(t)\}_{i=1}^{N(\varepsilon)}$  is a solution of the O.D.E. system

$$\begin{cases} \dot{X}_i^\varepsilon(t) = u^\varepsilon(t, X_i^\varepsilon(t)), \\ X_i^\varepsilon(0) = \alpha_i, \end{cases} \tag{A.3}$$

with  $u^\varepsilon$  defined as

$$u^\varepsilon(t, x) = K * \omega^\varepsilon(t, x) = \sum_{i=1}^{N(\varepsilon)} \Gamma_i^\varepsilon K_\varepsilon(x - X_i^\varepsilon(t)), \tag{A.4}$$

where  $K_\varepsilon = K * \varphi_\varepsilon$ . Note that, since  $\delta$  and  $h$  are  $\varepsilon$ -dependent, we only use the superscript, or subscript,  $\varepsilon$ . The ordinary differential Equations (A.3) are known as the *vortex-blob approximation*. In particular, the approximation of the initial vorticity and the initial velocity are given by

$$\omega^\varepsilon(0, x) = \sum_{i=1}^{N(\varepsilon)} \Gamma_i^\varepsilon \varphi_\varepsilon(x - \alpha_i), \quad u^\varepsilon(0, x) = \sum_{i=1}^{N(\varepsilon)} \Gamma_i^\varepsilon K_\varepsilon(x - \alpha_i). \tag{A.5}$$

It is not difficult to show the bound (see [16])

$$\sup_{t \in (0, T)} (\|u^\varepsilon(t, \cdot)\|_{L^\infty} + \|\nabla u^\varepsilon(t, \cdot)\|_{L^\infty}) \leq \frac{C}{\varepsilon^2}. \tag{A.6}$$

From (A.6) it follows that, for every fixed  $\varepsilon > 0$ , there exists a unique smooth solution  $\{X_i^\varepsilon(t)\}_{i=1}^{N(\varepsilon)}$  of the O.D.E. system (A.3), which implies that  $u^\varepsilon$  and  $\omega^\varepsilon$  are well-defined smooth functions. Note that  $u^\varepsilon$  and  $\omega^\varepsilon$  are not exact solutions of the Euler equations. Precisely, the approximate vorticity  $\omega^\varepsilon$  satisfies the following equation

$$\partial_t \omega^\varepsilon + u^\varepsilon \cdot \nabla \omega^\varepsilon = E_\varepsilon, \tag{A.7}$$

where, by a direct computation the error term is given by

$$E_\varepsilon(t, x) := \sum_{i=1}^{N(\varepsilon)} [u^\varepsilon(t, x) - u^\varepsilon(t, X_i^\varepsilon(t))] \cdot \nabla \varphi_\varepsilon(x - X_i^\varepsilon(t)) \Gamma_i^\varepsilon. \tag{A.8}$$

Concerning the approximate velocity  $u^\varepsilon$ , consider the quantity

$$w^\varepsilon = \partial_t u^\varepsilon + (u^\varepsilon \cdot \nabla) u^\varepsilon.$$

Since  $w^\varepsilon$  satisfies the system

$$\begin{cases} \operatorname{curl} w^\varepsilon = E_\varepsilon, \\ \operatorname{div} w^\varepsilon = \operatorname{div} \operatorname{div} (u^\varepsilon \otimes u^\varepsilon), \end{cases} \tag{A.9}$$

we derive that there exists a function  $p^\varepsilon$  such that

$$-\Delta p^\varepsilon = \operatorname{div} \operatorname{div} (u^\varepsilon \otimes u^\varepsilon),$$

and

$$w^\varepsilon = -\nabla p^\varepsilon + K * E_\varepsilon.$$

Then, the velocity given by the vortex-blob approximation verifies the following equations

$$\begin{cases} \partial_t u^\varepsilon + (u^\varepsilon \cdot \nabla) u^\varepsilon + \nabla p^\varepsilon = K * E_\varepsilon, \\ \operatorname{div} u^\varepsilon = 0. \end{cases} \tag{A.10}$$

Since  $u^\varepsilon$  is divergence-free,  $E_\varepsilon$  can be rewritten as  $E_\varepsilon(t, x) = \operatorname{div} F_\varepsilon(t, x)$  where

$$F_\varepsilon(t, x) := \sum_{i=1}^{N(\varepsilon)} [u^\varepsilon(t, x) - u^\varepsilon(t, X_i^\varepsilon(t))] \varphi_\varepsilon(x - X_i^\varepsilon(t)) \Gamma_i^\varepsilon. \tag{A.11}$$

Let  $\bar{\omega}^\varepsilon$  be the solution of the linear transport equation with vector field  $u^\varepsilon$ , that is

$$\begin{cases} \partial_t \bar{\omega}^\varepsilon + u^\varepsilon \cdot \nabla \bar{\omega}^\varepsilon = 0, \\ \bar{\omega}^\varepsilon(0, \cdot) = \omega_0^\varepsilon. \end{cases} \tag{A.12}$$

Since  $u^\varepsilon$  satisfies (A.6), there exists a unique smooth solution  $\bar{\omega}^\varepsilon$ , which is given by the formula

$$\bar{\omega}^\varepsilon(t, x) = \omega_0^\varepsilon((X^\varepsilon)^{-1}(t, \cdot)(x)), \tag{A.13}$$

where  $X^\varepsilon$  is the flow of  $u^\varepsilon$ , that is,

$$\begin{cases} \dot{X}^\varepsilon(t, x) = u^\varepsilon(t, X^\varepsilon(t, x)), \\ X^\varepsilon(0, x) = x. \end{cases} \tag{A.14}$$

Moreover, since  $\operatorname{div} u^\varepsilon = 0$ , we have

$$\|\bar{\omega}^\varepsilon(t, \cdot)\|_{L^p} = \|\omega_0^\varepsilon\|_{L^p} \leq \|\omega_0\|_{L^p}.$$

The following estimates between the  $L^p$  norms of  $\omega^\varepsilon$  and  $\bar{\omega}^\varepsilon$  hold true, see [2, 11].

LEMMA A.1. *Let  $\omega_0 \in L^1(\mathbb{R}^2)$  and let  $h = h(\varepsilon)$  be chosen as*

$$h(\varepsilon) = \frac{\varepsilon^4}{\exp(C_1 \varepsilon^{-2} \|\omega_0\|_{L^1 T})}, \tag{A.15}$$

where  $C_1 > 0$  is a positive constant. Then, the estimate

$$\sup_{0 \leq t \leq T} \|\omega^\varepsilon - \varphi_\varepsilon * \bar{\omega}^\varepsilon\|_{L^p} \leq C\varepsilon^{1+\frac{2}{p}} \quad (\text{A.16})$$

holds for all  $1 \leq p \leq \infty$ , where  $C > 0$  is a positive constant which does not depend on  $\varepsilon$ .

Moreover, with a suitable choice of the parameters in the definition of the vortex-blob method we also have that the error term  $F_\varepsilon$  goes to 0 in the limit, see [2].

LEMMA A.2. *Let  $\omega_0 \in L^p_c(\mathbb{R}^2)$  with  $p \geq 1$ , then the quantity  $F_\varepsilon$  defined in (A.11) satisfies*

$$\sup_{t \in [0, T]} \|F_\varepsilon(t, \cdot)\|_{L^1} \rightarrow 0, \quad \text{as } \varepsilon \rightarrow 0. \quad (\text{A.17})$$

Moreover, choosing  $h(\varepsilon) = C_1\varepsilon^6 \exp(-C_0\varepsilon^{-2})$  where  $C_1, C_0$  are positive constants, we have that  $F_\varepsilon$  satisfies the following additional bound

$$\|F_\varepsilon(t, \cdot)\|_{L^2} \leq C\delta^{-\beta} \varepsilon^{\frac{7}{3}} \|\omega_0\|_{L^1},$$

which goes to 0 choosing  $\delta$  as above and  $0 < \sigma < 1/7$ .

Finally, by showing the equi-integrability of the sequence  $\omega^\varepsilon$  one of the main results in [11] is the following.

THEOREM A.1. *Let  $\omega_0 \in L^1_c(\mathbb{R}^2)$  and  $\omega_0^\varepsilon$  defined as (A.1). Then the sequence  $\omega^\varepsilon$  as in (A.2) is equi-integrable in  $L^1((0, T) \times \mathbb{R}^2)$ . Moreover, there exists a function  $\omega \in C([0, T]; L^1(\mathbb{R}^2))$  such that, along a sub-sequence,*

$$\omega^\varepsilon \rightarrow \omega \quad \text{in } C([0, T]; L^1(\mathbb{R}^2)),$$

where  $\omega$  is a renormalized and Lagrangian solution of the two-dimensional Euler equations.

## REFERENCES

- [1] D.M. Ambrose, J.P. Kelliher, M.C. Lopes Filho, and H.J. Nussenzweig Lopes, *Serfati solutions to the 2D Euler equations on exterior domain*, J. Differ. Equ., **259:4509–4560**, 2015. [1](#), [4](#)
- [2] J.T. Beale, *The approximation of weak solutions to the 2-D Euler equations by vortex elements*, in J. Glimm and A.J. Majda (eds.), *Multidimensional Hyperbolic Problems and Computations*, IMA Vol. Math. Appl., **29:23–37**, 1991. [1](#), [2.2](#), [4](#), [4](#)
- [3] C. Bennett and R. Sharpley, *Interpolation of Operators*, Pure and Applied Mathematics, Academic Press, **129**, 1988. [2.1](#)
- [4] A. Bohun, F. Bouchut, and G. Crippa, *Lagrangian solutions to the 2D Euler system with  $L^1$  vorticity and infinite energy*, Nonlinear Anal. Theory Meth. Appl., **132:160–172**, 2016. [1](#), [2.2](#)
- [5] A. Bressan and R. Murray, *On self-similar solutions to the incompressible Euler equations*, J. Differ. Equ., **269:5142–5203**, 2020. [1](#)
- [6] A. Bressan and W. Shen, *A posteriori error estimates for self-similar solutions to the Euler equations*, Discrete Contin. Dyn. Syst. A, **41:113–130**, 2021. [1](#)
- [7] D. Chae, *Weak solutions of the 2-D Euler equations with initial vorticity in  $L(\log L)$* , J. Differ. Equ., **103:323–337**, 1993. [1](#), [2.2](#)
- [8] D. Chae, *Weak solutions of the 2-D incompressible Euler equations*, Nonlinear Anal. Theory Meth. Appl., **23:629–638**, 1994. [1](#), [2.2](#), [2.1](#), [4.2](#)
- [9] A. Cheskidov, M.C. Lopes Filho, H.J. Nussenzweig Lopes, and R. Shvydkoy, *Energy conservation in two-dimensional incompressible ideal fluids*, Commun. Math. Phys., **348:129–143**, 2016. [1](#), [2.3](#)
- [10] G. Ciampa, G. Crippa, and S. Spirito, *Strong convergence of the vorticity for the 2D Euler equations in the inviscid limit*, Arch. Ration. Mech. Anal., **240:295–326**, 2021. [1](#), [2.2](#), [2.3](#)

- [11] G. Ciampa, G. Crippa, and S. Spirito, *Weak solutions obtained by the vortex method for the 2D Euler equations are Lagrangian and conserve the energy*, J. Nonlinear Sci., **30**:2787–2820, 2020. [1](#), [2.2](#), [2.3](#), [4](#), [4](#)
- [12] P. Constantin, T.D. Drivas, and T.M. Elgindi, *Inviscid limit of vorticity distributions in Yudovich class*, Comm. Pure Appl. Math., **75**(1):60–82, 2022. [2.2](#)
- [13] G. Crippa, C. Nobili, C. Seis, and S. Spirito, *Eulerian and Lagrangian solutions to the continuity and Euler equations with  $L^1$  vorticity*, SIAM J. Math. Anal., **49**(5):3973–3998, 2017. [1](#)
- [14] G. Crippa and S. Spirito, *Renormalized solutions of the 2D Euler equations*, Commun. Math. Phys., **339**:191–198, 2015. [1](#)
- [15] J.M. Delort, *Existence de nappes de tourbillon en dimension deux*, J. Amer. Math. Soc., **4**:553–586, 1991. [1](#)
- [16] R.J. DiPerna and A. Majda, *Concentrations in regularizations for 2-D incompressible flow*, Comm. Pure Appl. Math., **40**:301–345, 1987. [1](#), [2](#), [2.2](#), [2.2](#), [2.2](#), [2.1](#), [4](#)
- [17] S. Lanthaler, S. Mishra, and C. Parés-Pulido, *On the conservation of energy in two-dimensional incompressible flows*, Nonlinearity, **34**:1084–1135, 2021. [1](#), [2.3](#)
- [18] L. Lichtenstein, *Über einige Existenzproblem der Hydrodynamik homogener, unzusammendrückbarer, reibungsloser Flüssigkeiten und die Helmholtzschen Wirbelsätze*, Math. Z., **23**:89–154, 1925. [1](#)
- [19] M.C. Lopes Filho, H.J. Nussenzweig Lopes, and E. Tadmor, *Approximate solutions of the incompressible Euler equations with no concentrations*, Ann. Inst. H. Poincaré Anal. Non Linéaire, **17**(3):317–412, 2000. [1](#), [2.3](#)
- [20] A.J. Majda and A.L. Bertozzi, *Vorticity and Incompressible Flow*, Cambridge Texts in Applied Mathematics, Cambridge Univ. Press, London/New York, **27**, 2002. [2.3](#)
- [21] F. Mengual and L. Székelyhidi Jr., *Dissipative Euler flows for vortex sheet initial data without distinguished sign*, Comm. Pure Appl. Math., 2022. [1](#)
- [22] H.J. Nussenzweig Lopes, C. Seis, and E. Wiedemann, *On the vanishing viscosity limit for 2D incompressible flows with unbounded vorticity*, Nonlinearity, **34**:3112–3121, 2021. [2.2](#)
- [23] P. Serfati, *Solutions  $C^\infty$  en temps,  $n$ -log Lipschitz bornées en espace et équation d'Euler*, C.R. Acad. Sci. Paris Sér. I Math., **320**:555–558, 1995. [1](#), [4](#)
- [24] S. Schochet, *The point-vortex method for periodic weak solutions of the 2-D Euler equations*, Comm. Pure Appl. Math., **49**:911–965, 1996. [1](#), [2.2](#), [2.1](#)
- [25] I. Vecchi and S.J. Wu, *On  $L^1$ -vorticity for 2-D incompressible flow*, Manuscripta Math., **78**(4):403–412, 1993. [1](#)
- [26] M. Vishik, *Instability and non-uniqueness in the Cauchy problem for the Euler equations of an ideal incompressible fluid. Part I*, arXiv preprint, [arXiv:1805.09426](#), 2018. [1](#)
- [27] M. Vishik, *Instability and non-uniqueness in the Cauchy problem for the Euler equations of an ideal incompressible fluid. Part II*, arXiv preprint, [arXiv:1805.09440](#), 2018. [1](#)
- [28] W. Wolibner, *Un théorème sur l'existence du mouvement plan d'un fluide parfait, homogène, incompressible, pendant un temps infiniment long*, Math. Z., **37**:698–726, 1933. [1](#)
- [29] V.I. Yudovich, *Non-stationary flows of an ideal incompressible fluid*, Ž. Vychisl. Mat. i Mat. Fiz., **3**:1032–1066, 1963. [1](#)