

Generating sets and representability for symplectic capacities

DUŠAN JOKSIMOVIĆ AND FABIAN ZILTENER

K. Cieliebak, H. Hofer, J. Latschev, and F. Schlenk (CHLS) posed the problem of finding a minimal generating set for the (symplectic) capacities on a given symplectic category. We show that if the category contains a certain one-parameter family of objects, then every countably Borel-generating set of (normalized) capacities has cardinality (strictly) bigger than the continuum. This appears to be the first result regarding the problem of CHLS, except for two results of D. McDuff about the category of ellipsoids in dimension 4.

We also prove that every finitely differentially generating set of capacities on a given symplectic category is uncountable, provided that the category contains a one-parameter family of symplectic manifolds that is “strictly volume-increasing” and “embedding-capacity-wise constant”. It follows that the Ekeland-Hofer capacities and the volume capacity do not finitely differentially generate all generalized capacities on the category of ellipsoids. This answers a variant of a question of CHLS.

In addition, we prove that if a given symplectic category contains a certain one-parameter family of objects, then almost no normalized capacity is domain- or target-representable. This provides some solutions to two central problems of CHLS.

1	Introduction	838
2	Main results and applications	848
3	Proof of Theorem 17(i,ii) (cardinality of the set of capacities)	865

D. Joksimović’s work on this article was funded by NWO (The Netherlands Organisation for Scientific Research) via the TOP grant 614.001.508. F. Ziltener worked on this article during a visit at FIM - the Institute for Mathematical Research at ETH Zürich. He gratefully acknowledges the hospitality of ETH.

4	Proof of Theorem 43 (cardinality of the set of capacities, more general setting)	877
5	Proof of Proposition 44 (sufficient conditions for being an I-collection)	890
6	Proof of Theorem 17(iii) (cardinality of a generating set)	895
7	Proof of Theorem 25 (uncountability of every generating set under a mild hypothesis)	900
	Appendix A Cardinality of the set of equivalence classes of pairs of manifolds and forms	901
	Appendix B Proof of Theorem 29 (monotone generation for ellipsoids)	905
	References	908

“The introduction of the cipher 0 or the group concept was general nonsense too, and mathematics was more or less stagnating for thousands of years because nobody was around to take such childish steps ...”

— Alexander Grothendieck

1. Introduction

1.1. The problems and special cases of the main results

In this section we state special cases of the main results, since they seem easier to digest. We postpone the statements of the general results until the next section.¹

Our first main result is concerned with generating sets of capacities on symplectic categories. To explain a special case of this result, let $n \in \mathbb{N}_0 :=$

¹In the spirit of Grothendieck we have aimed at formulating our main results in the greatest possible generality.² The anonymous referee for this article was afraid that this would scare off readers. For this reason we first state special cases of the results.

²Probably we failed miserably in this attempt.

$\{0, 1, \dots\}$. We denote by Symp^{2n} the category of all symplectic manifolds³ of dimension $2n$, with morphisms given by the symplectic embeddings⁴. Recall that a subcategory \mathcal{C}' of a category \mathcal{C} is called *isomorphism-closed (in \mathcal{C})* iff every isomorphism of \mathcal{C} starting at some object of \mathcal{C}' is a morphism of \mathcal{C}' .⁵

Definition ((weak) symplectic category). *A weak symplectic category in dimension $2n$ is a subcategory $\mathcal{C} = (\mathcal{O}, \mathcal{M})$ ⁶ of Symp^{2n} , such that for every $(M, \omega) \in \mathcal{O}$ and every $a \in (0, \infty)$ we have $(M, a\omega) \in \mathcal{O}$. We call such a \mathcal{C} a symplectic category if it is isomorphism-closed.*

Remark. Weak symplectic categories were first defined in [3, 2.1. Definition, p. 5], where the adjective “weak” is left out. We explain the reason for our change in terminology in Remark 11 on p. 849.

We denote by B_r^m (\overline{B}_r^m) the open (closed) ball of radius r around 0 in \mathbb{R}^m , and abbreviate

$$B := B_1^{2n}, \quad Z_r^{2n} := B_r^2 \times \mathbb{R}^{2n-2}, \quad Z := Z_1^{2n}.$$

We equip B_r^{2n} and Z_r^{2n} with the standard symplectic form ω_{st} . Let $\mathcal{C} = (\mathcal{O}, \mathcal{M})$ be a symplectic category.

Definition 1. *A generalized capacity on \mathcal{C} is a function*

$$c : \mathcal{O} \rightarrow [0, \infty]$$

with the following properties:

- (i) **(monotonicity)** *If (M, ω) and (M', ω') are two objects in \mathcal{O} between which there exists a \mathcal{C} -morphism, then*

$$c(M, \omega) \leq c(M', \omega').$$

- (ii) **(conformality)** *For every $(M, \omega) \in \mathcal{O}$ and $a \in (0, \infty)$ we have*

$$c(M, a\omega) = a c(M, \omega).$$

³In this article “manifold” refers to a smooth (C^∞) real finite-dimensional manifold. It is allowed to be disconnected and have boundary.

⁴By an embedding we mean an injective smooth immersion with continuous inverse. We do not impose any condition involving the boundaries of the two manifolds.

⁵In particular, it ends at some object of \mathcal{C}' .

⁶Here \mathcal{O} and \mathcal{M} denote the classes of objects and morphisms of \mathcal{C} , respectively.

Assume now that $B, Z \in \mathcal{O}$. Let c be a generalized capacity on \mathcal{C} . We call c a capacity iff it satisfies:

(iii) (**non-triviality**) $c(B) > 0$ and $c(Z) < \infty$.

We call it normalized iff it satisfies:

(iv) (**normalization**) $c(B) = c(Z) = \pi$.⁷

We denote

$$\text{Cap}(\mathcal{C}) := \{ \text{generalized capacity on } \mathcal{C} \}.$$

Remark. There is a set-theoretic issue with this definition, which we will resolve in Definition 12 in the next section. Compare to Remark 11.

Example 2 (embedding capacities). Let $\mathcal{C} = (\mathcal{O}, \mathcal{M})$ be a symplectic category in dimension $2n$ and (M, ω) an object of Symp^{2n} . We define the *domain-embedding capacity for (M, ω) on \mathcal{C}* to be the function

$$\begin{aligned} c_{M, \omega} &:= c_{M, \omega}^{\mathcal{C}} : \mathcal{O} \rightarrow [0, \infty], \\ c_{M, \omega}(M', \omega') &:= \sup \{ c \in (0, \infty) \mid \exists \text{ symplectic embedding} \\ &\quad (M, c\omega) \rightarrow (M', \omega') \}. \end{aligned}$$

We define the *target-embedding capacity for (M, ω) on \mathcal{C}* to be the function

$$\begin{aligned} c^{M, \omega} &:= c_{\mathcal{C}}^{M, \omega} : \mathcal{O} \rightarrow [0, \infty], \\ c^{M, \omega}(M', \omega') &:= \inf \{ c \in (0, \infty) \mid \exists \text{ symplectic embedding} \\ &\quad (M', \omega') \rightarrow (M, c\omega) \}. \end{aligned}$$

These are generalized capacities.⁸ We define the *Gromov width on \mathcal{C}* to be

$$(1) \quad w := \pi c_{B, \omega_{\text{st}}}^{\mathcal{C}}.$$

If $B, Z \in \mathcal{O}$, then by Gromov's nonsqueezing theorem the Gromov width is a normalized capacity.

⁷In [3, 2.1. Definition, p. 5] only the condition $c(B) = 1$ is imposed here. The second part of our first main result, Theorem 17 below, holds even with our stronger definition.

⁸In the definition on p. 13 in [3] (M, ω) is assumed to be an object of \mathcal{C} , and the morphisms in the definitions of the embedding capacities are asked to be \mathcal{C} -morphisms. However, in [3, Example 2, p. 14] the definition is applied with an (M, ω) that is not an object of \mathcal{C} . In order to make that example work, one needs to allow for Symp^{2n} -morphisms in the definition of the embedding capacities.

Capacities on the category of all symplectic manifolds of a fixed dimension were introduced by I. Ekeland and H. Hofer in [5, 6]. They measure how much a given symplectic manifold does not embed into another one. In [3] the notion of a symplectic capacity on a general weak symplectic category was introduced.⁹ For an overview over symplectic capacities we refer to [3, 13, 14] and references therein.

To define the notion of a CHLS capacity-generating set, let $f : [0, \infty]^\ell \rightarrow [0, \infty]^{\ell'}$ be a map. We call it *homogeneous* iff it is positively 1-homogeneous, i.e., iff $f(ax) = af(x)$ for every $a \in (0, \infty)$ and $x \in [0, \infty]^\ell$.¹⁰ We equip $[0, \infty]^\ell$ with the partial order given by $x \leq y$ iff $x_i \leq y_i$ for every $i \in \{1, \dots, \ell\}$. We call f *monotone* iff it preserves this partial order. As pointed out by the authors of [3] in that article, if $\ell' = 1$, f is homogeneous and monotone, and c_1, \dots, c_ℓ are generalized capacities, then $f \circ (c_1, \dots, c_\ell)$ is again a generalized capacity. Homogeneity and monotonicity are preserved under compositions.

Examples. *The following functions are homogeneous and monotone:*

- *maximum, minimum*
- *For every $a \in [0, \infty)^\ell$ and $p \in \mathbb{R} \setminus \{0\}$ the function*

$$f_{a,p}(x) := \sqrt[p]{\sum_{i=1}^{\ell} a_i x_i^p}.^{11}$$

In the case $a = (\frac{1}{\ell}, \dots, \frac{1}{\ell})$, $p = 1$ the function $f_{a,p}$ is the arithmetic mean, and in the case $a = (\frac{1}{\ell}, \dots, \frac{1}{\ell})$, $p = -1$ it is the harmonic mean.

- *For every $p \in [0, \infty)^\ell$ satisfying $\sum_{i=1}^{\ell} p_i = 1$ the function*

$$x \mapsto \prod_{i=1}^{\ell} x_i^{p_i}.$$

In the case $p = (\frac{1}{\ell}, \dots, \frac{1}{\ell})$ this is the geometric mean.

⁹For the set-theoretic reason mentioned in Remark 11 one needs to ask that the category is isomorphism-closed or small, in order to make sense of this definition.

¹⁰Here we use the convention $a \cdot \infty := \infty$ for every $a \in (0, \infty)$.

¹¹Here we use the conventions $\infty + a = \infty$ for every $a \in [0, \infty]$, $\infty^p = \infty$ for every $p > 0$, and $0^p := \infty$ and $\infty^p := 0$ for every $p < 0$.

Let \mathcal{C} be a symplectic category and \mathcal{G} a subset of $\text{Cap}(\mathcal{C})$. By a *finite homogeneous monotone combination of \mathcal{G}* we mean an expression of the form $f \circ (c_1, \dots, c_\ell)$, where $\ell \in \mathbb{N}_0$, $f : [0, \infty]^\ell \rightarrow [0, \infty]$ is homogeneous and monotone, and $c_1, \dots, c_\ell \in \mathcal{G}$. We define the *set CHLS generated by \mathcal{G}* to be the set of all functions $c : \mathcal{O} \rightarrow [0, \infty]$ that are the pointwise limit of a sequence of finite homogeneous monotone combinations of \mathcal{G} . Since pointwise limits preserve homogeneity and monotonicity, the set CHLS generated by \mathcal{G} consists again of generalized capacities.

By a *CHLS (generalized-)(capacity-)generating set (for \mathcal{C})* we mean a subset \mathcal{G} of $\text{Cap}(\mathcal{C})$, whose CHLS generated set equals $\text{Cap}(\mathcal{C})$.¹² The set CHLS generated by \mathcal{G} is obtained by combining capacities in a lot of ways. One may therefore expect that few capacities suffice to generate all the other capacities. It is tempting to even look for a generating set of capacities that is minimal, in the sense that none of its subsets is generating. This problem was posed by K. Cieliebak, H. Hofer, J. Latschek, and F. Schlenk (CHLS):

Problem 3 ([3], Problem 5, p. 17). *For a given symplectic category find a minimal CHLS capacity-generating set.*

In [3] CHLS also propose more restrictive notions of generating set, for example one in which the only allowed combining functions $f : [0, \infty]^\ell \rightarrow [0, \infty]$ are the minimum and the maximum. We call such a set *limit-min/max (capacity-)generating*. A concrete instance of Problem 3 for this variant of generation is the following.

Question 4. *For a given symplectic category, does there exist a countable¹³ (minimal) limit-min/max capacity-generating set?*

To our knowledge, up to now, Problem 3 and Question 4 have been completely open, except for the following two results of D. McDuff.¹⁴ The first result states that the ECH-capacities are generating in a weaker sense for the category of ellipsoids in dimension 4, see Theorem 29 on p. 865. The second result states that the Ekeland-Hofer capacities together with the volume capacity are not CHLS generating for the restriction category of (open) ellipsoids (as defined on p. 844) in dimension 4. (See [10, Corollary 1.4].)

¹²This notion is introduced in [3, Problem 5, p. 17], where it is just called a “generating system”. (The authors of that article do not explicitly state that \mathcal{G} should be a subset of $\text{Cap}(\mathcal{C})$, but presumably they implicitly ask for this.)

¹³By this we mean finite or countably infinite.

¹⁴There are of course some trivial cases in which Question 4 is easy, e.g. the case in which there are only finitely many \mathcal{C} -isomorphism classes.

By the *continuum* we mean the cardinality of the set of real numbers. The following special case of our first main result (Theorem 17 on p. 854) provides a negative answer to Question 4. In fact, it implies that there is not even a generating set of cardinality at most that of the continuum.

Theorem 5 (cardinality of generating set). *If $n \geq 2$ then every limit-min/max capacity-generating set for Symp^{2n} has cardinality (strictly) bigger than the continuum.*

This result follows immediately from Corollary 18(i) on p. 855, which follows from Theorem 17. This corollary greatly generalizes Theorem 5 by extending it to (differential) form categories containing a certain one-parameter family of objects and by replacing the notion of limit-min/max generation by the much weaker notion of countable Borel-generation. (See Definition 16 on p. 853 and Remarks 15 and 19.)¹⁵ The idea of proof of part of Theorem 17 is to use Stokes' theorem for helicity. (We will explain this in Subsection 2.4.)

Theorem 5 and - more generally - Corollary 18 diminish the hope of finding manageable generating sets of (generalized) symplectic capacities.

Our second main result states that every suitably generating set for the generalized capacities on a small weak form category is uncountable, if the category contains a one-parameter family of objects that is "strictly volume-increasing" and "embedding-capacity-wise constant". Morally speaking, this hypothesis is weaker than those of the first main result. As a special case the second main result holds for the category of ellipsoids in dimension at least 4. This answers a variant of a question of CHLS about the Ekeland-Hofer capacities.

To explain this, let (V, ω) be a (finite-dimensional) symplectic vector space. We abbreviate $V := (V, \omega)$ and define $\widetilde{\text{Op}}_V$ to be the category with objects given by all pairs $(U, \omega|_U)$, where U ranges over all open subsets of V , and morphisms between two objects U, U' given by the restrictions $\varphi|_U$, where φ ranges over all symplectomorphisms of V , such that $\varphi(U) \subseteq U'$. This is a small¹⁶ weak symplectic category. For such a category we define the notion of a generalized capacity as in Definition 1.

Let $i \in \mathbb{N}_0$. The i -th Ekeland-Hofer capacity c_i^{EH} is a capacity on $\widetilde{\text{Op}}_V$, which is defined as a certain min-max involving the symplectic action, see

¹⁵We also formulate a version of the result for normalized capacities. See Corollary 18(ii).

¹⁶A category is called small iff its objects and its morphisms form sets.

[5, 6] or [3, p. 7]. The capacity c_1^{EH} is normalized; the other Ekeland-Hofer capacities are not normalized.

The Ekeland-Hofer capacities are hard to compute. Their values are known for ellipsoids and polydisks, see [6, Proposition 4, p. 562] and [6, Proposition 5, p. 563].

Recall that a (bounded, open, full) ellipsoid in V is a set of the form $p^{-1}((-\infty, 0))$, where $p : V \rightarrow \mathbb{R}$ is a quadratic polynomial function whose second order part is positive definite. We equip each ellipsoid E with the restriction of ω to E . We define the *restriction category of ellipsoids* to be the full subcategory \widetilde{Ell}_V of \widetilde{Op}_V consisting of ellipsoids.¹⁷ The objects of \widetilde{Ell}_V are uniquely determined by the Ekeland-Hofer capacities, up to isomorphism, see [3, FACT 10, p. 27]. Therefore the following question seems natural:

Question 6 ([3], Problem 15, p. 29). *Do the Ekeland-Hofer capacities together with the volume capacity¹⁸ form a CHLS generating set for Ell_V ?¹⁹*

In the case $\dim V = 4$ this question was answered negatively by D. McDuff, see [10, Corollary 1.4]. A special case of our second main result, Theorem 7 below, answers Question 6 in the negative in dimension at least 4, if we replace “CHLS generating” by “finitely differentially generating”, as defined below. In fact, it states that every finitely differentially generating set on the category of ellipsoids is uncountable.

Let \mathcal{C} be a small weak symplectic category. We call a subset \mathcal{G} of $\text{Cap}(\mathcal{C})$ *finitely differentially (capacity-)generating for \mathcal{C}* iff for every $c \in \text{Cap}(\mathcal{C})$ there exist $\ell \in \mathbb{N}_0$, $c_1, \dots, c_\ell \in \mathcal{G}$, and a differentiable function $f : [0, \infty]^\ell \rightarrow [0, \infty]$, such that $c = f(c_1, \dots, c_\ell)$.²⁰ A special case of our second main result is the following.

¹⁷The word “restriction” refers to the fact that the morphisms of \widetilde{Ell}_V are restrictions of global symplectomorphisms. \widetilde{Ell}_V is a subcategory of the category Ell_V whose objects and morphisms are given by ellipsoids and *all* symplectic embeddings. We reserve the nicer notation Ell_V for the bigger category.

¹⁸One needs to include the volume capacity, since the Ekeland-Hofer capacities do not CHLS generate this capacity, see [3, Example 10, p. 28].

¹⁹In [3] it is only asked whether these generalized capacities generate all capacities, not all *generalized* capacities. However, from the discussion that precedes the question it is clear that the authors intended to include the word *generalized* here.

²⁰Here we view $[0, \infty]$ as a compact 1-dimensional manifold with boundary. Its Cartesian power is a manifold with boundary and corners. The function f is only assumed to be differentiable one time, with possibly discontinuous derivative.

Theorem 7 (uncountability of every generating set for ellipsoids). *Let V be a symplectic vector space of dimension at least 4. Then every finitely differentially capacity-generating set for \widetilde{Ell}_V is uncountable.*

Proof. This follows from Theorem 25 on p. 860 and Example 26 by considering the ellipsoids

$$M_a := \left\{ x = (x_1, \dots, x_n) \in \mathbb{R}^{2n} = (\mathbb{R}^2)^n \mid \sum_{i=1}^{n-1} \|x_i\|^2 + \frac{\|x_n\|^2}{a} < 1 \right\},$$

for $a \in A := [1, \infty)$. Here $\|\cdot\|$ denotes the Euclidean norm on \mathbb{R}^2 . Our hypothesis $n \geq 2$ guarantees that the inequality “ \leq ” in condition (10) on p. 860 holds. □

The idea of proof of Theorem 25 is to use Lebesgue’s Monotone Differentiation Theorem. (We explain this in Subsection 2.4.)

By Theorem 7, in particular, the Ekeland-Hofer capacities together with the volume capacity do not finitely generate the set of all generalized capacities on \widetilde{Ell}_V . This provides a negative answer to the variant of Question 6 involving the notion of finite differentiable generation. We will generalize Theorem 7 to small weak form categories containing a certain type of one-parameter family of objects. (See Theorem 25 on p. 860.)

Our first main result (Theorem 17 below) also has immediate applications to two questions that CHLS prominently posed as Problems 1 and 2 in their article [3]. To explain these problems, let $n \in \mathbb{N}$ and $\mathcal{C} = (\mathcal{O}, \mathcal{M})$ be a symplectic category in dimension $2n$.

Definition (representability). *Let c be a capacity on \mathcal{C} . We call c (symplectically) domain-/ target-representable iff there exists a symplectic manifold (M, ω) , for which $c = c_{M, \omega} / c = c^{M, \omega}$ (defined as in Example 2). We call it connectedly target-representable iff there exists a connected symplectic manifold (M, ω) , for which $c = c^{M, \omega}$.*

Remark. By definition, the topology of a manifold is second countable. Without this condition every capacity would be target-representable, if all objects of \mathcal{C} are connected, see [3, Example 2, p. 14].

Question 8 (target-representability, [3], p. 14, Problem 1). *Which (generalized) capacities on \mathcal{C} are connectedly target-representable?*²¹

Question 9 (domain-representability, [3], p. 14, Problem 2). *Which (generalized) capacities on \mathcal{C} are domain-representable?*²²

In particular, one may wonder about the following:

Question 10. *Is every generalized capacity connectedly target-representable?*

If the answer to this question is “yes”, then this simplifies the study of capacities, since we may then identify every capacity with some symplectic manifold that target-represents it.

Apart from some elementary examples, up to now, the answers to Questions 8,9, and 10 appear to be completely unknown. In order to answer Question 10 negatively, it seems that we need to understand all symplectic embeddings from objects of \mathcal{C} to all connected symplectic manifolds. At first glance this looks like a hopeless enterprise.

The following application of the first main result may therefore come as a surprise. Namely, the answer to Question 10 is “no”, if the symplectic category is of dimension at least 4 and contains a certain one-parameter family of objects. In fact, the answer remains “no”, even if we ask the question only for *normalized* capacities and drop the word “connectedly”. Perhaps all the more unexpectedly, *almost no* normalized capacity is target-representable. Here we say that *almost no* element of a given infinite set has a given property iff the subset of all elements with this property has smaller cardinality than the whole set. Similarly, almost no normalized capacity is domain-representable. The following application of the first main result concerns the special case of these statements for the whole symplectic category.

Theorem (representability). *For every $n \geq 2$ almost no normalized capacity on Symp^{2n} is domain- or target-representable.*

²¹In [3] this question is asked, based on the definition of the embedding capacities on p. 13 in that article. In that definition the embedding capacity is only defined for objects of \mathcal{C} . However, in [3, Example 2, p. 14] the authors use the definition with a symplectic manifold that is not an object of \mathcal{C} . This suggests that CHLS are interested in Question 8 with the modified definition given in Example 2. Compare to footnote 8.

²²A remark similar to footnote 21 applies.

This is an immediate consequence of Corollary 20 on p. 857.²³ It follows that there are as many normalized capacities on Symp^{2n} that are neither domain- nor target-representable, as there are normalized capacities overall (namely $\mathcal{P}(\mathbb{R})$ -many²⁴). This provides some answers to Questions 8,9, and 10.

1.2. Organization of this article

In Section 2 we state the main results in the general setting of (weak) differential form categories and deduce the applications about generating sets and representability. We also present the ideas of proof of the main results. Furthermore, we discuss the related result of D. McDuff about monotone capacity-generation for the category of ellipsoids in dimension 4, and a potential application of our proof technique to morphism detection.

In Section 3 we formulate Theorem 43, which generalizes part of the first main result (Theorem 17(i,ii)). It states that the cardinality of the set of (normalized) capacities equals that of $\mathcal{P}(\mathbb{R})$ for every form category containing each disjoint union $M_a \sqcup M_{-a}$ for a suitable one-parameter family of manifolds with forms $(M_a, \omega_a)_{a \in A_0}$. This family needs to satisfy the following crucial condition. We denote by I_a the set of connected components of the boundary of M_a , and $I := (I_a)_{a \in A_0}$. Then the collection of boundary helicities associated with $(M_a, \omega_a)_{a \in A_0}$ is an I -collection. We introduce the notions of helicity and of an I -collection in this section. We also state Proposition 44, which provides sufficient criteria for the helicity hypothesis of Theorem 43.

In Sections 4 and 5 we prove Theorem 43 and Proposition 44.

Section 6 contains the proof of the last part of the first main result (Theorem 17(iii)), which states that every set of cardinality at most that of \mathbb{R} countably Borel-generates a set of cardinality at most that of \mathbb{R} .

Section 7 is devoted to the proof of our second main result, Theorem 25, stating that every finitely differentially generating set of capacities is uncountable if the category contains a certain type of one-parameter family of objects.

In Section A we prove an auxiliary result, which states that the set of diffeomorphism classes of manifolds has cardinality that of \mathbb{R} . We also show

²³In that corollary we use a definition of representability that allows for the representing pair (M, ω) to be a general manifold with a two-form.

²⁴Here $\mathcal{P}(S)$ denotes the power set of a set S .

that the same holds for the set of all equivalence classes of pairs (M, ω) , where M is a manifold and ω a differential form on M .

Finally, in Section B we deduce Theorem 29 (monotone generation for ellipsoids) from McDuff's characterization of the existence of symplectic embeddings between ellipsoids.

1.3. Acknowledgments

The authors would like to thank the anonymous referee for suggesting to make the introduction more readable by mentioning only special cases of the main results. We are grateful to Felix Schlenk for sharing his expertise on symplectic capacities. We also thank Urs Frauenfelder for an interesting discussion, Timofey Rodionov and Jiří Spurný for their help with a question about Baire and Borel hierarchies, and Asaf Karagila for his help with a question about aleph and bet numbers.

2. Main results and applications

In this section we state the main results of this article in the general setting and deduce the applications about generating sets and representability, taking care of the set-theoretic issue that was mentioned in the previous section.

2.1. Cardinalities of the set of capacities and of the generated set

The first main result provides conditions under which every generating set of capacities on a differential form category is bigger than the continuum. To define the notion of such a category, let $m, k \in \mathbb{N}_0 := \{0, 1, \dots\}$. We define $\Omega^{m,k}$ to be the following category:

- Its objects are pairs (M, ω) , where M is a manifold of dimension m , and ω is a differential k -form on M .
- Its morphisms are embeddings that intertwine the differential forms.

Definition. A weak (m, k) -(differential) form category is a subcategory $\mathcal{C} = (\mathcal{O}, \mathcal{M})$ of $\Omega^{m,k}$, such that if $(M, \omega) \in \mathcal{O}$ and $a \in (0, \infty)$ then $(M, a\omega) \in \mathcal{O}$. We call such a \mathcal{C} a (m, k) -form category iff it is also isomorphism-closed.

A (weak) symplectic category (in dimension $2n$) is a (weak) $(2n, 2)$ -form category whose objects are symplectic manifolds.

Examples ((weak) (m, k) -form category).

- (i) Let \mathfrak{M} be a diffeomorphism class of smooth manifolds of dimension m . The full subcategory of $\Omega^{m,k}$ whose objects (M, ω) satisfy $M \in \mathfrak{M}$, is an (m, k) -form category.
- (ii) Let (M, ω) be an object of $\Omega^{m,k}$. Consider the category with objects given by all pairs $(U, \omega|_U)$, where U ranges over all open subsets of M , and morphisms between two objects U, U' given by the restrictions $\varphi|_U$, where φ ranges over all isomorphisms of (M, ω) , such that $\varphi(U) \subseteq U'$. This is a small weak (m, k) -form category, which is not isomorphism-closed, hence not an (m, k) -form category.

Remark 11 (isomorphism-closedness). Symplectic categories were first defined in [3, 2.1. Definition, p. 5]. In that definition isomorphism-closedness is not assumed. However, this condition is needed in order to avoid the following set-theoretic issue in the definition of the notion of a symplectic capacity on a given symplectic category \mathcal{C} .

This article is based on ZFC, the Zermelo-Fraenkel axiomatic system together with the axiom of choice. A category is a pair consisting of classes of objects and morphisms. Formally, in ZFC there is no notion of a “class” that is not a set. The system *can* handle a “class” that is determined by a wellformed formula, such as the “class” of all sets or the “class” of all symplectic manifolds, by rewriting every statement involving the “class” as a statement involving the formula.

However, it is not possible in ZFC to define the “class” of all maps between two classes, even if the target class is a set. In particular, it is a priori not possible to define the “class” of all symplectic capacities on a given symplectic category. Our assumption that \mathcal{C} is isomorphism-closed makes it possible to define this “class” even as a set, see below.

We now define the notion of a (generalized) capacity on a given form category. Let S be a set. By $|S|$ we denote the (*von Neumann*) *cardinality of S* , i.e., the smallest (von Neumann) ordinal that is in bijection with S . For every pair of sets S, S' we denote by S'^S the set of maps from S to S' . For every pair of cardinals α, β ²⁵ we also use β^α to denote the cardinality of β^α . Recursively, we define $\beth_0 := \aleph_0$, and for every $i \in \mathbb{N}_0$, the cardinal $\beth_{i+1} := 2^{\beth_i}$.²⁶

²⁵i.e., cardinalities of some sets

²⁶ \beth (bet) is the second letter of the Hebrew alphabet.

Let $\mathcal{C} = (\mathcal{O}, \mathcal{M})$ be an (m, k) -form category. We define the set

$$(2) \quad \mathcal{O}_0 := \{(M, \omega) \in \mathcal{O} \mid \text{The set underlying } M \text{ is a subset of } \mathbb{Q}_1.\}$$

Definition 12. A generalized capacity on \mathcal{C} is a function

$$c : \mathcal{O}_0 \rightarrow [0, \infty]$$

with the following properties:

(i) (**monotonicity**) If (M, ω) and (M', ω') are two objects in \mathcal{O}_0 between which there exists a \mathcal{C} -morphism, then

$$c(M, \omega) \leq c(M', \omega').$$

(ii) (**conformality**) For every $(M, \omega) \in \mathcal{O}_0$ and $a \in (0, \infty)$ we have

$$c(M, a\omega) = a c(M, \omega).$$

Assume now that $k = 2$, $m = 2n$ for some integer n , and that \mathcal{O}_0 contains some objects B_0, Z_0 that are isomorphic to B, Z (the open unit ball and cylinder). Let c be a generalized capacity on \mathcal{C} . We call c a capacity iff it satisfies:

(iii) (**non-triviality**) $c(B_0) > 0$ and $c(Z_0) < \infty$.²⁷

We call it normalized iff it satisfies:

(iv) (**normalization**) $c(B_0) = c(Z_0) = \pi$.

We denote by

$$\text{Cap}(\mathcal{C}), \quad \mathcal{N}\text{Cap}(\mathcal{C})$$

the sets of generalized and normalized capacities on \mathcal{C} . If \mathcal{C} is a symplectic category then we call a (generalized/ normalized) capacity on \mathcal{C} also a (generalized/ normalized) symplectic capacity.

Example 13 (embedding capacities). Let $\mathcal{C} = (\mathcal{O}, \mathcal{M})$ be an (m, k) -form category and (M, ω) an object of $\Omega^{m, k}$. We define \mathcal{O}_0 as in (2) and the

²⁷These conditions do not depend on the choices of B_0, Z_0 , since c is invariant under isomorphisms by monotonicity.

domain-embedding capacity for (M, ω) on \mathcal{C} to be the function

$$(3) \quad c_{M,\omega} := c_{M,\omega}^{\mathcal{C}} : \mathcal{O}_0 \rightarrow [0, \infty],$$

$$c_{M,\omega}(M', \omega') := \sup \{c \in (0, \infty) \mid \exists \Omega^{m,k}\text{-morphism } (M, c\omega) \rightarrow (M', \omega')\}.$$

We define the target-embedding capacity for (M, ω) on \mathcal{C} to be the function

$$c^{M,\omega} := c_{\mathcal{C}}^{M,\omega} : \mathcal{O}_0 \rightarrow [0, \infty],$$

$$c^{M,\omega}(M', \omega') := \inf \{c \in (0, \infty) \mid \exists \Omega^{m,k}\text{-morphism } (M', \omega') \rightarrow (M, c\omega)\}.$$

These are generalized capacities.

Remarks 14 (set of capacities and isomorphism-closedness). (i)

The collections $\mathcal{C}ap(\mathcal{C})$ and $\mathcal{N}C\mathcal{C}ap(\mathcal{C})$ are indeed sets, since \mathcal{O}_0 is a set.

(ii) Heuristically, let us denote by $\widetilde{\mathcal{C}ap}(\mathcal{C})$ the “subclass” of “[0, ∞]^ℳ” consisting of all “functions” satisfying conditions the monotonicity and conformality conditions of Definition 12. Formally, the restriction from \mathcal{O} to \mathcal{O}_0 induces a bijection between $\widetilde{\mathcal{C}ap}(\mathcal{C})$ and $\mathcal{C}ap(\mathcal{C})$.²⁸ This means that our definition of a generalized capacity corresponds to the intuition behind the usual “definition”. Here we use isomorphism-closedness of \mathcal{C} . Compare to Remark 11.

(iii) Isomorphism-closedness of \mathcal{C} implies that there is a canonical bijection between $\mathcal{C}ap(\mathcal{C})$ and the set of generalized capacities that we obtain by replacing \mathcal{O}_0 by any subset of \mathcal{O} that contains an isomorphic copy of each element of \mathcal{O} . Such a subset can for example be obtained by replacing \beth_1 in (2) by any set of cardinality at least \beth_1 .²⁹ This means that our definition of $\mathcal{C}ap(\mathcal{C})$ is natural.

(iv) Example 13 generalizes Example 2, taking care of the set-theoretic issue mentioned in Remark 11.

²⁸This follows from the fact that every object of $\Omega^{m,k}$ is isomorphic to one whose underlying set is a subset of \beth_1 , and the assumption that \mathcal{C} is isomorphism-closed. To prove the fact, recall that by definition, the topology of every manifold M is second countable. Using the axiom of choice, it follows that its underlying set has cardinality $\leq \beth_1$. This means that there exists an injective map $f : M \rightarrow \beth_1$. Consider now an object (M, ω) of $\Omega^{m,k}$. Pushing forward the manifold structure and ω via a map f , we obtain an object of $\Omega^{m,k}$ isomorphic to (M, ω) , whose underlying set is a subset of \beth_1 . This proves the fact.

²⁹This follows from an argument as in the last footnote.

Let \mathcal{G} be a subset of $\mathcal{C}ap(\mathcal{C})$. We define the set CHLS generated by \mathcal{G} as on p. 842, except that we ask the domain of each function c to be \mathcal{O}_0 rather than \mathcal{O} . The reason for this change is the set-theoretic issue discussed in Remark 11. (See also Remarks 14.) As mentioned in Section 1.1, in [3, Problem 5, p. 17] K. Cieliebak, H. Hofer, J. Latschev, and F. Schlenk posed the following problem:

Problem. *For a given symplectic category find a minimal CHLS capacity-generating set.*

In particular, we may ask whether there exists a *countable* generating set. Our first main result, Theorem 17 below, answers this question in the negative for a notion of a generation that, morally speaking, is much weaker than CHLS generation. The theorem states that in dimension at least 4 the cardinality of the set of generalized (or normalized) capacities on \mathcal{C} is \beth_2 , provided that the category contains a certain one-parameter family of objects.³⁰ Its last part implies that a set of $[0, \infty]$ -valued functions of cardinality at most \beth_1 countably Borel-generates a set of cardinality at most \beth_1 in the sense of Definition 16 below.

As an immediate consequence, every countably Borel-generating set for $\mathcal{C}ap(\mathcal{C})$ (or $\mathcal{N}C\mathcal{C}ap(\mathcal{C})$) has cardinality bigger than the continuum. See Corollary 18 below. Countable Borel-generation is a weak notion of generation. (Compare to Remark 15 below.) It is weaker than the notion of limit-min/max generation (as defined on p. 842).³¹ Hence Corollary 18 makes a statement about a large class of “generating sets of capacities”.

This corollary diminishes the hope of finding manageable generating sets of (generalized) symplectic capacities.

To state our first main result, we need the following. Let (X, τ) be a topological space. Recall that the (τ) -Borel σ -algebra of X is the smallest σ -algebra containing the topology of X . We call its elements (τ) -Borel sets.

Remark 15 (Borel sets). Consider the real line $X = \mathbb{R}$. The axiom of choice (AC) implies that there exist subsets of \mathbb{R} that are not Lebesgue-measurable, hence not Borel-measurable. However, all subsets occurring in practice are Borel. Furthermore, for any concretely described subset of \mathbb{R} ,

³⁰It is formulated for a form category, not just for a symplectic category.

³¹This follows from the fact that a given set of capacities countably Borel-generates a larger set than it limit-min/max generates. Compare to Remark 19.

it appears to be difficult to prove (using AC) that it is indeed not Borel-measurable.³²

Let now (X, τ) and (X', τ') be topological spaces. A map $f : X \rightarrow X'$ is called (τ, τ') -Borel-measurable iff the pre-image under f of every τ' -Borel set in X' is a τ -Borel set.³³ In particular, every continuous map is Borel-measurable. Borel-measurability is preserved under composition. It is preserved under pointwise limits of sequences if X' is metrizable. This yields many examples of Borel-measurable maps. In fact, all maps occurring in practice are Borel-measurable.

Let S, S' be sets. We denote

$$S'^S := \{\text{map from } S \text{ to } S'\}.$$

For every subset $\mathcal{G} \subseteq S'^S$ we denote by

$$(4) \quad \text{ev}_{\mathcal{G}} : S \rightarrow S'^{\mathcal{G}}, \quad \text{ev}_{\mathcal{G}}(s)(u) := u(s),$$

the evaluation map. If (X, τ) is a topological space then we denote by τ_S the product topology on X^S .

Definition 16 (countably Borel-generated set). *Let S be a set, (X, τ) a topological space, and $\mathcal{G} \subseteq X^S$. We define the set countably (τ) -Borel-generated by \mathcal{G} to be*

$$\langle \mathcal{G} \rangle := \{f \circ \text{ev}_{\mathcal{G}_0} \mid \mathcal{G}_0 \subseteq \mathcal{G} \text{ countable, } f : X^{\mathcal{G}_0} \rightarrow X : (\tau_{\mathcal{G}_0}, \tau)\text{-Borel-measurable}\} \subseteq X^S.$$

For every subset $\mathcal{F} \subseteq X^S$ we say that \mathcal{G} countably (τ) -Borel-generates at least \mathcal{F} iff $\mathcal{F} \subseteq \langle \mathcal{G} \rangle$.

We denote by $\text{int } S$ the interior of a subset S of a topological space. Let V be a vector space, $S \subseteq V$, $A \subseteq \mathbb{R}$, and $n \in \mathbb{N}_0$. We denote $AS := \{av \mid a \in A, v \in S\}$. In the case $A = \{a\}$ we also denote this set by aS . We call S

³²An example of such a subset A was provided by N. Luzin. It can be obtained from [9, Exercise (27.2), p. 209] via [9, Exercise (3.4)(ii), p. 14]. This set is Σ^1_1 -analytic, see [9, Definitions (22.9), p. 169, (21.13), p. 156]. It follows from a theorem of Souslin, [9, (14.2) Theorem, p. 85] and the definition of Σ^1_1 -analyticity that A is not Borel.

³³This happens if and only if the pre-image under f of every element of τ' is a τ -Borel set.

strictly starshaped around 0 iff $[0, 1)S \subseteq \text{int } S$. For every $i \in \{1, \dots, n\}$ we denote by $\text{pr}_i : V^n = V \times \dots \times V \rightarrow V$ the canonical projection onto the i -th component. For every multilinear form ω on V we denote

$$\omega^{\oplus n} := \sum_{i=1}^n \text{pr}_i^* \omega.$$

For every $r \in (1, \infty)$ we define the closed spherical shell of radii 1, r in \mathbb{R}^m to be

$$Sh_r^m := \overline{B}_r^m \setminus B_1^m.$$

We equip $Sh_r := Sh_r^{2n}$ with the standard symplectic form ω_{st} . The first main result of this article is the following.

Theorem 17 (cardinalities of the set of (normalized) capacities and of the generated set). *The following statements hold:*

(i) *Let $k, n \in \{2, 3, \dots\}$ with k even, and $\mathcal{C} = (\mathcal{O}, \mathcal{M})$ be a (kn, k) -form category. Then the cardinality of $\text{Cap}(\mathcal{C})$ equals \beth_2 , provided that there exist*

- *a (real) vector space V of dimension k ,*
- *a volume form Ω on V ,³⁴*
- *a nonempty compact submanifold K of V^n (with boundary) that is strictly starshaped around 0,*
- *a number $r \in (1, \sqrt[kn]{2})$,*

such that defining $M_a := (r + a)K \setminus \text{int } K$ and equipping this manifold with the restriction of $\Omega^{\oplus n}$, we have

$$(5) \quad M_a \sqcup M_{-a} \in \mathcal{O}, \quad \forall a \in (0, r - 1).^{35}$$

³⁴By this we mean a nonvanishing top degree skewsymmetric multilinear form.

³⁵Here $A \sqcup B$ denotes the disjoint union of two sets A, B . This can be defined in different ways, e.g. as the set consisting of all pairs $(0, a), (1, b)$, with $a \in A, b \in B$, or alternatively pairs $(1, a), (2, b)$. Based on this, we obtain two definitions of the disjoint union of two objects of $\Omega^{kn, k}$. The disjoint union defined in either way is isomorphic to the one defined in the other way. Since we assume \mathcal{C} to be isomorphism-closed, condition (5) does not depend on the choice of how we define the disjoint union.

(ii) Let $n \in \{2, 3, \dots\}$ and $\mathcal{C} = (\mathcal{O}, \mathcal{M})$ be a $(2n, 2)$ -form category that contains the objects B and Z . The cardinality of $\mathcal{N}Cap(\mathcal{C})$ equals \beth_2 , provided that there exists $r \in (1, \sqrt[2^n]{2})$ satisfying

$$(6) \quad Sh_{r-a} \sqcup Sh_{r+a} \in \mathcal{O}, \quad \forall a \in (0, r - 1).$$

(iii) Let S be a set and (X, τ) a separable metrizable topological space. If a subset of X^S has cardinality at most \beth_1 , then the set it countably τ -Borel-generates has cardinality at most \beth_1 .

This result has the following immediate application. We define \mathcal{O}_0 as in (2), and τ_0 to be the standard topology on $[0, \infty]$, w.r.t. which it is homeomorphic to the interval $[0, 1]$.

Corollary 18 (cardinality of a generating set).

- (i) Under the hypotheses of Theorem 17(i) every subset of $[0, \infty]^{\mathcal{O}_0}$ that countably τ_0 -Borel-generates at least $Cap(\mathcal{C})$ has cardinality bigger than \beth_1 .
- (ii) Under the hypotheses of Theorem 17(ii) every subset of $[0, \infty]^{\mathcal{O}_0}$ that countably τ_0 -Borel-generates at least $\mathcal{N}Cap(\mathcal{C})$ has cardinality bigger than \beth_1 .

This corollary answers Question 4 (p. 842) negatively for every symplectic category satisfying the hypotheses of Theorem 17(i). An example of such a category is the category of all symplectic manifolds of some fixed dimension, which is at least 4.

Remarks (cardinalities of the set of (normalized) capacities and of the generated set).

- As Corollary 18 holds for (kn, k) -form categories with k even and $n \geq 2$, the fact that generating sets of capacities are large, is not a genuinely symplectic phenomenon.
- The proof of Theorem 17(ii) shows that the cardinality of the set of *discontinuous* normalized capacities is \beth_2 . This improves the result of K. Zehmisch and the second author that discontinuous capacities exist, see [17].³⁶

³⁶The proof of [17, Theorem 1.2] actually shows that the spherical shell capacities used in that proof are all different. This implies that the set of discontinuous normalized symplectic capacities has cardinality at least \beth_1 .

- The statements of Theorem 17(i,ii) and thus of Corollary 18 hold in a more general setting, see Theorem 43 and Proposition 44 below. In particular, let V, Ω be as in Theorem 17(i), $j \in \{1, 2, \dots\}$, and for each $a \in \mathbb{R}$ let M_a be the complement of j disjoint connected open sets in some compact submanifold of V^n . The cardinality of $\mathcal{C}ap(\mathcal{C})$ equals \beth_2 , provided that $M_a \sqcup M_{-a} \in \mathcal{O}$ ³⁷ for every a , the volumes of the open sets are all equal (also for different a), the volume of each M_a is small enough and strictly increasing in a , the infimum of these volumes is positive, and each M_a is 1-connected.
- Morally, Corollary 18 implies that every generating set of capacities has as many elements as there are capacities. More precisely, we denote by ZF the Zermelo-Fraenkel axiomatic system, and $ZFC := ZF + AC$. We claim that ZFC is consistent with the statement that under the hypotheses of Theorem 17(i) every subset of $[0, \infty]^{\mathcal{O}_0}$ that countably Borel-generates at least $\mathcal{C}ap(\mathcal{C})$ has the same cardinality as $\mathcal{C}ap(\mathcal{C})$ (namely \beth_2) ³⁸.

To see this, assume that the generalized continuum hypothesis (GCH) holds. This means that for every infinite cardinal α there is no cardinal strictly between α and 2^α . In particular, there is no cardinal strictly between \beth_1 and $\beth_2 = 2^{\beth_1}$. Hence under the hypotheses of Theorem 17(i) by Corollary 18(i) every subset of $[0, \infty]^{\mathcal{O}_0}$ that countably Borel-generates at least $\mathcal{N}Cap(\mathcal{C})$ has cardinality at least \beth_2 . By Theorem 17(i) this equals the cardinality of $\mathcal{C}ap(\mathcal{C})$. Since GCH is consistent with ZFC ³⁹, the claim follows.

Remark 19 (comparison of different notions of generating sets).

Let \mathcal{C} be a symplectic category and \mathcal{G} a CHLS generating set (as defined on p. 842), with the extra condition that each combining function $f : [0, \infty]^\ell \rightarrow [0, \infty]$ is Borel-measurable. Then \mathcal{G} countably Borel-generates $\mathcal{C}ap(\mathcal{C})$. (See

³⁷In particular we assume here that M_a is a smooth submanifold of V^n .

³⁸provided that ZF is consistent

³⁹provided that ZF is consistent

Definition 16.)⁴⁰ This holds in particular if \mathcal{G} is limit-min/max generating (as defined on p. 842).

Definition 16 relaxes the conditions in the definition of a CHLS generating set in two ways:

- The combining functions are allowed to depend on countably many variables (elements of the generating set), not just on finitely many variables.
- The assumption that the combining functions are homogeneous and monotone is omitted.

2.2. Representability of symplectic capacities and morphism detection

The following corollary is a direct consequence of Theorem 17(ii). Let \mathcal{C} be an (m, k) -form category. We say that a generalized capacity c on \mathcal{C} is *domain-/ target-representable* iff there exists an object (M, ω) of $\Omega^{m,k}$, such that $c = c_{(M,\omega)}$ / $c = c^{(M,\omega)}$ (defined as in Example 13). We say that *almost no* element of a given infinite set has a given property iff the subset of all elements with this property has smaller cardinality than the whole set.

Corollary 20 (representability). *Under the hypotheses of Theorem 17(ii) almost no normalized capacity on \mathcal{C} is domain- or target-representable.*

It follows that under the hypotheses of Theorem 17(ii) there are as many normalized capacities that are neither domain- nor target-representable, as there are normalized capacities overall (namely \beth_2). This provides some answers to Questions 8,9, and 10 (p. 845). The statement of Corollary 20 holds in particular for \mathcal{C} given by the category of all symplectic manifolds of some fixed dimension, which is at least 4.

⁴⁰To see this, let $c \in \text{Cap}(\mathcal{C})$. We choose a sequence of Borel-measurable combining functions and finite sets of generalized capacities in \mathcal{G} as in the definition of a CHLS generating set. We define \mathcal{G}_0 to be the set of all capacities occurring in the sequence. Each combining function gives rise to a Borel-measurable function from $[0, \infty]^{\mathcal{G}_0}$ to $[0, \infty]$. Its restriction to the image of $\text{ev}_{\mathcal{G}_0}$ is measurable w.r.t. the σ -algebra induced by the Borel σ -algebra. By assumption the sequence of these restrictions converges pointwise. The limit f is again measurable. Since its target space is $[0, \infty]$, an argument involving approximations by simple functions shows that f extends to a Borel-measurable function on $[0, \infty]^{\mathcal{G}_0}$. Hence \mathcal{G}_0 and f satisfy the conditions of Definition 16, as desired.

Proof of Corollary 20. The set of isomorphism classes of $2n$ -dimensional manifolds together with 2-forms has cardinality \beth_1 . This follows from Corollary 58 on p. 904. The image of this set under the map $[(M, \omega)] \mapsto c_{M, \omega}$ is the set of all domain-representable capacities. It follows that at most \beth_1 normalized capacities are domain-representable. A similar statement holds for target-representation. The statement of Corollary 20 now follows from Theorem 17(ii). \square

The proof technique for Corollary 18 can potentially also be used to show that certain sets of capacities do not detect morphisms. To explain this, let $\mathcal{C} = (\mathcal{O}, \mathcal{M})$ be an (m, k) -form category and $\mathcal{G} \subseteq \text{Cap}(\mathcal{C})$. We say that \mathcal{G} detects morphisms iff for each pair of objects $(M, \omega), (M', \omega')$ of \mathcal{C} the following holds. Assume that $c(M, \omega) \leq c(M', \omega')$, for every $c \in \mathcal{G}$. Then there exists a \mathcal{C} -morphism from (M, ω) to (M', ω') . CHLS asked the following question in the case in which \mathcal{C} is a symplectic category and $\mathcal{G} = \text{Cap}(\mathcal{C})$ (see [3, Question 1, p. 20]):

Question 21. *Does \mathcal{G} detect morphisms?*⁴¹

Remark (monotone generation and detection of morphisms). We equip the set $S := \mathcal{O}_0$ with the pre-order given by $(M, \omega) \leq (M', \omega')$ iff there exists a \mathcal{C} -morphism from (M, ω) to (M', ω') . We also equip this set with the $(0, \infty)$ -action given by rescaling of forms. Suppose the following:

- (*) Every subset of $\text{Cap}(\mathcal{C})$ that monotonely generates in the sense of Definition 28 on p. 864, has cardinality bigger than \beth_1 .

Let $\mathcal{G} \subseteq \text{Cap}(\mathcal{C})$ be a subset of cardinality at most \beth_1 . Then \mathcal{G} does not detect morphisms, thus the answer to Question 21 is “no”. To see this, observe that by our assumption (*) the set \mathcal{G} does not monotonely generate. Therefore, by Proposition 60 on p. 906, it is not almost order-reflecting. Hence \mathcal{G} is not order-reflecting, i.e., it does not detect morphisms.

By Corollary 18, under the hypotheses of Theorem 17(i), condition (*) is satisfied with “monotonely generates” replaced by “countably τ_0 -Borel-generates”. Therefore potentially, the proof technique for Corollary 18 may be adapted, in order to provide a negative answer to Question 21 under

⁴¹CHLS do not use our terminology of “morphism detection”. Instead, the title of the subsection in which they ask their Question 1, is “Recognition”. We think that the expression “ \mathcal{G} detects morphisms” more accurately describes the condition that \mathcal{G} determines whether there exists a morphism between two given objects.

suitable conditions on \mathcal{C} that do not involve (*), if the cardinality of \mathcal{G} is at most \beth_1 .

2.3. Uncountability of every generating set under a mild hypothesis

Our second main result states that every finitely differentially capacity-generating set for a small weak form category is uncountable, if the category contains a one-parameter family of objects that is “strictly volume-increasing” and “embedding-capacity-wise constant”. Morally speaking, this hypothesis is weaker than those of the first main result. In order to state the result, we recall the notion of finite differentiable generation from Section 1.1, reformulating and generalizing it slightly:

Definition (finite differentiable generation). *Let S be a set, and $\mathcal{F}, \mathcal{G} \subseteq [0, \infty]^S$. We say that \mathcal{G} finitely differentially generates at least \mathcal{F} iff the following holds. For every $F \in \mathcal{F}$ there exists a finite subset $\mathcal{G}_0 \subseteq \mathcal{G}$ and a differentiable function $f : [0, \infty]^{\mathcal{G}_0} \rightarrow [0, \infty]$ ⁴², such that $F = f \circ \text{ev}_{\mathcal{G}_0}$.*

Let now $k, n \in \mathbb{N} := \{1, 2, \dots\}$ and (M, ω) be an object of $\Omega^{kn, k}$. We call ω *maxipotent* iff $\omega^{\wedge n} = \omega \wedge \dots \wedge \omega$ does not vanish anywhere.

Remark 22 (maxipotency and nondegeneracy). Let V be a (real) vector space and $k \in \mathbb{N}$. We call a k -linear form ω on V *nondegenerate* iff interior multiplication with ω is an injective map from V to the space of $(k - 1)$ -linear forms. Let $k, n \in \mathbb{N}$ and assume that $\dim V = kn$. We call a skewsymmetric k -form ω on V *maxipotent* iff $\omega^{\wedge n} \neq 0$. Every maxipotent form on V is nondegenerate. The converse holds if and only if $k = 1$, $k = 2$, or $n = 1$.

Let (M, ω) be a maxipotent object of $\Omega^{kn, k}$. We equip M with the orientation induced by $\omega^{\wedge n}$ and define

$$(7) \quad \text{Vol}(M) := \text{Vol}(M, \omega) := \frac{1}{n!} \int_M \omega^{\wedge n}.$$

⁴²Here we view $[0, \infty]$ as a compact 1-dimensional manifold with boundary. The set $[0, \infty]^{\mathcal{G}_0}$ carries a canonical structure of a smooth finite-dimensional manifold with boundary and corners. The function f is only assumed to be differentiable one time, with possibly discontinuous derivative.

Remark 23 (volume). Assume that k is odd. Then we have $\omega \wedge \omega = 0$, and therefore $\text{Vol}(M, \omega) = 0$ in the case $n \geq 2$.

Let $\mathcal{C} = (\mathcal{O}, \mathcal{M})$ be a small weak (kn, k) -form category. We define the notion of a generalized capacity as in Definition 12.

Remark 24. Since \mathcal{C} is small, Definition 12 and our original Definition 1 are equivalent in the sense that capacities in either sense correspond to each other in a canonical way. This follows from Remark 14(iii). Recall here that a capacity in the sense of Definition 12 (Definition 1) is a function with domain $\mathcal{O}_0(\mathcal{O})$.

Our second main result is the following.

Theorem 25 (uncountability of every generating set under a mild hypothesis). *Every subset of $\text{Cap}(\mathcal{C})$ that finitely differentially generates (at least) $\text{Cap}(\mathcal{C})$, is uncountable, provided that there exists an interval A of positive length and a map $M : A \rightarrow \mathcal{O}$, such that*

$$(8) \quad M_a := M(a) \text{ is maxipotent for every } a \in A,$$

$$(9) \quad \text{Vol} \circ M \text{ is continuous and strictly increasing,}$$

$$(10) \quad c_{M_a}(M_{a'}) = 1, \forall a, a' \in A : a \leq a'.$$

Remarks.

- Condition (8) ensures that the volume of each M_a is well-defined. Hence condition (9) makes sense.
- Condition (10) means that M is “embedding-capacity-wise constant”, in the sense that the composition of the map $\{(a, a') \in A^2 \mid a \leq a'\} \ni (a, a') \mapsto (M_a, M_{a'})$ with the “embedding capacity function” $(X, X') \mapsto c_X(X')$ is constant.
- Assume that there exists a map M satisfying (8,9). Then we have $n > 0$. If $n \geq 2$, then k is even. This follows from Remark 23. Assume that there exists a map satisfying (8,9,10). Then we have $k > 0$. If each M_a is compact, then $n \neq 1$. This follows from Moser’s isotopy argument.

Example 26. Let $n \geq 2$ and A be an interval of positive length. We denote by \mathcal{U} the set of all open subsets of \mathbb{R}^{2n} that contain B_1^{2n} and are contained in Z_1^{2n} . We equip each element of \mathcal{U} with the restriction of the form ω_{st} . Let $M :$

$\widetilde{A} \rightarrow \mathcal{U}$ be an increasing map in the sense that $a \leq a'$ implies that $M(a) \subseteq M(a')$. If M also satisfies (9) then it satisfies all conditions of Theorem 25. The inequality “ \leq ” in condition (10) follows from Gromov’s nonsqueezing theorem.

As an application of Theorem 25 we obtain Theorem 7 (p. 845, uncountability of every generating set for ellipsoids).

Remark. Theorem 7 is concerned with the *weak* symplectic category \widetilde{Ell}_V of ellipsoids in V . This is not a symplectic category, since it is not isomorphism-closed in $\Omega^{2n,2}$, where $2n := \dim V$. This is the reason for formulating Theorem 25 for a (small) *weak* form category.

- Remarks.**
- The hypotheses of Corollary 18(i) (i.e., of Theorem 17(i)) and of Theorem 25 do not imply each other. However, morally, the hypotheses of Corollary 18(i) are more restrictive than that of Theorem 25. This becomes literally true if we modify the hypotheses of this corollary by replacing the disjoint union $M_a \sqcup M_{-a}$ by M_a .
 - On the other hand, the conclusion of Corollary 18(i) is stronger than that of Theorem 25.

2.4. Ideas of proof

The idea of the proof of Theorem 17(i) is the following. Recall the definition (3) of the domain-embedding capacity $c_M := c_{M,\omega}$. We choose V, Ω, K, r and define M_a as in the hypothesis of the theorem. We define $W_a := M_a \sqcup M_{-a}$. For each $A \in \mathcal{P}((0, r - 1))$ ⁴³ we define

$$c_A := \sup_{a \in A} c_{W_a}.$$

This is a capacity, satisfying

$$(11) \quad c_A(W_a) = 1, \quad \forall a \in A,$$

$$(12) \quad \sup_{a \in (0, r - 1) \setminus A} c_A(W_a) < 1.$$

The second statement follows from Stokes’ Theorem for helicity. Helicity assigns a real number to an exact k -form on a closed oriented manifold of dimension $kn - 1$, where $n \geq 2$. (To build some intuition, see the explanations

⁴³Here $\mathcal{P}(S)$ denotes the power set of a set S .

on p. 871 and Figures 1,2,3 on p. 872–873.) Helicity generalizes contact volume. The conditions (11,12) imply that $c_A \neq c_{A'}$ if $A \neq A' \in \mathcal{P}((0, r - 1))$. Since the cardinality of $\mathcal{P}((0, r - 1))$ equals \beth_2 , it follows that the cardinality of $\mathcal{C}ap(\mathcal{C})$ is at least \beth_2 .

On the other hand, we denote by S the set of isomorphism classes of objects of \mathcal{C} . This set has cardinality \beth_1 . Since $\mathcal{C}ap(\mathcal{C})$ can be viewed as a subset of $[0, \infty]^S$, it has cardinality at most \beth_2 , hence equal to \beth_2 .

A refined version of this argument shows Theorem 17(ii), i.e., that $|\mathcal{N}Cap(\mathcal{C})| = \beth_2$. For this we normalize each capacity c_A , by replacing it by the maximum of c_A and the Gromov width.

Remark (helicity argument). In [17] K. Zehmisch and F. Ziltener used helicity to show that the spherical capacity is discontinuous on some smooth family of ellipsoidal shells. This argument is related to the proof of Theorem 17(i,ii).

The proof of Theorem 17(iii) is based on the fact that the set of Borel-measurable maps from a second countable space to a separable metrizable space has cardinality at most \beth_1 . The proof of this uses the following well-known results:

- Every map f with target a separable metric space is determined by the pre-images under f of balls with rational radii around points in a countable dense subset.
- The σ -algebra generated by a collection of cardinality at most \beth_1 has itself cardinality at most \beth_1 . The proof of this uses transfinite induction.

The idea of the proof of Theorem 25 is to use Lebesgue's Monotone Differentiation Theorem, which states that every monotone function on an interval is differentiable almost everywhere. It follows that for every countable set \mathcal{G} of capacities, there exists a point $a_0 \in A$ at which the function $c \circ M$ is differentiable, for every $c \in \mathcal{G}$. On the other hand, our conditions on the map $M : A \rightarrow \mathcal{O}$ imply that the function $c_{M_{a_0}} \circ M$ is not differentiable at a_0 . It follows that \mathcal{G} does not finitely differentiably generate $c_{M_{a_0}}$.

Remark (diagonal argument). This idea of the proof is remotely reminiscent of Cantor's second diagonal argument, which shows that the set of real numbers is uncountable. Namely, consider the open sentence P given by:

$P(a, b)$: “The function $c_{M_a} \circ M$ is differentiable at b .”

The proof of Theorem 25 exploits the fact that P is false along the diagonal, that is, $P(a, a)$ is false for all a .

2.5. McDuff’s characterization of existence of symplectic embeddings for ellipsoids in dimension 4 and monotone generation

In this subsection we recall a result of D. McDuff, which states that the ECH-capacities are monotonely generating for the category of ellipsoids in dimension 4. On ellipsoids, these capacities are given by the following. Let $n, j \in \mathbb{N}_0$. We define the function

$$\mathcal{N}_j^n : [0, \infty)^n \rightarrow [0, \infty),$$

$$\mathcal{N}_j^n(a) := \min \left\{ b \in [0, \infty) \mid j+1 \leq \# \left\{ m \in \mathbb{N}_0^n \mid m \cdot a = \sum_{i=1}^n m_i a_i \leq b \right\} \right\}.$$

Remark. The sequence $(\mathcal{N}_j^n(a))_{j \in \mathbb{N}_0}$ is obtained by arranging all the non-negative integer combinations of a_1, \dots, a_n in increasing order, with repetitions.

We define the ellipsoid

$$E(a) := \left\{ x = (x_1, \dots, x_n) \in \mathbb{R}^{2n} = (\mathbb{R}^2)^n \mid \sum_{i=1}^n \frac{\|x_i\|^2}{a_i} < 1 \right\}.$$

(Here $\|\cdot\|$ denotes the Euclidean norm on \mathbb{R}^2 .) We equip this manifold with the standard symplectic form.

Let $V := (V, \omega)$ be a symplectic vector space. We denote by $\mathcal{O}_V^{\text{Ell}}$ the set of all pairs $(E, \omega|_E)$, where E is a (bounded, open, full) ellipsoid in V , and by $\mathcal{M}_V^{\text{Ell}}$ the set of all symplectic embeddings between elements of $\mathcal{O}_V^{\text{Ell}}$. We define the *category of (open) ellipsoids in V* to be the pair $\text{Ell}_V := (\mathcal{O}_V^{\text{Ell}}, \mathcal{M}_V^{\text{Ell}})$. This is a *small* weak symplectic category. For such a category we may view a generalized capacity as a monotone and conformal function on the *whole* set of objects. (Compare to Remark 24.) For every $j \in \mathbb{N}_0$ we define the function

(13)
$$c_j^V : \mathcal{O}_V^{\text{Ell}} \rightarrow [0, \infty),$$

by setting $c_j^V(E) := \mathcal{N}_j^n(a)$, where $a \in [0, \infty)^n$ is such that E is affinely symplectomorphic to $E(a)$. This number is well-defined, i.e., such an a exists (see [12, Lemma 2.43]) and $\mathcal{N}_j^n(a)$ does not depend on its choice. The latter is true, since if $E(a)$ and $E(a')$ are affinely symplectomorphic, then a and a' are permutations of each other. (See [12, Lemma 2.43].) The following result is due to M. Hutchings.

Theorem 27 (ECH-capacities). *If $\dim V = 4$ then for every $j \in \mathbb{N}_0$ the function c_j^V is a generalized capacity on Ell_V .*

Proof. Conformality follows from the definition of \mathcal{N}_j^n . Monotonicity was proved by M. Hutchings in [8, Proposition 1.2, Theorem 1.1]. □

Remarks.

- The function c_j^V is the restriction of the j -th ECH-capacity to Ell_V , see [8, Proposition 1.2].
- The category \widetilde{Ell}_V that we considered on p. 844 is a subcategory of Ell_V .

McDuff proved that the set of all c_j^V (with $j \in \mathbb{N}_0$) monotonely generates all generalized capacities. To explain this, we equip the interval $(0, \infty)$ with multiplication and let it act on the extended interval $[0, \infty]$ via multiplication. Let S, S' be sets. We fix $(0, \infty)$ -actions on S and S' and call a map $f : S \rightarrow S'$ (*positively 1-homogeneous*) iff it is $(0, \infty)$ -equivariant.

Recall that a preorder on a set S is a reflexive and transitive relation on S . We call a map f between two preordered sets *monotone (or increasing)* if it preserves the preorders, i.e., if $s \leq s'$ implies that $f(s) \leq f(s')$. Let (S, \leq) be a preordered set. We fix an order-preserving $(0, \infty)$ -action on S . We define the set of (*generalized*) *capacities on S* to be

$$(14) \quad \text{Cap}(S) := \{c \in [0, \infty]^S \mid c \text{ monotone and } (0, \infty)\text{-equivariant}\}.$$

We equip the set $[0, \infty]^S$ with the preorder

$$x \leq x' \iff x(s) \leq x'(s), \forall s \in S.$$

Let $\mathcal{G} \subseteq \text{Cap}(S)$.

Definition 28 (monotone and homogeneous monotone generation). *We say that \mathcal{G} monotonely (capacity-)generates iff for every $c \in \text{Cap}(S)$*

there exists a monotone function $F : [0, \infty]^{\mathcal{G}} \rightarrow [0, \infty]$, such that $c = F \circ \text{ev}_{\mathcal{G}}$. We say that \mathcal{G} homogeneously and monotonely (capacity-)generates iff the function F above can also be chosen to be homogeneous.

Remark (monotone versus homogeneous and monotone generation). The set \mathcal{G} monotonely generates if and only if it homogeneously and monotonely generates. The “only if”-direction follows by considering the monotonization (see p. 906 below) of the restriction of a function F as in Definition 28 to the image of $\text{ev}_{\mathcal{G}}$. Here we use that every $c \in \text{Cap}(S)$ is homogeneous, and thus $F|_{\text{im}(\text{ev}_{\mathcal{G}})}$ is homogeneous, as well as Remark 61 on p. 906.

Let $V := (V, \omega)$ be a symplectic vector space. Recall the definition (13) of the capacity c_j^V . The next result easily follows from McDuff’s solution of the embedding problem for ellipsoids in dimension 4. (See Section B.)

Theorem 29 (monotone generation for ellipsoids in dimension 4). *If $\dim V = 4$ then the set of all c_j^V (with $j \in \mathbb{N}_0$) monotonely generates (the generalized capacities on the category of ellipsoids Ell_V).*

This theorem provides a positive answer to the variant of Question 4 with “limit-min/max generating” replaced by “monotonely generating”. Monotone generation is (possibly nonstrictly) weaker than CHLS generation, since the pointwise limit of monotone functions is monotone. To deduce the theorem from McDuff’s result, we will characterize monotone generation in terms of almost order-reflexion.

3. Proof of Theorem 17(i,ii) (cardinality of the set of capacities)

In this section we prove Theorem 17(i,ii), based on a more general result, Theorem 43 below. That result states that the set of generalized capacities on a given (kn, k) -category \mathcal{C} has cardinality \beth_2 , provided that \mathcal{C} contains each disjoint union $M_a \sqcup M_{-a}$ for a suitable one-parameter family of manifolds with forms $(M_a, \omega_a)_{a \in A_0}$. A crucial hypothesis is that the collection of boundary helicities associated with $(M_a, \omega_a)_{a \in A_0}$, is an I -collection.

We also state Proposition 44, which provides sufficient conditions for this hypothesis to be satisfied.

3.1. (Boundary) helicity of an exact differential form

In this subsection we introduce the notion of helicity of an exact form, and based on this, the notion of boundary helicity.

Let $k, n \in \mathbb{N}_0$ be such that $n \geq 2$, N a closed⁴⁴ $(kn - 1)$ -manifold, O an orientation on N , and σ an exact k -form on N .

Definition 30 (helicity). *We define the helicity of (N, O, σ) to be the integral*

$$(15) \quad h(N, O, \sigma) := \int_{N, O} \alpha \wedge \sigma^{\wedge(n-1)},$$

where α is an arbitrary primitive of σ , and $\int_{N, O}$ denotes integration over N w.r.t. O .

We show that this number is well-defined, i.e., it does not depend on the choice of the primitive α . Let α and α' be primitives of σ . Then $\alpha' - \alpha$ is closed, and therefore

$$(\alpha' - \alpha) \wedge \sigma^{\wedge(n-1)} = (-1)^{k-1} d \left((\alpha' - \alpha) \wedge \alpha \wedge \sigma^{\wedge(n-2)} \right).$$

Here we used that $n \geq 2$. Using Stokes' Theorem and our assumption that N has no boundary, it follows that

$$\int_{N, O} (\alpha' - \alpha) \wedge \sigma^{\wedge(n-1)} = 0.$$

Therefore, the integral (15) does not depend on the choice of α .

Remark 31 (case k odd, case $n = 1$). The helicity vanishes if k is odd. This follows from the equality

$$\alpha \wedge (d\alpha)^{n-1} = \frac{1}{2} d \left(\alpha^{\wedge 2} \wedge (d\alpha)^{n-2} \right),$$

which holds for every even-degree form α , and from Stokes' Theorem. The helicity is not well-defined in the case $n = 1$. Namely, in this case $\dim N = k - 1$, and therefore every $(k - 1)$ -form is a primitive of the k -form 0. Hence the integral (15) depends on the choice of a primitive.

⁴⁴This means compact and without boundary.

Remark 32 (orientation). Denoting by \bar{O} the orientation opposite to O , we have

$$h(N, \bar{O}, \sigma) = -h(N, O, \sigma).$$

Remark 33 (rescaling). For every $c \in \mathbb{R}$ we have

$$h(N, O, c\sigma) = c^n h(N, O, \sigma).$$

This follows from a straight-forward argument.

Remark 34 (naturality). Let N and N' be closed $(kn - 1)$ -manifolds, O an orientation on N , σ an exact k -form on N , and $\varphi : N \rightarrow N'$ a (smooth) embedding. We denote

$$\varphi_*(N, O, \sigma) := (\varphi(N), \varphi_*O, \varphi_*\sigma)$$

(push-forwards of the orientation and the form). A straight-forward argument shows that

$$h(\varphi_*(N, O, \sigma)) = h(N, O, \sigma).$$

Remark 35 (helicity of a vector field). In the case $k = 2$ and $n = 2$ the integral (15) equals the helicity of a vector field V on a three-manifold N , which is dual to the two-form σ , via some fixed volume form. See [1, Definition 1.14, p. 125]. This justifies the name ‘‘helicity’’ for the function h .

The helicity of the boundary of a compact manifold equals the volume of the manifold. This is a crucial ingredient of the proofs of the main results and the content of the following lemma. Let M be a manifold, $N \subseteq M$ a sub-manifold, and ω a differential form on M . We denote by ∂M the boundary of M , and

$$(16) \quad \omega_N := \text{pullback of } \omega \text{ by the canonical inclusion of } N \text{ into } M.$$

If O is an orientation on M and N is contained in ∂M , then we define

$$(17) \quad O_N := O_N^M := \text{orientation of } N \text{ induced by } O.$$

Let $k, n \in \mathbb{N}_0$, such that $n \geq 2$, (M, O) be a compact oriented (smooth) manifold of dimension kn and ω an exact k -form on M .

Lemma 36 (Stokes’ theorem for helicity). *The following equality holds:*

$$\int_{M,O} \omega^{\wedge n} = h(\partial M, O_{\partial M}, \omega_{\partial M}).$$

Remark. The left hand side of this equality is $n!$ times the signed volume of M associated with O and ω .

Proof of Lemma 36. Choosing a primitive α of ω , we have

$$\omega^{\wedge n} = d(\alpha \wedge \omega^{\wedge(n-1)}),$$

and therefore, by Stokes’ Theorem,

$$\int_{M,O} \omega^{\wedge n} = \int_{\partial M, O_{\partial M}} \alpha \wedge \omega^{\wedge(n-1)} = h(\partial M, O_{\partial M}, \omega_{\partial M}).$$

This proves Lemma 36. □

This lemma has the following consequence. We denote

$$I_M := \{\text{connected component of } \partial M\}.$$

Definition 37 (boundary helicity). *We define the boundary helicity of (M, O, ω) to be the function*

$$h_M := h_{M,O,\omega} : I_M \rightarrow \mathbb{R}, \quad h_{M,O,\omega}(i) := h(i, O_i, \omega_i),$$

Corollary 38 (Stokes’ theorem for helicity). *The following equality holds:*

$$\int_{M,O} \omega^{\wedge n} = \sum_{i \in I_M} h(i, O_i, \omega_i).$$

Proof. This directly follows from Lemma 36. □

3.2. I -collections

An I -collection is collection $f = (f_a)_{a \in A_0}$ of real-valued functions with finite domains, such that the supremum of a certain set of numbers is less than 1. The set consists of all numbers C for which $A \cup B$ is nonempty, where A and B are certain sets of partitions, which depend on f and C . I -collections will occur in the generalized main result, Theorem 43 below. Namely, one hypothesis of this result is that the boundary helicities of a certain collection of manifolds and forms, are an I -collection.

Definition 39. Let I and I' be finite sets. an (I, I') -partition is a partition \mathcal{P} of the disjoint union $I \sqcup I'$, such that

$$(18) \quad \forall J \in \mathcal{P} : |J \cap I| = 1.$$

Let $f : I \rightarrow \mathbb{R}$, $f' : I' \rightarrow \mathbb{R}$, and $C \in (0, \infty)$. For every $J \subseteq I \sqcup I'$ we define

$$(19) \quad \sum_{J, f, f', C} := -C \sum_{i \in J \cap I} f(i) + \sum_{i' \in J \cap I'} f'(i').$$

A (f, f', C) -partition is an (I, I') -partition \mathcal{P} such that

$$(20) \quad \sum_{J, f, f', C} \geq 0, \quad \forall J \in \mathcal{P}.$$

Definition 40. Let I^+, I^-, I' be finite sets. We denote $I := I^+ \sqcup I^-$. an (I^+, I^-, I') -partition is a partition \mathcal{P} of $I \sqcup I'$ with the following properties:

- (a) There exists a unique element of \mathcal{P} that intersects both I^+ and I^- in exactly one point.
- (b) All other $J \in \mathcal{P}$ intersect I in exactly one point.

Let $f^\pm : I^\pm \rightarrow \mathbb{R}$, $f' : I' \rightarrow \mathbb{R}$, and $C \in (0, \infty)$. We denote by $f := f^+ \sqcup f^- : I \rightarrow \mathbb{R}$ the disjoint union of functions.⁴⁵ A (f^+, f^-, f', C) -partition is an (I^+, I^-, I') -partition satisfying (20).

Remark 41. Every (I^+, I^-, I') -partition \mathcal{P} satisfies

$$|\mathcal{P}| = |I| - 1.$$

Let A_0 be an interval and I a collection of finite sets indexed by A_0 , i.e., a map from A_0 to the class of all finite sets. We denote $I_a := I(a)$. Let $f = (f_a : I_a \rightarrow \mathbb{R})_{a \in A_0}$ be a collection of functions. We define

$$(21) \quad C_0^f := \sup \{ C \in (0, \infty) \mid \exists a, a' \in A_0 : a > a', \exists (f_a, f_{a'}, C)\text{-partition} \},$$

$$(22) \quad C_1^f := \sup \{ C \in (0, \infty) \mid \exists a, a' \in A_0 \cap (0, \infty) : a < a', \exists (f_a, f_{-a}, f_{a'}, C)\text{-partition} \}.$$

Here we use the convention that $\sup \emptyset := 0$.

⁴⁵This is the function defined by $f(i) := f^\pm(i)$ if $i \in I^\pm$.

Definition 42 (I-collection). *We call f an I-collection iff the following holds:*

$$(23) \quad C_0^f < 1,$$

$$(24) \quad C_1^f < 1.$$

Remark. *The condition of being an I-collection is invariant under rescaling by some positive constant.*

3.3. Cardinality of the set of capacities in a more general setting, sufficient conditions for being an I-collection, proof of Theorem 17(i,ii)

Theorem 17(i,ii) is a special case of the following more general result. We call a k -form ω on a kn -manifold *maxipotent* iff $\omega^{\wedge n} = \omega \wedge \dots \wedge \omega$ does not vanish anywhere.⁴⁶ In this case we denote

$$O_\omega := \text{orientation on } M \text{ induced by } \omega^{\wedge n}.$$

Recall that B, Z denote the unit ball and the standard symplectic cylinder, ω_{st} the standard symplectic form, $c_{M,\omega}$ the domain-embedding capacity for (M, ω) as in (3), and w the Gromov width as in (1).

Theorem 43 (cardinality of the set of (normalized) capacities, more general setting). *The following holds:*

(i) *Let $k, n \in \{2, 3, \dots\}$ with k even, and $\mathcal{C} = (\mathcal{O}, \mathcal{M})$ be a (kn, k) -form category. Then the cardinality of $\text{Cap}(\mathcal{C})$ equals \beth_2 , provided that there exist an interval A_0 around 0 of positive length, and a collection $(M_a, \omega_a)_{a \in A_0}$ of objects of $\Omega^{kn, k}$, such that for every $a \in A_0$, M_a is nonempty, compact, and 1-connected,⁴⁷ ω_a is maxipotent and exact, and the following holds:*

- (a) $(W_a, \eta_a) := (M_a \sqcup M_{-a}, \omega_a \sqcup \omega_{-a}) \in \mathcal{O}$, for every $a \in A_0 \cap (0, \infty)$.
- (b) We denote by I_a the set of connected components of ∂M_a , and $I := (I_a)_{a \in A_0}$. The collection of boundary helicities $f := (h_{M_a, O_{\omega_a}, \omega_a})_{a \in A_0}$ is an I-collection.

(ii) *Let $n \in \{2, 3, \dots\}$ and $\mathcal{C} = (\mathcal{O}, \mathcal{M})$ be a $(2n, 2)$ -form category that contains the objects B and Z . Then the cardinality of $\mathcal{NCap}(\mathcal{C})$ equals*

⁴⁶See Remark 22 for the relation between maxipotency and nondegeneracy.

⁴⁷This means connected and simply connected.

\beth_2 , provided that there exist A_0 and $(M_a, \omega_a)_{a \in A_0}$ as in (i), such that also the following holds:

- (a) $\sup_{a \in A_0} w(M_a, \omega_a) < 1$
- (b) $\sup_{a \in A_0} c_{M_a, \omega_a}(Z, \omega_{st}) \leq \pi$

We will prove this theorem in Section 4. The idea of the proof is to consider the family of capacities

$$c_A := \sup_{a \in A} c_{W_a, \eta_a}, \quad A \in \mathcal{P}(A_0 \cap (0, \infty)).$$

Hypothesis (ib) implies that there exists $c_0 < 1$ such that for all $a \neq a' \in (0, \infty)$ and $c \geq c_0$, the pair $(W_a, c\eta_a)$ does not embed into $(W_{a'}, \eta_{a'})$. See the explanations below. It follows that

$$\sup \{c_A(W_a, \eta_a) \mid a \in A_0 \cap (0, \infty) \setminus A\} < 1, \quad \forall A.$$

Since also $c_A(W_a, \eta_a) = 1$, for every $a \in A$, it follows that

$$c_A \neq c_{A'}, \quad \text{if } A \neq A'.$$

Since the cardinality of $\mathcal{P}((0, \infty))$ equals \beth_2 , it follows that the cardinality of $\mathcal{Cap}(\mathcal{C})$ is at least \beth_2 . On the other hand, we denote by S the set of isomorphism classes of symplectic manifolds. This set has cardinality \beth_1 . Since $\mathcal{Cap}(\mathcal{C})$ can be viewed as a subset of $[0, \infty]^S$, it has cardinality at most \beth_2 , hence equal to \beth_2 .

A refined version of this argument shows that $|\mathcal{N}\mathcal{Cap}(\mathcal{C})| = \beth_2$. For this we normalize each capacity c_A , by replacing it by the maximum of c_A and the Gromov width. Hypothesis (iia) guarantees that the modified capacities are still all different from each other. Hypothesis (iib) guarantees that they are normalized.

To understand the reason why no big multiple of (W_a, η_a) embeds into $(W_{a'}, \eta_{a'})$, consider the case in which each M_a is a spherical shell in a symplectic vector space, with inner radius 1 and outer radius $r + a$ for some fixed $r > 1$. Assume that $(M_a, c\omega_a)$ embeds into $(M_{a'}, \omega_{a'})$ in such a way that the image of the inner boundary sphere of M_a wraps around the inner boundary sphere of $M_{a'}$. By Corollary 38 (Stokes' Theorem for helicity) and Remark 32 the difference of the helicities of these spheres equals the enclosed volume on the right hand side. Since this volume is nonnegative, it follows that $c \geq 1$. Using our hypothesis (ib) that the collection of boundary helicities is an I -collection, it follows that $a \leq a'$.

It follows that if $a > a'$ then no multiple of W_a (symplectically) embeds into $W_{a'}$ in such a way that the inner boundary sphere of M_a wraps around one of the two inner boundary spheres of $W_{a'}$. Figure 1 illustrates this. In contrast with this, Figure 2 shows a possible embedding. In this case our helicity hypothesis (ib) implies that the rescaling factor is small.




Figure 1: If $a > a'$ then no multiple of the red spherical shell M_a (symplectically) embeds into the blue shell $M_{a'}$ in such a way that the inner boundary sphere of the red shell wraps around the inner boundary sphere of the blue shell, since our helicity hypothesis (ib) forces the rescaling factor to be at least 1.

If $a < a'$ then M_a embeds into $M_{a'}$ (without rescaling). However, there is not enough space left for M_{-a} . See Figure 3.

In the proof of Theorem 17(i) we will use the following sufficient criterion for condition (ib) of Theorem 43. For every finite set S and every function $f : S \rightarrow \mathbb{R}$ we denote

$$(25) \quad \sum f := \sum_{s \in S} f(s).$$

Let A_0 be an interval, $I := (I_a)_{a \in A_0}$ a collection of finite sets, and $f = (f_a : I_a \rightarrow \mathbb{R})_{a \in A_0}$ a collection of functions. We define the disjoint unions of I and

f to be

$$\begin{aligned} \sqcup I &:= \sqcup_{a \in A_0} I_a := \{(a, i) \mid a \in A_0, i \in I_a\}, \\ \sqcup f &: \sqcup I \rightarrow \mathbb{R}, \quad \sqcup f(a, i) := f_a(i). \end{aligned}$$




Figure 2: A possible embedding of $(W_a, c\eta_a)$ into $(W_{a'}, \eta_{a'})$ in the case $a > a'$. The constant c needs to be small (even if a is close to a'), since the volume of the hole enclosed by the image of M_a equals minus c times the helicity of the inner boundary sphere of M_a . Here we use again our helicity hypothesis (ib).




Figure 3: An attempt for an embedding of W_a into $M_{a'}$ in the case $a < a'$ (without rescaling). The image of M_{-a} overlaps itself, since there is not enough space left in $M_{a'}$.

Proposition 44 (sufficient conditions for being an I -collection).

The collection f is an I -collection if there exists $\ell \in \mathbb{N}_0$, such that the following holds:

(i) For all $a \in A_0$ we have

$$\begin{aligned}
 (26) \quad & |I_a| = \ell, \\
 (27) \quad & f_a \geq -1, \\
 (28) \quad & f_a^{-1}(-1) \neq \emptyset, \\
 (29) \quad & |f_a^{-1}((0, \infty))| = 1, \\
 (30) \quad & \sum f_a \leq 1.
 \end{aligned}$$

(ii) For all $a, a' \in A_0$ we have

$$(31) \quad \sum f_a > \sum f_{a'}, \quad \text{if } a > a'.$$

(iii) We have

$$(32) \quad \sup \left(\text{im} \left(\bigsqcup f \right) \cap (-\infty, 0] \right) < -1 + \inf_{a \in A_0} \sum f_a.$$

If $\ell \geq 4$ then we have

$$(33) \quad \sup \bigsqcup f < 2 \inf \left(\text{im} \left(\bigsqcup f \right) \cap (0, \infty) \right) + 1.$$

We will prove this proposition in Section 5.

Remark. The conditions (28,29) imply that $\ell \geq 2$.

Proof of Theorem 17(i,ii). **(i):** We choose V, Ω, K, r as in the hypothesis. We define

$$(34) \quad \omega := \Omega^{\oplus n}.$$

Since by hypothesis, k is even and Ω is a volume form, the form ω is maxipotent, i.e., $\omega^{\wedge n}$ is a volume form. We denote by O the orientation on V^n

induced by this form. Since by hypothesis, K is nonempty and strictly star-shaped around 0, its interior contains 0. It follows that

$$(35) \quad C := \int_{K,O} \omega^{\wedge n} > 0.^{48}$$

By hypothesis, we have

$$(36) \quad a_1 := \min \{r - 1, \sqrt[kn]{2} - r\} > 0.$$

We choose $a_0 \in (0, a_1)$ and define $A_0 := [-a_0, a_0]$. For every $a \in A_0$ we define

$$(37) \quad M_a := (r + a)K \setminus \text{int } K,$$

$$(38) \quad \omega_a := C^{-\frac{1}{n}} \omega|_{M_a},$$

$$(39) \quad I_a := \{\text{connected component of } \partial M_a\},$$

$$I := (I_a)_{a \in A_0}.$$

The form ω_a is well-defined, since $C > 0$. We check the hypotheses of Theorem 43(i). Let $a \in A_0$. The set M_a is compact. Since K is strictly starshaped around 0, M_a is a smooth submanifold of V^n that continuously deformation retracts onto ∂K . The manifold ∂K is homeomorphic to the sphere S_1^{kn-1} . Since by hypothesis $k, n \geq 2$, this sphere is 1-connected. Hence the same holds for M_a . The form Ω is exact. Hence the same holds for ω and thus for ω_a .

Condition (ia) is satisfied by our hypothesis and the rescaling property for a (kn, k) -form category. We show that the collection of boundary helicities

$$(40) \quad f := (f_a := h_{M_a, O_{\omega_a}, \omega_a})_{a \in A_0}$$

satisfies (ib). We check the hypotheses of Proposition 44. Let $s \in (0, \infty)$. We denote by O^s the orientation on $\partial(sK)$ induced by O and sK . By Lemma 36 we have

$$(41) \quad h(\partial(sK), O^s, \omega_{\partial(sK)}) = \int_{sK, O} \omega^{\wedge n} = C s^{kn}.$$

⁴⁸Here we view V^n as a manifold and ω as a differential form on it.

For every connected component i of ∂M_a we denote by O_i the orientation of i induced by O, M_a . Using (41,38) and Remarks 33,32, we obtain

$$(42) \quad h(i, O_i, (\omega_a)_i) = \begin{cases} (r+a)^{kn}, & \text{for } i = \partial((r+a)K), \\ -1, & \text{for } i = \partial K. \end{cases}$$

Here we used that the orientation of ∂K induced by O and M_a is the opposite of O^1 . It follows that

$$(43) \quad \sum f_a := \sum_{i \in I_a} f_a(i) = -1 + (r+a)^{kn} \in [-1 + (r-a_0)^{kn}, -1 + (r+a_0)^{kn}], \forall a \in A_0.$$

Since $a_0 < a_1 \leq r-1$, we have $-1 + (r-a_0)^{kn} > 0$. Hence by (43), we have $\inf_{a \in A_0} \sum f_a > 0$. Using (42), it follows that condition (32) is satisfied.

Since $a_0 < a_1 \leq \sqrt[kn]{2} - r$, we have $-1 + (r+a_0)^{kn} < 1$. Using (43), it follows that $\sup_{a \in A_0} \sum f_a < 1$. Hence inequality (30) is satisfied. The collection f also satisfies the other hypotheses of Proposition 44. Applying this proposition, it follows that f is an I -collection. Hence condition (ib) is satisfied.

Therefore, all hypotheses of Theorem 43(i) are satisfied. Applying this theorem, it follows that the cardinality of $Cap(\mathcal{C})$ equals \beth_2 . This proves Theorem 17(i).

To prove (ii), assume that the hypotheses of this part of the theorem are satisfied. We choose $r \in (1, \sqrt[2n]{2})$ satisfying (6). We define $V := \mathbb{R}^2, \Omega$ to be the standard area form on $\mathbb{R}^2, K := \bar{B}_1^{2n}$, and a_1 as in (36). We choose $a_0 \in (0, a_1)$, and define $A_0 := [-a_0, a_0]$ and (M_a, ω_a) as in (37,38). The triple (V, Ω, K) satisfies the conditions of part (i) of Theorem 17. Hence by what we proved above, the collection $(M_a, \omega_a), a \in A_0$, satisfies the conditions of Theorem 43(i). Let $a \in A_0$. By (34) we have $\omega = \omega_{st}$. Using (35), it follows that

$$C = \int_{K=\bar{B}_1^{2n}} \omega_{st}^{\wedge n} = \pi^n,$$

and therefore,

$$(44) \quad (M_a, \omega_a) = \left(Sh_{r+a}, \frac{1}{\pi} \omega_{st} | M_a \right).$$

We check condition (iia) of Theorem 43. Let $a \in A_0$. It follows from (44) that the symplectic volume of (M_a, ω_a) is $\frac{-1+(r+a)^{2n}}{\pi^n}$ times the volume of

the unit ball. Therefore, we have

$$w(M_a, \omega_a) \leq \sqrt[n]{-1 + (r + a)^{2n}}.$$

Using the inequalities $a \leq a_0 < a_1 \leq \sqrt[2n]{2} - r$, it follows that condition (iia) is satisfied.

We check (iib). Let $a \in A_0$. Since $r + a \geq r - a_0 > r - a_1 \geq 1$, the shell Sh_{r+a} contains the sphere S_{r+a}^{2n-1} . Using $r + a > 1$, $\omega_a = \frac{1}{\pi}\omega_{st}$, and $n \geq 2$, it follows from [15, Corollary 5, p. 8] (spherical nonsqueezing) that $(M_a, b\omega_a)$ does not symplectically embed into Z for any $b \geq \pi$. Hence (iib) holds.

Therefore, all hypotheses of Theorem 43(ii) are satisfied. Applying this part of the theorem, it follows that the cardinality of $\mathcal{NCap}(\mathcal{C})$ equals \beth_2 . This proves Theorem 17(ii). □

4. Proof of Theorem 43 (cardinality of the set of capacities, more general setting)

As mentioned, the idea of proof of Theorem 43 is that our helicity hypothesis (ib) and Stokes' Theorem for helicity imply that for $a \neq a'$ only small multiples of (W_a, η_a) embed into $(W_{a'}, \eta_{a'})$. The idea behind this is that every embedding φ of M_a into $M_{a'}$ gives rise to a partition of the disjoint union of the sets of connected components of ∂M_a and $\partial M_{a'}$. The elements of this partition consist of components that lie in the same connected component of the complement of $\varphi(\text{Int } M)$. Here $\text{Int } M$ denotes the interior of M as a manifold with boundary, and we identify each component of ∂M_a with its image under φ .

Stokes' Theorem for helicity implies that the inequality (20) is satisfied. Together with a similar argument in which we consider embeddings of W_a into $M_{a'}$, it follows that the partition satisfies the conditions of Definitions 39,40. Combining this with our helicity hypothesis (ib), it follows that indeed only small multiples of W_a embed into $W_{a'}$.

Lemmata 47 and 49 below will be used to make this argument precise. To formulate the first lemma, we need the following.

Remark 45 (pullback of relation). Let S', S be sets, R a relation on S , and $f : S' \rightarrow S$ a map. Denoting by \times the Cartesian product of maps, the set

$$R' := f^*R := (f \times f)^{-1}(R)$$

is a relation on S' . If R is reflexive/ symmetric/ transitive, then the same holds for R' .

Let X be a topological space. We define

$$(45) \quad \mathcal{C}_X := \{\text{path-connected subset of } X\}$$

and the relation \sim_X on \mathcal{C}_X by

$$(46) \quad A \sim_X B : \iff \exists \text{ continuous path starting in } A \text{ and ending in } B.$$

This is an equivalence relation.

Let M and M' be topological manifolds of the same dimension, and $\varphi : M \rightarrow M'$ a topological embedding, i.e., a homeomorphism onto its image. We denote by $\text{Int}(M)$ and ∂M the interior and the boundary of M as a manifold with boundary. We denote

$$(47) \quad I := I_M := \{\text{connected component of } \partial M\}, \quad I' := I_{M'}$$

$$(48) \quad P := M' \setminus \varphi(\text{Int}(M)).$$

We define

$$\Phi : \mathcal{P}(M) \rightarrow \mathcal{P}(M'), \quad \Phi(A) := \text{image of } A \text{ under } \varphi,$$

$$\Psi : I \sqcup I' \rightarrow \mathcal{P}(P), \quad \Psi := \Phi \text{ on } I, \quad \Psi := \text{id on } I',$$

$$\sim^\varphi := \Psi^* \sim_P,$$

$$\mathcal{P}^\varphi := \text{partition of } I \sqcup I' \text{ associated with } \sim^\varphi.$$

Remark 46 (partition induced by embedding). For every path-component P_0 of P we define

$$(49) \quad \begin{aligned} J^\varphi(P_0) &:= \Psi^{-1}(\mathcal{P}(P_0)) \\ &= \{i \in I \mid \Phi(i) \in \mathcal{P}(P_0)\} \sqcup (I' \cap \mathcal{P}(P_0)). \end{aligned}$$

The map

$$J^\varphi : \{\text{path-component } P_0 \text{ of } P : J^\varphi(P_0) \neq \emptyset\} \rightarrow \mathcal{P}^\varphi$$

is well-defined and a bijection.

For every field F and $i \in \mathbb{N}_0$ we denote by $H_i(M; F)$ the degree i singular homology of M with coefficients in F .

Lemma 47 (partition associated with an embedding). *Assume that M, M' are compact, M' is connected, $\partial M' \neq \emptyset$, and that there exists a field F , for which $H_1(M'; F)$ vanishes. Then the following holds:*

- (i) *If M is nonempty and connected then \mathcal{P}^φ is an $(I_M, I_{M'})$ -partition.*
- (ii) *If M consists of precisely two connected components M^+ and M^- then \mathcal{P}^φ is an $(I_{M^+}, I_{M^-}, I_{M'})$ -partition.*

Recall that the first statement means that condition (18) is satisfied, i.e., $|J \cap I_M| = 1$ for every $J \in \mathcal{P}^\varphi$. The idea of proof of the inequality ≤ 1 is the following. Each J corresponds to a path-component P_0 of the complement of $\varphi(\text{Int } M)$. Suppose that there exists J that intersects I_M in at least two points i_0, i_1 (= components of ∂M). Then there is a path in P_0 joining $\varphi(i_0)$ and $\varphi(i_1)$. By connecting this path with a path in $\varphi(M)$ with the same endpoints, we obtain a loop in M' that intersects i_0 and i_1 in one point each. See Figure 4.




Figure 4: The blue region is the image of M under φ , and the red and green regions are the path-components of the complement of $\varphi(\text{Int } M)$. The red region contains the images of two connected components i_0, i_1 of the boundary of M . The yellow loop intersects these images in one point each.

Hence the algebraic intersection number of this loop with i_0 equals 1. In particular, it represents a nonzero first homology class. Hence the hypothesis that the first homology of M' vanishes, is violated. It follows that $|J \cap I_M| \leq 1$.

In order to make this argument precise one needs to ensure that the algebraic intersection number equals the “naïve intersection number”. For simplicity, we therefore use an alternative method of proof, which is based on a certain Mayer-Vietoris sequence for singular homology. We need the following.

Remark 48 (embedding is open, boundary). We denote by $\partial^X S$ the boundary of a subset S of a topological space X . Let M, M' be topological manifolds of the same dimension n , and $\varphi : M \rightarrow M'$ an injective continuous map. By invariance of the domain, in every pair of charts for $\text{Int } M$ and M' , the map φ sends every open subset of \mathbb{R}^n to an open subset of \mathbb{R}^n . It follows that the set $\varphi(\text{Int } M)$ is open in M' . This implies that

$$\varphi(\partial M) \subseteq \partial^{M'} \varphi(\text{Int } M),$$

and if M is compact, then equality holds.

Suppose now that M is nonempty and compact, $\partial M = \emptyset$, and M' is connected. Then M' has no boundary, either. To see this, observe that $\varphi(M)$ is compact, hence closed in M' . Since $M = \text{Int } M$, as mentioned above, $\varphi(M)$ is also open. Since M' is connected, it follows that $\varphi(M) = M'$. Since in every pair of charts for M and M' , φ sends every open subset of \mathbb{R}^n to an open subset of \mathbb{R}^n , it follows that $\partial M' = \emptyset$.

Proof of Lemma 47. Assume that M, M' are compact, $M \neq \emptyset$, M' is connected, and $\partial M' \neq \emptyset$. We denote

$$I := I_M, \quad I' := I_{M'}, \quad P := M' \setminus \varphi(\text{Int}(M)),$$

and by k the number of connected components of M .

Claim 1. *We have*

$$(50) \quad |\mathcal{P}^\varphi| = |I| + 1 - k.$$

Proof of Claim 1. Let P_0 be a path-component of P .

Claim 2. *P_0 intersects $\varphi(\partial M)$.*

Proof of Claim 2. By Remark 48 we have $\partial M \neq \emptyset$. Since by hypothesis, M' is connected, there exists a continuous path $x' : [0, 1] \rightarrow M'$ that starts in

P_0 and ends at $\varphi(\partial M)$. Since M is compact, the same holds for ∂M , and hence for $\varphi(\partial M)$. Hence the minimum

$$t_0 := \min \{t \in [0, 1] \mid x'(t) \in \varphi(\partial M)\}$$

exists. By Remark 48 the set $\varphi(\text{Int } M)$ is open in M' . It follows that $x'(t_0) \notin \varphi(\text{Int } M)$, and hence $x'([0, t_0]) \subseteq P = M' \setminus \varphi(\text{Int } M)$. (In the case $t_0 = 0$ this holds, since $x'(0) \in P_0 \subseteq P$.) It follows that $x'(t_0) \in P_0$. Since also $x'(t_0) \in \varphi(\partial M)$, it follows that $P_0 \cap \varphi(\partial M) \neq \emptyset$. This proves Claim 2. \square

Claim 2 implies that the set $J^\varphi(P_0)$ (defined as in (49)) is nonempty. Hence by Remark 46 we have

$$(51) \quad |\{\text{path-component of } P\}| = |\mathcal{P}^\varphi|.$$

By M. Brown's Collar Neighbourhood Theorem [2] (see also [4, Theorem, p. 180]) there exists an open subset V of M and a (strong) deformation retraction h of V onto ∂M . We define

$$A := \varphi(M), \quad B := M' \setminus \varphi(M \setminus V).$$

Extending $\varphi \circ h_t \circ \varphi^{-1} : \varphi(V) \rightarrow \varphi(V)$ by the identity, we obtain a map $h' : [0, 1] \times B \rightarrow B$. Since by Remark 48, the restriction of φ to $\text{Int } M$ is open, the map h' is continuous, and therefore a deformation retraction of B onto P .

We choose a field F as in the hypothesis, and denote by H_i singular homology in degree i with coefficients in F . Since P is a deformation retract of B , these spaces have isomorphic H_0 . Combining this with (51), it follows that

$$(52) \quad \begin{aligned} |\mathcal{P}^\varphi| &= |\{\text{path-component of } P\}| \\ &= \dim H_0(P) \\ &= \dim H_0(B). \end{aligned}$$

The interiors of A and B cover M' . Therefore, the Mayer-Vietoris Theorem implies that there is an exact sequence

$$\dots \rightarrow H_1(M') \rightarrow H_0(A \cap B) \rightarrow H_0(A) \oplus H_0(B) \rightarrow H_0(M') \rightarrow 0.$$

Since by hypothesis, $H_1(M') = 0$, it follows that

$$(53) \quad \dim H_0(B) = \dim H_0(A \cap B) + \dim H_0(M') - \dim H_0(A).$$

Since $A \cap B = \varphi(V)$ and φ is a homeomorphism onto its image, we have $H_0(A \cap B) \cong H_0(V)$. Since V deformation retracts onto ∂M , we have $H_0(V) \cong H_0(\partial M)$, hence $H_0(A \cap B) \cong H_0(\partial M)$. Since ∂M is a topological manifold, its path-components are precisely its connected components. Recalling the definition (47) of I , it follows that

$$(54) \quad \dim H_0(A \cap B) = |I|.$$

Since by hypothesis M' is connected, we have

$$(55) \quad \dim H_0(M') = 1.$$

Since $A := \varphi(M)$, we have $H_0(A) \cong H_0(M)$, and therefore

$$\dim H_0(A) = k.$$

Combining this with (52,53,54,55), equality (50) follows. This proves Claim 1. □

Remark 46 and Claim 2 imply that every element of \mathcal{P}^φ intersects I .

We prove (i). Assume that M is connected. Then by Claim 1, we have $|\mathcal{P}^\varphi| = |I|$. It follows that $|J \cap I| = 1$, for every $J \in \mathcal{P}^\varphi$. Hence \mathcal{P}^φ is an (I, I') -partition. This proves (i).

Assume now that M^\pm are as in the hypothesis of (ii). By Claim 1 we have $|\mathcal{P}^\varphi| = |I| - 1$. Since every element of \mathcal{P}^φ intersects I , it follows that there exists a unique $J_0 \in \mathcal{P}^\varphi$, such that $|J_0 \cap I| = 2$, and

$$(56) \quad |J \cap I| = 1, \quad \forall J \in \mathcal{P}^\varphi \setminus \{J_0\}.$$

By Remark 46 there exists a unique path-component P_0 of P , such that $J_0 = J^\varphi(P_0)$.

Claim 3. *We have*

$$J_0 \cap I^- \neq \emptyset \neq J_0 \cap I^+.$$

Proof of Claim 3. We denote by P_0^+ the path-component of $M' \setminus \varphi(\text{Int}(M^+))$ containing P_0 . Assume by contradiction that $P_0^+ \cap \varphi(M^-) = \emptyset$. Then we

have

$$P_0^+ = P_0, \quad J^{\varphi|M^+}(P_0^+) = J^{\varphi}(P_0) = J_0, \quad J_0 \cap I = J_0 \cap I^+.$$

Since $|J_0 \cap I| = 2$, we obtain a contradiction with (i), with I, φ replaced by $I^+, \varphi|M^+$. Hence we have

$$P_0^+ \cap \varphi(M^-) \neq \emptyset.$$

It follows that there exists a continuous path $x' : [0, 1] \rightarrow M' \setminus \varphi(\text{Int}(M^+))$ that starts at P_0 and ends at $\varphi(M^-)$. Since M is compact, the same holds for $\varphi(M^-)$. Hence the minimum

$$t_0 := \min \{t \in [0, 1] \mid x'(t) \in \varphi(M^-)\}$$

exists. By Remark 48 the set $\varphi(\text{Int } M^-)$ is open. It follows that $x'(t_0) \notin \varphi(\text{Int } M^-)$, hence $x'([0, t_0]) \subseteq P$, and therefore

$$(57) \quad x'(t_0) \in P_0.$$

On the other hand $x'(t_0) \in \varphi(M^-) \subseteq \overline{\varphi(\text{Int } M^-)}$, and therefore

$$x'(t_0) \in \partial^{M'} \varphi(\text{Int } M^-) = \varphi(\partial M^-).$$

Here we used Remark 48. Combining this with (57), it follows that $P_0 \cap \varphi(\partial M^-) \neq \emptyset$, and therefore $J_0 \cap I^- \neq \emptyset$.

An analogous argument shows that $J_0 \cap I^+ \neq \emptyset$. This proves Claim 3. □

By Claim 3 and (56) \mathcal{P}^φ is an (I^+, I^-, I') -partition. This proves (ii) and completes the proof of Lemma 47. □

The second ingredient of the proof of Theorem 43 is the following. Let $k, n \in \mathbb{N}_0$ with $n \geq 2$, M, M' be compact (smooth) manifolds of dimension kn , ω, ω' exact maxipotent k -forms on M, M' , $c \in (0, \infty)$, and $\varphi : M \rightarrow M'$ a (smooth) orientation preserving embedding that intertwines $c\omega$ and ω' . We denote by O, O' the orientations of M, M' induced by ω, ω' . Recall Definitions 30,37 of (boundary) helicity.

Lemma 49 (helicity inequality). *Condition (20) holds with $\mathcal{P} = \mathcal{P}^\varphi$, $f = h_{M,O,\omega}$, $f' = h_{M',O',\omega'}$, and $C := c^n$.*

The reason for this is that the left hand side of (20) is the volume of the path-component of the complement of $\varphi(\text{Int } M)$, determined by J . To make this precise, we need the following.

Remark 50. Let X and X' be topological spaces and $f : X \rightarrow X'$ be continuous. Recall the definitions (45,46) of \mathcal{C}_X and \sim_X .

(i) The map

$$f_* : \mathcal{C}_X \rightarrow \mathcal{C}_{X'}, \quad f_*(A) := f(A),$$

is well-defined. Furthermore, we have

$$f_* \times f_*(\sim_X) \subseteq \sim_{X'} .$$

(ii) Assume that $X = X'$ and for every $x \in X$ there exists a continuous path from x to $f(x)$. Then for every pair $A, B \in \mathcal{C}_X$ we have

$$f_*(A) \sim_{f(X)} f_*(B) \Rightarrow A \sim_X B.$$

This follows from transitivity of \sim_X .

Proof of Lemma 49. Let $M, O, \omega, M', O', \omega', c, \varphi$ be as in the hypothesis. We define $I = I_M, I' = I_{M'}$ as in (47). Consider first the **case** in which

$$(58) \quad \varphi(\partial M) \cap \partial M' = \emptyset.$$

Then the set

$$P := M' \setminus \varphi(\text{Int } M)$$

is a smooth submanifold of M' . Let $i \in I$. We denote $\hat{i} := \varphi(i)$. We define O_N^M as in (17), and abbreviate

$$O_i := O_i^M, \quad O_{\hat{i}} := (O')_{\hat{i}}^P.$$

Recall that \overline{O} denotes the orientation opposite to O . Since φ intertwines O, O' , and $P, \varphi(M)$ lie on opposite sides of \hat{i} , we have

$$(59) \quad (\varphi|i)_* \overline{O_i} = \overline{(\varphi|i)_* O_i} = O_{\hat{i}}.$$

Recall the definition (16) of ω_N . Since φ intertwines $c\omega, \omega'$, we have

$$(60) \quad (\varphi|i)_* c\omega_i = \omega'_{\hat{i}}.$$

We have

$$\begin{aligned}
 -c^n h(i, O_i, \omega_i) &= c^n h(i, \overline{O_i}, \omega_i) && \text{(by Remark 32)} \\
 &= h(i, \overline{O_i}, c\omega_i) && \text{(by Remark 33)} \\
 &= h((\varphi|i)_*(i, \overline{O_i}, c\omega_i)) && \text{(by Remark 34)} \\
 (61) \qquad &= h(\widehat{i}, O_{\widehat{i}}, \omega'_{\widehat{i}}) && \text{(using } \widehat{i} = \varphi(i), (59,60)).
 \end{aligned}$$

Let P_0 be a path-component of P . We define $J := J^\varphi(P_0)$ as in (49). Using $h_{M,O,\omega}(i) = h(i, O_i, \omega_i)$ and (61), we have

$$\begin{aligned}
 & -c^n \sum_{i \in J \cap I} h_{M,O,\omega}(i) + \sum_{i' \in J \cap I'} h_{M',O',\omega'}(i') \\
 &= \sum_{\widehat{i} \in I_{P_0}} h_{P_0,O'|_{P_0},\omega'|_{P_0}}(\widehat{i}) \\
 &= \int_{P_0,O'|_{P_0}} \omega'^n \qquad \text{(using Corollary 38)} \\
 &\geq 0.
 \end{aligned}$$

Hence the statement of Lemma 49 holds in the case (58).

Consider now the general situation. Let $(K_i, r_i)_{i \in I}$ be a collection, where for each $i \in I$, K_i is a compact connected neighbourhood of i that is a (smooth) submanifold of M (with boundary), and $r_i : K_i \rightarrow i$ is a continuous retraction, such that the sets K_i , $i \in I$, are disjoint. We denote by $\text{int}(K_i)$ the interior of K_i in M . We define

$$\begin{aligned}
 \widetilde{M} &:= M \setminus \bigcup_{i \in I} \text{int}(K_i), \\
 \widetilde{\varphi} &:= \varphi|_{\widetilde{M}}, \\
 \widetilde{I}_i &:= I_{K_i} \setminus \{i\}, \forall i \in I, \quad \widetilde{I} := I_{\widetilde{M}}.
 \end{aligned}$$

We define

$$(62) \qquad \widetilde{\sim} : \mathcal{P}(I \sqcup I') \rightarrow \mathcal{P}(\widetilde{I} \sqcup I'), \quad \widetilde{J} := (J \setminus I) \cup \bigcup_{i \in J \cap I} \widetilde{I}_i.$$

The set \widetilde{M} is a submanifold of M , and

$$(63) \qquad \widetilde{\varphi}(\partial \widetilde{M}) \cap \partial M' = \emptyset.$$

Claim 1.

$$(64) \quad \mathcal{P}^{\tilde{\varphi}} = \tilde{\mathcal{P}}^{\tilde{\varphi}} := \{\tilde{J} \mid J \in \mathcal{P}^{\varphi}\}.$$

Proof of Claim 1. We define

$$\tilde{P} := M' \setminus \varphi(\text{Int}(\tilde{M})), \quad r : \tilde{P} \rightarrow P,$$

by setting

$$r := \begin{cases} \varphi \circ r_i \circ \varphi^{-1} & \text{on } \varphi(K_i), \text{ with } i \in I, \\ r = \text{id} & \text{on } M' \setminus \varphi(M). \end{cases}$$

Since the sets K_i are disjoint, the map r is well-defined. Since by hypothesis, φ is an embedding between two manifolds of the same dimension, the map r is continuous. Let $i \in I$. Since K_i is path-connected and r_i is a retraction onto the subset i of K_i , the hypotheses of Remark 50(ii) are satisfied with $f = r$. Applying this remark, it follows that for every pair \tilde{A}, \tilde{B} of path-connected subsets of \tilde{P} we have

$$\tilde{A} \sim_{\tilde{P}} \tilde{B} \iff r(\tilde{A}) \sim_{r(\tilde{P})=P} r(\tilde{B}).$$

This implies that if $i_0, i_1 \in I, \tilde{i}_k \in \tilde{I}_{i_k}$, for $k = 0, 1$, and $i'_0, i'_1 \in I'$ then

$$\tilde{i}_0 \sim_{\tilde{\varphi}} \tilde{i}_1 \iff i_0 \sim_{\varphi} i_1, \quad i'_0 \sim_{\tilde{\varphi}} i'_1 \iff i'_0 \sim_{\varphi} i'_1, \quad \tilde{i}_0 \sim_{\tilde{\varphi}} i'_0 \iff i_0 \sim_{\varphi} i'_0.$$

Equality (64) follows. This proves Claim 1. □

We abbreviate

$$h_M := h_{M,O,\omega}.$$

Recall the definition (19). Using (63), by what we already proved, condition (20) holds with I replaced by $\tilde{I}, \mathcal{P} := \mathcal{P}^{\tilde{\varphi}}, f := h_{\tilde{M}}, f' := h_{M'}$, and $C := c^n$. Using Claim 1, it follows that

$$(65) \quad \sum_{\tilde{J}, h_{\tilde{M}}, h_{M'}, c^n} \geq 0, \quad \forall J \in \mathcal{P}^{\varphi}.$$

We denote by $\partial^X S$ the boundary of a subset S of a topological space X . For every $i \in I$ Remark 32 and Lemma 36 imply that

$$(66) \quad \begin{aligned} h_{\tilde{M}}(\partial^M K_i) &= -h_{K_i}(\partial^M K_i) \\ &= h_M(i) - \int_{K_i} \omega^{\wedge n}, \end{aligned}$$

where the integral is w.r.t. the orientation $O|K_i$. Let $J \in \mathcal{P}^\varphi$. Recalling the definition (62) of $\tilde{}$ and using (66), we have

$$\sum_{\tilde{i} \in \tilde{J} \cap \tilde{I}} h_{\tilde{M}}(\tilde{i}) = \sum_{i \in J \cap I} \left(h_M(i) - \int_{K_i} \omega^{\wedge n} \right).$$

Combining this with (65) and recalling the definition (19), it follows that

$$\sum_{J, h_M, h_{M'}, c^n} \geq -c^n \sum_{i \in J \cap I} \int_{K_i} \omega^{\wedge n}.$$

Since this holds for every choice of $(K_i)_{i \in I}$, it follows that $\sum_{J, h_M, h_{M'}, c^n} \geq 0$. Hence condition (20) holds with $\mathcal{P} := \mathcal{P}^\varphi$, $f := h_M$, $f' := h_{M'}$, and $C := c^n$. This proves Lemma 49. \square

Remark (helicity inequality). Under the hypotheses of this lemma, the set $M' \setminus \varphi(\text{Int}(M))$ need not be a submanifold of M' , since $\varphi(\partial M)$ may intersect $\partial M'$. This is the reason for the construction of \tilde{M} in the proof of this lemma.

We are now ready for the proof of Theorem 43.

Proof of Theorem 43. Assume that there exist $A_0, (M_a, \omega_a)_{a \in A_0}$ as in the hypothesis of (i). Let $a \in A_0 \cap (0, \infty)$. We define

$$(W_a, \eta_a) := (M_a \sqcup M_{-a}, \omega_a \sqcup \omega_{-a}).$$

Let $A \in \mathcal{P}(A_0 \cap (0, \infty))$. Recall the definition (2) of \mathcal{O}_0 . We define the function

$$c_A := \sup_{a \in A} c_{W_a, \eta_a} : \mathcal{O}_0 \rightarrow [0, \infty].$$

If $k = 2$ and the ball B lies in \mathcal{O} , then we define the function \tilde{c}_A : by

$$(67) \quad \tilde{c}_A := \max \{c_A, w\} : \mathcal{O}_0 \rightarrow [0, \infty].$$

The functions c_A and \tilde{c}_A are generalized capacities on \mathcal{C} .

Claim 1. (i) *The map $\mathcal{P}(A_0 \cap (0, \infty)) \ni A \mapsto c_A \in \text{Cap}(\mathcal{C})$ is injective.*

Assume now that the hypotheses of Theorem 43(ii) are satisfied.

(ii) *The map $\mathcal{P}(A_0 \cap (0, \infty)) \ni A \mapsto \tilde{c}_A \in \text{Cap}(\mathcal{C})$ is injective.*

(iii) For every $A \in \mathcal{P}(A_0 \cap (0, \infty))$ the capacity \tilde{c}_A is normalized.

Proof of Claim 1. We denote

$$h_M := h_{M,O,\omega}, \quad f_a := h_{M_a}, \quad f := (f_a)_{a \in A_0},$$

and define C_0^f, C_1^f as in (21,22). Let $a \neq a' \in A_0 \cap (0, \infty)$, and $c \in (0, \infty)$, such that there exists a \mathcal{C} -morphism φ from $(W_a, c\eta_a)$ to $(W_{a'}, \eta_{a'})$.

Case A: There exist such a φ and $b \in \{a, -a\}, b' \in \{a', -a'\}$, such that $b > b'$ and $\varphi(M_b) \subseteq M_{b'}$. We denote

$$M := M_b, \quad \omega := \omega_b, \quad M' := M_{b'}, \quad \omega' := \omega_{b'}, \quad I := I_M, \quad I' := I_{M'}.$$

Let $d \in A_0$. By hypotheses M_d is nonempty, compact, and 1-connected. Since by hypothesis $n \geq 2 > 0$ and ω_d is maxipotent and exact, we have $\partial M_d \neq \emptyset$. Hence the hypotheses of Lemma 47(i) are satisfied. Applying this lemma, it follows that \mathcal{P}^φ is an (I, I') -partition. By Lemma 49 the set \mathcal{P}^φ is an $(h_M, h_{M'}, c^n)$ -partition. It follows that

$$(68) \quad c^n \leq C_0^f.$$

Consider now the **case** that is complementary to Case A. Then $a < a'$ and there exists a morphism φ from $(W_a, c\eta_a)$ to $(W_{a'}, \eta_{a'})$, such that $\varphi(W_a) \subseteq M_{a'}$. Lemmata 47(ii) and 49 imply that \mathcal{P}^φ is an $(h_{M_a}, h_{M_{-a}}, h_{M_{a'}}, c^n)$ -partition. It follows that $c^n \leq C_1^f$. Combining this with (68), in any case we have

$$c^n \leq C := \max \{C_0^f, C_1^f\}.$$

It follows that

$$\begin{aligned} & \sup \{c \in (0, \infty) \mid \exists a \neq a' \in A_0 \cap (0, \infty) \exists \text{ morphism } (W_a, c\eta_a) \rightarrow (W_{a'}, \eta_{a'})\} \\ & \leq \sqrt[n]{C} \\ & < 1 \quad (\text{using our hypothesis (ib) and Definition 42}). \end{aligned}$$

It follows that

$$(69) \quad c_A(W_{a'}, \eta_{a'}) < 1, \quad \forall A \in \mathcal{P}(A_0 \cap (0, \infty)), \quad a' \in A_0 \cap (0, \infty) \setminus A. \quad 49$$

Let $A \neq A' \in \mathcal{P}(A_0 \cap (0, \infty))$. Assume first that $A' \setminus A \neq \emptyset$. We choose $a' \in A' \setminus A$. Since $c_{A'}(W_{a'}, \eta_{a'}) \geq 1$,⁵⁰ inequality (69) implies that $c_A \neq c_{A'}$. This also holds in the case $A \setminus A' \neq \emptyset$, by an analogous argument. This proves statement (i).

We prove (ii). Combining inequality (69) with our hypothesis (iia), we have

$$\tilde{c}_A(W_{a'}, \eta_{a'}) < 1, \quad \forall A \in \mathcal{P}(A_0 \cap (0, \infty)), \quad a' \in A_0 \cap (0, \infty) \setminus A.$$

Hence an argument as above shows that the map $\mathcal{P}(A_0 \cap (0, \infty)) \ni A \mapsto \tilde{c}_A$ is injective. This proves (ii).

We prove (iii). Let $A \in \mathcal{P}(A_0 \cap (0, \infty))$. By our definition (67) we have

$$(70) \quad \pi = w(B) \leq \tilde{c}_A(B).$$

Since B symplectically embeds into Z , we have $c_{M,\omega}(B) \leq c_{M,\omega}(Z)$ for every object (M, ω) of \mathcal{C} . It follows that

$$(71) \quad \tilde{c}_A(B) \leq \tilde{c}_A(Z).$$

Our hypothesis (iib) and Gromov’s Nonsqueezing Theorem imply that $\tilde{c}_A(Z) \leq \pi$. Combining this with (70,71), it follows that \tilde{c}_A is normalized. This proves (iii) and therefore Claim 1. \square

Claim 1(i) implies that

$$(72) \quad |\mathcal{C}ap(\mathcal{C})| \geq |\mathcal{P}(A_0 \cap (0, \infty))| = \beth_2,$$

where in the second inequality we used our hypothesis that A_0 is an interval of positive length. On the other hand, by Corollary 58 in the appendix the set \mathcal{O}_0 has cardinality at most \beth_1 . It follows that

$$|\mathcal{C}ap(\mathcal{C})| \leq |[0, \infty]^{\mathcal{O}_0}| \leq \beth_1^{\beth_1} = \beth_2.$$

Combining this with (72), the statement of Theorem 43(i) follows.

The statement of Theorem 43(ii) follows from an analogous argument, using parts (ii,iii) of Claim 1. This completes the proof of Theorem 43. \square

⁴⁹A priori the function $c := c_A$ is only defined on the set \mathcal{O}_0 . For a general $(M, \omega) \in \mathcal{O}$ we define $c(M, \omega) := c(M_0, \omega_0)$, where (M_0, ω_0) is an arbitrary object of \mathcal{O}_0 isomorphic to (M, ω) .

⁵⁰In fact equality holds, but we do not use this.

5. Proof of Proposition 44 (sufficient conditions for being an I -collection)

Proof of Proposition 44. Let $I = (I_a), f = (f_a)$ be as in the hypothesis. To simplify notation, we canonically identify the collection f with its disjoint union $\bigsqcup f : \bigsqcup I \rightarrow \mathbb{R}$.

Claim 1. *Let $a, a' \in A_0$. If $a > a'$ then for every partition \mathcal{P} of $I_a \sqcup I_{a'}$ there exists $J \in \mathcal{P}$, such that*

$$(73) \quad \sum_{i \in J \cap I_a} f(i) > \sum_{i' \in J \cap I_{a'}} f(i').$$

Proof of Claim 1. This follows from hypothesis (31). □

By hypothesis (26) there exists k , such that $|I_a| = k + 1$, for every $a \in A_0$. By hypothesis (29) for every $a \in A_0$ the set $f_a^{-1}((0, \infty))$ contains a unique element p_a . Hypotheses (30,27) imply that

$$(74) \quad f(p_a) \leq k + 1, \quad \forall a \in A_0.$$

Recalling the notation (25), we have

$$(75) \quad \inf_{a \in A_0} \sum f_a > 0, \quad (\text{using (32,28)})$$

$$(76) \quad f(p_a) > 1, \quad \forall a \in A_0 \quad (\text{using (75,28)}).$$

Claim 2. *If $k = 1$ or 2 then the inequality (33) holds.*

Proof. For every $a \in A_0$ we have

$$\begin{aligned} f(p_a) &= \sum f_a - \sum_{n \in I_a \setminus \{p_a\}} f(n) \\ &\geq \inf_b \sum f_b + 1 - (k - 1) \sup(\text{im}(f) \cap (-\infty, 0]) \quad (\text{using (28)}) \\ &> k + (2 - k) \inf_b \sum f_b \quad (\text{using (32)}) \\ &\geq k \quad (\text{using that } k = 1 \text{ or } 2, \text{ and (75)}). \end{aligned}$$

Using (74), it follows that (33) holds. This proves Claim 2. □

We now check the conditions (23,24) of Definition 42.

Condition (23): Let $a, a' \in A_0$ be such that $a > a'$, $C \in (0, \infty)$ and \mathcal{P} be an $(f_a, f_{a'}, C)$ -partition. If $C \geq 1$ then Claim 1 implies that condition (20) in Definition 39 with $I := I_a, I' := I_{a'}$ is violated. It follows that $C < 1$.

We denote by J_0 the unique element of \mathcal{P} containing p_a .

Claim 3. *We have $p_{a'} \in J_0$.*

Proof of Claim 3. By Definition 39 we have $|J_0 \cap I_a| = 1$. It follows that $J_0 \cap I_a = \{p_a\}$. Therefore, by condition (20) applied to $J := J_0$, we have

$$Cf(p_a) \leq \sum_{i' \in J_0 \cap I_{a'}} f(i').$$

Since $Cf(p_a) > 0$ and $p_{a'}$ is the only point in $I_{a'}$ at which f is positive, Claim 3 follows. □

Claim 4. *We have $f_{a'}^{-1}(-1) \subseteq J_0$.*

Proof of Claim 4. Let $J \in \mathcal{P} \setminus \{J_0\}$. By (18) the set $J \cap I_a$ consists of a unique element i . Hypothesis (27) and the inequality $C < 1$ imply that $Cf(i) > -1$. Combining this with (20), it follows that

$$(77) \quad \sum_{i' \in J \cap I_{a'}} f(i') > -1.$$

Since J and J_0 are disjoint, Claim 3 implies that $p_{a'} \notin J$. Therefore, (77) implies that $J \cap I_{a'} \cap f^{-1}(-1) = \emptyset$. Since this holds for every $J \in \mathcal{P} \setminus \{J_0\}$, and \mathcal{P} covers $I_{a'}$, it follows that $I_{a'} \cap f^{-1}(-1) \subseteq J_0$. This proves Claim 4. □

Claims 3,4 and hypothesis (28) imply that $|J_0 \cap I_{a'}| \geq 2$. Since $|I_a| = |I_{a'}| = k + 1$ and $p_a \in J_0 \cap I_a$, it follows that

$$(78) \quad |(I_a \sqcup I_{a'}) \setminus J_0| \leq 2k - 1.$$

The condition (18) implies that $|\mathcal{P} \setminus \{J_0\}| = |I_a| - 1 = k$. Since the elements of $\mathcal{P} \setminus \{J_0\}$ are disjoint and their union is contained in $(I_a \sqcup I_{a'}) \setminus J_0$, using (78), it follows that there exists $J_1 \in \mathcal{P} \setminus \{J_0\}$ satisfying $|J_1| \leq 1$. Since $|J_1 \cap I_a| = 1$, it follows that

$$(79) \quad J_1 \cap I_{a'} = \emptyset.$$

The facts $J_1 \neq J_0$, and that p_a lies in J_0 and is the only point of I_a at which f is positive, imply that $\sum_{i \in J_1 \cap I_a} f(i) \leq \sup(\text{im}(f) \cap (-\infty, 0])$. Using (79)

and recalling the definition (19), it follows that

$$(80) \quad \sum_{J_1, f_a, f_{a'}, C} \geq -C \sup(\text{im}(f) \cap (-\infty, 0]).$$

Summing up the inequality (20) over all $J \in \mathcal{P} \setminus \{J_1\}$ and adding (80), we obtain

$$-C \sum f_a + \sum f_{a'} \geq -C \sup(\text{im}(f) \cap (-\infty, 0]).$$

It follows that

$$C \left(-\sup(\text{im}(f) \cap (-\infty, 0]) + \inf_a \sum f_a \right) \leq \sum f_{a'} \leq 1 \quad (\text{using hypothesis (30)}).$$

Combining this with hypothesis (32), it follows that $C_0^f < 1$. Hence f satisfies (23).

Condition (24): Let $a, a' \in (0, \infty)$, such that $a < a'$, $C \in (0, \infty)$ and \mathcal{P} be an $(f_a, f_{-a}, f_{a'}, C)$ -partition. We denote by $J_0 \in \mathcal{P}$ the unique element that contains p_a . We will show that \mathcal{P} and J_0 look like in Figure 5.




Figure 5: The dots in the first row constitute the set I_a , which contains the point p_a , and similarly for I_{-a} and $I_{a'}$. The blue and black sets denote the elements of the partition \mathcal{P} . We show below that except for p_a , the blue set J_0 also contains $p_{-a}, p_{a'}$, and an element of $I_{a'}$ at which f takes on the value -1 . Note that J_0 intersects both I_a and I_{-a} in exactly one point, and that the other elements of \mathcal{P} intersect $I_a \sqcup I_{-a}$ in exactly one point.

Claim 5. *We have $p_{a'}, p_{-a} \in J_0$.*

Proof of Claim 5. We show that $p_{a'} \in J_0$. Conditions (a,b) of Definition 40 with $I^\pm := I_{\pm a}$ imply that $J_0 \cap I_{\pm a}$ is empty or a singleton. Combining this

with the fact that $p_a \in J_0$, hypothesis (27), and (76), we obtain

$$\sum_{i \in J_0 \cap (I_a \sqcup I_{-a})} f(i) > 0.$$

Using condition (20) with $J = J_0$, it follows that $p_{a'} \in J_0$.

To show that $p_{-a} \in J_0$, let $J \in \mathcal{P} \setminus \{J_0\}$. Since $p_{a'} \in J_0$, it does not lie in J . It follows that $\sum_{i' \in J \cap I_{a'}} f(i') \leq 0$. Using (20) with $I = I_a \sqcup I_{-a}$, it follows that

$$(81) \quad \sum_{i \in J \cap (I_a \sqcup I_{-a})} f(i) \leq 0.$$

Conditions (a,b) of Definition 40 with $I^\pm := I_{\pm a}$ imply that $J \cap I_{\pm a}$ is empty or a singleton. Using hypothesis (27) and (81), it follows that $J \cap I_{-a}$ is empty or consists of one element i , satisfying $f(i) \leq 1$. Using (76), it follows that $p_{-a} \notin J$. Since this holds for every $J \in \mathcal{P} \setminus \{J_0\}$, it follows that $p_{-a} \in J_0$. This proves Claim 5. \square

Claim 6. *We have $C < 1$.*

Proof of Claim 6. By Remark 41 we have $|\mathcal{P}| = 2k + 1$. Since $|I_{a'}| = k + 1$, $k \geq 1$, and the elements of \mathcal{P} are disjoint, it follows that there exists $J_1 \in \mathcal{P}$, such that

$$(82) \quad J_1 \cap I_{a'} = \emptyset.$$

Claim 5 implies that $J_1 \neq J_0$, and hence that $p_a, p_{-a} \notin J_1$. By Definition 40(b) we have

$$(83) \quad J_1 \cap I_a \sqcup I_{-a} = \{n\}, \quad \text{for some point } n.$$

By (32) we have

$$(84) \quad f(n) < -1 + \inf_{b \in A_0} \sum f_b.$$

Denoting

$$\sum_J := \sum_{i \in J \cap (I_a \sqcup I_{-a})} f(i), \quad \sum'_J := \sum_{i' \in J \cap I_{a'}} f(i'),$$

we have

$$\begin{aligned}
 1 &\geq \sum f_{a'} \quad (\text{using 30}) \\
 &= \sum_{J \in \mathcal{P}} \sum'_J \\
 &= \sum_{J \in \mathcal{P}} \left(-C \sum_J + \sum'_J \right) + C \sum (f_a + f_{-a}) \\
 &\geq -C \sum_{J_1} + \sum'_{J_1} + 2C \inf_{b \in A_0} \sum f_b \quad (\text{using (20) with } J \in \mathcal{P} \setminus \{J_1\}) \\
 &> C \left(1 + \inf_{b \in A_0} \sum f_b \right) \quad (\text{using (83,84,82)}).
 \end{aligned}$$

Using (75), it follows that $C < 1$. This proves Claim 6. □

Claim 7. *We have $f_{a'}^{-1}(-1) \subseteq J_0$.*

Proof of Claim 7. Let $J \in \mathcal{P} \setminus \{J_0\}$. By Claim 5 we have $p_{-a} \in J_0$. Since also $p_a \in J_0$, by Definition 40(b), it follows that $|J \cap (I_a \sqcup I_{-a})| = 1$. Using hypothesis (27) and (20), it follows that

$$\begin{aligned}
 \sum_{i' \in J \cap I_{a'}} f(i') &\geq -C \\
 (85) \qquad \qquad \qquad &> -1 \quad (\text{by Claim 6}).
 \end{aligned}$$

By Claim 5 we have $p_{a'} \in J_0$. Hence this point does not lie in J . Therefore, (85) implies that $J \cap I_{a'} \cap f^{-1}(-1) = \emptyset$. Since this holds for every $J \in \mathcal{P} \setminus \{J_0\}$, and \mathcal{P} covers $I_{a'}$, it follows that $I_{a'} \cap f^{-1}(-1) \subseteq J_0$. This proves Claim 7. □

Claim 5 and Definition 40(a) imply that $J_0 \cap (I_a \sqcup I_{-a}) = \{p_a, p_{-a}\}$, and therefore,

$$(86) \qquad \sum_{i \in J_0 \cap (I_a \sqcup I_{-a})} f(i) = f(p_a) + f(p_{-a}).$$

Claim 7 and hypothesis (28) imply that

$$\sum_{i' \in J_0 \cap I_{a'}} f(i') \leq f(p_{a'}) - 1.$$

Combining this with (86) and (20) with $J = J_0$, it follows that

$$C(f(p_a) + f(p_{-a})) \leq f(p_{a'}) - 1.$$

It follows that

$$\begin{aligned} C &\leq \frac{f(p_{a'}) - 1}{f(p_a) + f(p_{-a})} \\ &\leq \frac{\sup_b f(p_b) - 1}{2 \inf_b f(p_b)} \\ &< 1 \quad (\text{using (33)}). \end{aligned}$$

Here in the case $k = 1$ or 2 we use Claim 2. It follows that $C_1^f < 1$. Hence f satisfies (24). This completes the proof of Proposition 44. \square

6. Proof of Theorem 17(iii) (cardinality of a generating set)

The proof of Theorem 17(iii) is based on the following lemma. For every set S we denote by $\mathcal{P}(S)$ its power set. For every subcollection $\mathcal{C} \subseteq \mathcal{P}(X)$ we denote by $\sigma(\mathcal{C})$ the σ -algebra generated by \mathcal{C} . It is given by

$$\sigma(\mathcal{C}) := \bigcap_{\mathcal{A} \text{ } \sigma\text{-algebra on } X: \mathcal{C} \subseteq \mathcal{A}} \mathcal{A}.$$

A measurable space is a pair (X, \mathcal{A}) , where X is a set and \mathcal{A} a σ -algebra on X . Let $(X, \mathcal{A}), (X', \mathcal{A}')$ be measurable spaces. A map $f : X \rightarrow X'$ is called $(\mathcal{A}, \mathcal{A}')$ -measurable iff $f^{-1}(A') \in \mathcal{A}$, for all $A' \in \mathcal{A}'$. We denote

$$\mathcal{M}(\mathcal{A}, \mathcal{A}') := \{(\mathcal{A}, \mathcal{A}')\text{-measurable map: } X \rightarrow X'\}.$$

Lemma 51 (cardinality of the set of measurable maps). *Let X, X' be sets and $\mathcal{C} \subseteq \mathcal{P}(X), \mathcal{C}' \subseteq \mathcal{P}(X')$ be subcollections. Assume that $|\mathcal{C}| \leq \beth_1, |\mathcal{C}'| \leq \beth_0 = \aleph_0$, and*

$$(87) \quad \forall x' \in X' : \bigcap_{C' \in \mathcal{C}': x' \in C'} C' = \{x'\}.$$

We define $\mathcal{A} := \sigma(\mathcal{C}), \mathcal{A}' := \sigma(\mathcal{C}')$. Then $\mathcal{M}(\mathcal{A}, \mathcal{A}')$ has cardinality at most \beth_1 .

For the proof of this lemma we need the following.

Lemma 52 (cardinality of σ -algebra). *Let X be a set and $\mathcal{C} \subseteq \mathcal{P}(X)$ be a subcollection of cardinality at most \aleph_1 . Then $\sigma(\mathcal{C})$ has cardinality at most \aleph_1 .*

The proof of this lemma is based on the following. Let S be a set, $F : \mathcal{P}(S) \rightarrow \mathcal{P}(S)$, such that

$$(88) \quad A \subseteq F(A), \quad \forall A \in \mathcal{P}(S).$$

Let $A \in \mathcal{P}(S)$. We define $\langle F, A \rangle$, the *set generated by F, A* , to be the smallest fixed point of F containing A . This is the set given by

$$\langle F, A \rangle = \bigcap \{B \in \mathcal{P}(S) \mid A \subseteq B = F(B)\}.$$
⁵¹

Lemma 53 (cardinality of generated set). *The set $\langle F, A \rangle$ has cardinality at most \aleph_1 , if the following conditions are satisfied:*

- (a) *F is monotone, i.e., $B \subseteq C$ implies that $F(B) \subseteq F(C)$.*
- (b) *$|A| \leq \aleph_1$.*
- (c) *If $|B| \leq \aleph_1$ then $|F(B)| \leq \aleph_1$, for every $B \in \mathcal{P}(S)$.*
- (d) *If $B \in \mathcal{P}(S)$ satisfies*

$$(89) \quad F(C) \subseteq B, \quad \forall \text{ countable subset } C \subseteq B,$$

then B is a fixed point of F .

Proof of Lemma 53. We denote by ω_1 the smallest uncountable (von Neumann) ordinal, i.e., the set of countable ordinals. We define $A_0 := A$, and using transfinite recursion, for every $\alpha \leq \omega_1$, we define

$$(90) \quad A_\alpha := \begin{cases} F(A_\beta), & \text{if } \alpha = \beta + 1, \\ \bigcup_{\beta < \alpha} A_\beta, & \text{if } \alpha \neq 0 \text{ is a limit ordinal.} \end{cases}$$

(A limit ordinal is an ordinal for which there does not exist any ordinal β for which $\alpha = \beta + 1$.)

⁵¹This intersection is well-defined, since the collection of all admissible B is nonempty. It contains $B = S$.

Claim 1. *We have*

$$\langle F, A \rangle \subseteq A_{\omega_1}.$$

Proof of Claim 1. Since $A_0 \subseteq A_{\omega_1}$, it suffices to show that A_{ω_1} is a fixed point of F .

Claim 2. *Condition (89) is satisfied with $B = A_{\omega_1}$.*

Proof of Claim 2. Let $C \subseteq A_{\omega_1}$ be a countable subset. The definition (90), condition (88), and transfinite induction imply that for every pair α, β of ordinals, we have

$$(91) \quad \alpha \leq \beta \Rightarrow A_\alpha \subseteq A_\beta.$$

We choose a collection $(\alpha_c)_{c \in C}$ of countable ordinals, such that $c \in A_{\alpha_c}$, for every $c \in C$. The ordinal

$$\alpha := \sup_{c \in C} \alpha_c := \bigcup_{c \in C} \alpha_c$$

is countable, and therefore less than ω_1 . For every $c \in C$, we have $\alpha_c \leq \alpha$, and thus by (91), $A_{\alpha_c} \subseteq A_\alpha$. It follows that $C \subseteq A_\alpha$, and therefore,

$$\begin{aligned} F(C) &\subseteq F(A_\alpha) && \text{(using (a))} \\ &= A_{\alpha+1} && \text{(using (90))} \\ &\subseteq A_{\omega_1} && \text{(using } \alpha + 1 < \omega_1 \text{ and (91)).} \end{aligned}$$

This proves Claim 2. □

By this claim and (d) the set A_{ω_1} is a fixed point of F . This proves Claim 1. □

For every ordinal α we denote by $P(\alpha)$ the statement “ $|A_\alpha| \leq \beth_1$ ”.

Claim 3. *The statement $P(\alpha)$ is true for all $\alpha \leq \omega_1$.*

Proof of Claim 3. We prove this by transfinite induction. Let $\alpha \leq \omega_1$ and assume that the statement holds for all $\beta < \alpha$. If $\alpha = 0$ then $P(0)$ holds by our hypothesis (b). If $\alpha = \beta + 1$ for some β then $P(\alpha)$ holds by (90) and our hypothesis (c). If $\alpha \neq 0$ is a limit ordinal, then $P(\alpha)$ holds by (90), our induction hypothesis, and the fact $|\alpha| \leq |\omega_1| \leq \beth_1$. This completes the inductive step. Claim 3 now follows from transfinite induction. □

Lemma 53 follows from Claims 1 and 3. \square

Proof of Lemma 52. This follows from Lemma 53 with

$$S := \mathcal{P}(X), A := \mathcal{C}, F(\mathcal{D}) := \left\{ \bigcup \mathcal{E} \mid \mathcal{E} \subseteq \mathcal{D} \text{ countable} \right\} \cup \{X \setminus E \mid E \in \mathcal{D}\}.$$

To see that (d) holds, let $B = \mathcal{D} \in \mathcal{P}(S)$ be such that (89) holds. It suffices to show that \mathcal{D} is closed under countable unions and complements. Let $\mathcal{E} \subseteq \mathcal{D}$ be a countable subcollection. We have

$$\begin{aligned} \bigcup \mathcal{E} &\in F(\mathcal{E}) \\ &\subseteq \mathcal{D} \quad (\text{using (89)}). \end{aligned}$$

Hence \mathcal{D} is closed under countable unions. Let now $E \in \mathcal{D}$. We have

$$\begin{aligned} X \setminus E &\in F(\{E\}) \\ &\subseteq \mathcal{D} \quad (\text{using (89)}). \end{aligned}$$

Hence \mathcal{D} is closed under complements. It follows that \mathcal{D} is a fixed point of F . This proves (d) and completes the proof of Lemma 52. \square

Proof of Lemma 51. Recall that for every pair of sets S, S' we denote by S'^S the set of maps from S to S' . Let $f \in \mathcal{M}(\mathcal{A}, \mathcal{A}')$ and $x' \in X'$. Our hypothesis that $|\mathcal{C}'| \leq \aleph_0$ and (87) imply that the set $\{x'\}$ is a countable intersection of elements of \mathcal{C}' . Hence it lies in \mathcal{A}' . It follows that $f^{-1}(x') \in \mathcal{A}$. The following map is therefore well-defined:

$$\iota : \mathcal{M}(\mathcal{A}, \mathcal{A}') \rightarrow \mathcal{A}^{X'}, \quad \iota(f)(x') := f^{-1}(x').$$

We define the map

$$\varphi : \mathcal{M}(\mathcal{A}, \mathcal{A}') \rightarrow \mathcal{A}^{\mathcal{C}'}, \quad \varphi(f)(C') := f^{-1}(C'),$$

$$\psi : \mathcal{A}^{\mathcal{C}'} \rightarrow \mathcal{A}^{X'}, \quad \psi(A)(x') := \bigcap_{C' \in \mathcal{C}': x' \in C'} A(C').$$

Our hypothesis $|\mathcal{C}'| \leq \aleph_0$ implies that $\psi(A)(x')$ is a countable intersection of elements of \mathcal{A} , hence an element of \mathcal{A} . It follows that ψ is well-defined. For

every $f \in \mathcal{M}(\mathcal{A}, \mathcal{A}')$ and $x' \in X'$, we have

$$\begin{aligned} \iota(f)(x') &= f^{-1}(x') \\ &= f^{-1}\left(\bigcap_{C' \in \mathcal{C}': x' \in C'} C'\right) \quad (\text{by (87)}) \\ &= \bigcap_{C' \in \mathcal{C}': x' \in C'} f^{-1}(C') \\ &= (\psi(\varphi(f)))(x'). \end{aligned}$$

Hence the equality $\iota = \psi \circ \varphi$ holds. Since ι is injective, it follows that φ is injective. Our hypothesis that $|\mathcal{C}| \leq \beth_1$ and Lemma 52 imply that $|\mathcal{A} = \sigma(\mathcal{C})| \leq \beth_1$. Since $|\mathcal{C}'| \leq \aleph_0$, it follows that $|\mathcal{A}^{C'}| \leq \beth_1$. Since φ maps $\mathcal{M}(\mathcal{A}, \mathcal{A}')$ to $\mathcal{A}^{C'}$, it follows that $|\mathcal{M}(\mathcal{A}, \mathcal{A}')| \leq \beth_1$. This proves Lemma 51. \square

In the proof of Theorem 17(iii) we will also use the following.

Remarks 54.

- (i) Every countable product of second countable topological spaces is second countable.
- (ii) Let (X, τ) be a topological space and \mathcal{B} a basis of τ . Then the following inequality holds:

$$|\tau| \leq 2^{|\mathcal{B}|}$$

Proof of Theorem 17(iii). Let \mathcal{G}_0 be a countable subset of X^S . We equip $X^{\mathcal{G}_0}$ with the product topology $\tau_{\mathcal{G}_0}$. We define $\mathcal{A}_{\mathcal{G}_0}, \mathcal{A}$ to be the Borel σ -algebras of $\tau_{\mathcal{G}_0}, \tau$.

Claim 1. *The set $\mathcal{M}(\mathcal{A}_{\mathcal{G}_0}, \mathcal{A})$ has cardinality at most \beth_1 .*

Proof of Claim 1. Our assumption that τ is separable and metrizable, implies that it is second countable. Hence by Remark 54(i), the same holds for $\tau_{\mathcal{G}_0}$. Hence by Remark 54(ii), we have

$$(92) \quad |\tau_{\mathcal{G}_0}| \leq 2^{\aleph_0} = \beth_1.$$

We have $\mathcal{A}_{\mathcal{G}_0} = \sigma(\tau_{\mathcal{G}_0})$. Since τ is separable, there exists a countable τ -dense subset A of X . We define \mathcal{C} to be the collection of all open balls with rational radius around points in A . Since A is dense, every element of τ is a union of elements of \mathcal{C} . Since A is countable, the set \mathcal{C} is countable. It follows that

$\mathcal{A} = \sigma(\tau) = \sigma(\mathcal{C})$. Since τ is separable and metrizable, the condition (87) with \mathcal{C}' replaced by \mathcal{C} is satisfied. Using (92), it follows that the hypotheses of Lemma 51 are satisfied with $\mathcal{C}, \mathcal{C}'$ replaced by $\tau_{\mathcal{G}_0}, \mathcal{C}$. Applying this lemma, it follows that $|\mathcal{M}(\mathcal{A}_{\mathcal{G}_0}, \mathcal{A})| \leq \beth_1$. This proves Claim 1. \square

Let \mathcal{G} be a subset of X^S of cardinality at most \beth_1 . By Definition 16 the set countably Borel-generated by \mathcal{G} is given by

$$\langle \mathcal{G} \rangle := \{f \circ \text{ev}_{\mathcal{G}_0} \mid \mathcal{G}_0 \subseteq \mathcal{G} \text{ countable, } f \in \mathcal{M}(\mathcal{A}_{\mathcal{G}_0}, \mathcal{A})\}.$$

The set of all countable subsets of \mathcal{G} has cardinality at most $\beth_1^{\aleph_0} = \beth_1$. Using Claim 1, it follows that

$$|\langle \mathcal{G} \rangle| \leq \beth_1^2 = \beth_1.$$

This proves Theorem 17(iii). \square

7. Proof of Theorem 25 (uncountability of every generating set under a mild hypothesis)

Proof of Theorem 25. Let $\mathcal{C} = (\mathcal{O}, \mathcal{M}), A, M$ be as in the hypothesis. W.l.o.g. we may assume that A is open. Our hypothesis (9) implies that the function $\text{Vol}^{\frac{1}{n}} \circ M : A \rightarrow \mathbb{R}$ is continuous and strictly increasing. Hence it is injective with image \tilde{A} given by an interval. We define

$$\tilde{M} := M \circ \left(\text{Vol}^{\frac{1}{n}} \circ M\right)^{-1} : \tilde{A} \rightarrow \mathcal{O}.$$

Let $\tilde{a}_0 \in \tilde{A}$. We define

$$g_{\tilde{a}_0} := c_{\tilde{M}_{\tilde{a}_0}} \circ \tilde{M} : \tilde{A} \rightarrow \mathbb{R}.$$

Claim 1. *This function is not differentiable at \tilde{a}_0 .*

Proof of Claim 1. We have

$$\text{Vol}^{\frac{1}{n}} \circ \tilde{M} = \text{id}.$$

It follows that

$$(93) \quad g_{\tilde{a}_0}(\tilde{a}) \leq \frac{\tilde{a}}{\tilde{a}_0}, \quad \forall \tilde{a} \in \tilde{A} \cap (0, \tilde{a}_0).$$

Our hypothesis (10) implies that

$$g_{\tilde{a}_0}(\tilde{a}) = 1, \quad \forall \tilde{a} \in \tilde{A} \cap [\tilde{a}_0, \infty).$$

Combining this with (93), it follows that $g_{\tilde{a}_0}$ is not differentiable at \tilde{a}_0 . This proves Claim 1. □

Let now \mathcal{G} be a countable subset of $Cap(\mathcal{C})$. Let $c \in \mathcal{G}$. The inequality “ \geq ” in our hypothesis (10) implies that the function $c \circ M$ is increasing. It follows that the same holds for $c \circ \tilde{M}$. Therefore, by Lebesgue’s Monotone Differentiation Theorem the function $c \circ \tilde{M}$ is differentiable⁵² almost everywhere, see e.g. [16, p. 156, Theorem 1.6.25]. Since \mathcal{G} is countable, it follows that the set of all points in \tilde{A} at which the function $c \circ \tilde{M}$ is differentiable, for every $c \in \mathcal{G}$, has full Lebesgue measure. Since A has positive length, the same holds for \tilde{A} . It follows that there exists a point $\tilde{a}_0 \in \tilde{A}$ at which $c \circ \tilde{M}$ is differentiable, for every $c \in \mathcal{G}$.

Let \mathcal{G}_0 be a finite subset of \mathcal{G} , and $f : [0, \infty]^{\mathcal{G}_0} \rightarrow [0, \infty]$ a differentiable function. We define $ev_{\mathcal{G}_0}$ as in (4). Since $c \circ \tilde{M}$ is differentiable at \tilde{a}_0 for every $c \in \mathcal{G}_0$, the same holds for the map $ev_{\mathcal{G}_0} \circ \tilde{M} : \tilde{A} \rightarrow [0, \infty]^{\mathcal{G}_0}$. It follows that the composition $f \circ ev_{\mathcal{G}_0} \circ \tilde{M}$ is differentiable at \tilde{a}_0 . Using Claim 1, it follows that

$$f \circ ev_{\mathcal{G}_0} \circ \tilde{M} \neq g_{\tilde{a}_0} = c_{\tilde{M}_{\tilde{a}_0}} \circ \tilde{M},$$

and therefore that $f \circ ev_{\mathcal{G}_0} \neq c_{\tilde{M}_{\tilde{a}_0}}$. Hence \mathcal{G}_0 does not finitely differentially generate $c_{\tilde{M}_{\tilde{a}_0}}$. This proves Theorem 25. □

Appendix A. Cardinality of the set of equivalence classes of pairs of manifolds and forms

In this section we prove that the set of diffeomorphism types of smooth manifolds has cardinality at most \aleph_1 . We also prove that the same holds for the set of all equivalence classes of pairs (M, ω) , where M is a manifold, and ω is a differential form on M . We used this in the proof of Theorem 43, to estimate the cardinality of the set of (normalized) capacities from above.

In order to deal with a certain set-theoretic issue, we explain how to make the class of all diffeomorphism types a set. Let A, B be sets and $S : A \rightarrow B$ a map. Let $a \in A$. We denote $S_a := S(a)$. Recall that in ZFC “everything”

⁵²in the usual sense

is a set, in particular S_a . Recall also that the disjoint union of S is defined to be

$$\bigsqcup S := \{(a, s) \mid s \in S_a\}.$$

We denote

$$H^n := \{x \in \mathbb{R}^n \mid x_n \geq 0\}.$$

Let S be a set. By an atlas on S we mean a subset

$$\mathcal{A} \subseteq \bigsqcup_{U \in \mathcal{P}(S)} (H^n)^U,$$

such that

$$\bigcup_{(U, \varphi) \in \mathcal{A}} U = S,$$

for every $(U, \varphi) \in \mathcal{A}$ the map φ is injective, and for all $(U, \varphi), (U', \varphi') \in \mathcal{A}$ the set $\varphi(U \cap U')$ is open (in H^n) and the transition map

$$\varphi' \circ \varphi^{-1} : \varphi(U \cap U') \rightarrow H^n$$

is smooth. We call an atlas maximal iff it is not contained in any strictly larger atlas. By a *(smooth finite-dimensional real) manifold (with boundary)* we mean a pair $M = (S, \mathcal{A})$, where S is a set and \mathcal{A} is a maximal atlas on S , such that the induced topology is Hausdorff and second countable. We denote by \beth_1 the (von Neumann) cardinal $2^{\beth_0} = \aleph_1$, and by \sim the diffeomorphism relation on

$$(A.1) \quad \mathcal{M}_0 := \{(S, \mathcal{A}) \mid S \subseteq \beth_1, (S, \mathcal{A}) \text{ is a manifold}\}.$$

This means that $M \sim M'$ iff M and M' are diffeomorphic. We define the *set of diffeomorphism types (of manifolds)* to be

$$\mathfrak{M} := \{ \sim\text{-equivalence class}\}.$$

Remarks 55 (diffeomorphism types).

- The above definition overcomes the set-theoretic issue that the “set” of diffeomorphism classes of all manifolds (without any restriction on the underlying set) is not a set (in ZFC).
- Every manifold M is diffeomorphic to one whose underlying set is a subset of \beth_1 . To see this, note that using second countability and

the axiom of choice, the set underlying M has cardinality $\leq \beth_1$. This means that there exists an injective map $f : M \rightarrow \beth_1$. Pushing forward the manifold structure via f , we obtain a manifold whose underlying set is a subset of \beth_1 , as claimed.

- By the last remark, heuristically, there is a canonical bijection between \mathfrak{M} and the “set” of diffeomorphism classes of all manifolds.
- One may understand \mathfrak{M} in a more general way as follows. Let \mathcal{M} be a set consisting of manifolds, such that every manifold is diffeomorphic to some element of \mathcal{M} . For example, let S be a set of cardinality at least \beth_1 and define \mathcal{M} to be the set of all manifolds whose underlying set is a subset of S . The set \mathfrak{M} is in bijection with the set of all diffeomorphism classes of elements of \mathcal{M} .

Proposition 56. *The set \mathfrak{M} has cardinality at most \beth_1 .*

In the proof of this result we will use the following.

Remark 57 (Whitney’s Embedding Theorem). Let $n \in \mathbb{N}_0$ and M be a (smooth) manifold of dimension n . There exists a (smooth) embedding of M into \mathbb{R}^{2n+1} with closed image. To see this, consider the double \widetilde{M} of M , which is obtained by gluing two copies of M along the boundary. By Whitney’s Embedding Theorem there exists an embedding of \widetilde{M} into \mathbb{R}^{2n+1} with closed image, see e.g. [7, 2.14. Theorem, p. 55]⁵³. Composing such an embedding with one of the two canonical inclusions of M in \widetilde{M} , we obtain an embedding of M into \mathbb{R}^{2n+1} with closed image, as desired.

Proof of Proposition 56. We define

$$\mathcal{M} := \bigsqcup_{m \in \mathbb{N}_0} \{\text{submanifold of } \mathbb{R}^m\}.$$

Claim 1. *We have $|\mathcal{M}| \leq \beth_1$.*

⁵³In this section of Hirsch’s book \widetilde{M} manifolds are not allowed to have boundary. This is the reason for considering \widetilde{M} , rather than M .

Proof. Let $n, m \in \mathbb{N}_0$. The topological space $\mathbb{N}_0 \times H^n$ is separable. Since $|\mathbb{R}^m| \leq \beth_1$, it follows that

$$(A.2) \quad |C(\mathbb{N}_0 \times H^n, \mathbb{R}^m)| \leq \beth_1.$$

Let $n \in \mathbb{N}_0$ and $(m, M) \in \mathcal{M}$, such that M is of dimension n . Since M is second countable, there exists a surjective map $\psi : \mathbb{N}_0 \times H^n \rightarrow M$ whose restriction to $\{i\} \times H^n$ is an embedding, for every $i \in \mathbb{N}_0$. It follows that M lies in the image of the map

$$C(\mathbb{N}_0 \times H^n, \mathbb{R}^m) \rightarrow \mathcal{P}(\mathbb{R}^m), \quad f \mapsto \text{im}(f).$$

Combining this with (A.2), it follows that $|\mathcal{M}| \leq \beth_1$. This proves Claim 1. □

Let $n \in \mathbb{N}_0$. We choose an injection $\alpha : \mathbb{R}^{2n+1} \rightarrow \beth_1$, and consider the push-forward map

$$\alpha_* : \mathcal{M} \rightarrow \mathfrak{M}, \quad \alpha_*(S, \mathcal{A}) := [\alpha(S), \alpha_*\mathcal{A}].$$

Remark 57 implies that this map is surjective. Using Claim 1, it follows that $|\mathfrak{M}| \leq \beth_1$. This proves Proposition 56. □

We define \mathcal{M}_0 as in (A.1),

$$\begin{aligned} \Omega(M) &:= \{\text{differential form on } M\}, \\ \Omega_0 &:= \bigsqcup_{M \in \mathcal{M}_0} \Omega(M), \end{aligned}$$

the equivalence relation \approx on Ω_0 by

$$(M, \omega) \approx (M', \omega) : \iff \exists \text{ diffeomorphism } \varphi : M \rightarrow M' : \varphi^*\omega' = \omega,$$

$$\text{and } \bar{\Omega} := \Omega_0 / \approx .$$

Remark. Philosophically, this is the “set” of all equivalence classes of pairs (M, ω) , where M is an arbitrary manifold and ω is a differential form on M . The above definition makes this idea precise.

Corollary 58. *The set $\bar{\Omega}$ has cardinality at most \beth_1 .*

Proof of Corollary 58. If M, M' are manifolds and $\varphi : M \rightarrow M'$ is a diffeomorphism then

$$(A.3) \quad \varphi^* : \Omega(M') \rightarrow \Omega(M) \text{ is a bijection.}$$

We denote by $\Pi : \Omega_0 \rightarrow \overline{\Omega}$ and $\pi : \mathcal{M}_0 \rightarrow \mathfrak{M}$ the canonical projections, and by $f : \Omega_0 \rightarrow \mathcal{M}_0, f((M, \omega)) := M$, the forgetful map. We define $F : \overline{\Omega} \rightarrow \mathfrak{M}$ to be the unique map satisfying $F \circ \Pi = \pi \circ f$. Let $\mathcal{M} \in \mathfrak{M}$. Choosing $M \in \mathcal{M}$, we have

$$(A.4) \quad \begin{aligned} F^{-1}(\mathcal{M}) &= \Pi((F \circ \Pi)^{-1}(\mathcal{M})) \\ &= \Pi((\pi \circ f)^{-1}(\mathcal{M})) \\ &= \Pi(f^{-1}(\mathcal{M})) \quad (\text{using that } \pi^{-1}(\mathcal{M}) = \mathcal{M}) \\ &= \Pi(f^{-1}(M) = \Omega(M)) \quad (\text{using (A.3)}). \end{aligned}$$

Since M is separable and $|TM| \leq \beth_1$, we have $|C(M, TM)| \leq \beth_1$. Using $\Omega(M) \subseteq C(M, TM)$, (A.4), and Proposition 56, it follows that

$$\left| \overline{\Omega} = \bigcup_{\mathcal{M} \in \mathfrak{M}} F^{-1}(\mathcal{M}) \right| \leq \beth_1^2 = \beth_1.$$

This proves Corollary 58. □

Remark. Let $n \geq 2$. Then the set of diffeomorphism types of manifolds of dimension n has cardinality equal to \beth_1 . To see this, we choose a countable set \mathcal{M} of nondiffeomorphic connected n -manifolds. The map

$$\{0, 1\}^{\mathcal{M}} \ni u \mapsto \bigsqcup_{M \in \mathcal{M}: u(M)=1} M \in \{n\text{-manifold}\}$$

is injective. Hence the set of diffeomorphism types of manifolds of dimension n has cardinality $\geq \beth_1$. Combining this with Proposition 56, it follows that this cardinality equals \beth_1 , as claimed.

Appendix B. Proof of Theorem 29 (monotone generation for ellipsoids)

Theorem 29 follows from McDuff’s characterization of the existence of symplectic embeddings between ellipsoids, and the fact that monotone generation is equivalent to almost order-reflexion. To explain this, let (S, \leq) be

a preordered set. We fix an order-preserving $(0, \infty)$ -action on S . We define the *order-capacity function* $c^{\leq} : S \times S \rightarrow [0, \infty]$ by

$$c^{\leq}(s, s') := \sup \{a \in (0, \infty) \mid as \leq s'\}.$$

Remark 59. For every $s \in S$ the function $c^{\leq}(s, \cdot)$ is a capacity, as defined in (14).

Let $\mathcal{G} \subseteq \text{Cap}(S)$. We call \mathcal{G} *almost order-reflecting* iff for all $s, s' \in S$ the following holds:

$$c(s) \leq c(s'), \forall c \in \mathcal{G} \Rightarrow c^{\leq}(s, s') \geq 1.$$

Remark. A map f between two preordered sets is called *order-reflecting* if $f(s) \leq f(s')$ implies that $s \leq s'$. The set \mathcal{G} is almost order-reflecting iff its evaluation map is “almost” order-reflecting, in the sense that $\text{ev}_{\mathcal{G}}(s) \leq \text{ev}_{\mathcal{G}}(s')$ implies that for every $a_0 \in (0, 1)$ there exists an $a \in [a_0, \infty)$, such that $as \leq s'$.

Proposition 60 (characterization of monotone generation). *The set \mathcal{G} monotonely generates if and only if it is almost order-reflecting.*

In the proof of this result we use the following. Let $(X, \leq), (X', \leq')$ be preordered sets, $X_0 \subseteq X$, and $f : X_0 \rightarrow X'$. We define the *monotonization of f* to be the map $F : X \rightarrow X'$ given by

$$F(x) := \sup \{f(x_0) \mid x_0 \in X_0 : x_0 \leq x\}.$$

Remarks 61 (monotonization).

- (i) The map F is monotone.
- (ii) If X and X' are equipped with order-preserving $(0, \infty)$ -actions and f is homogeneous, then its *monotonization* is homogeneous.
- (iii) If f is monotone then it agrees with the restriction of F to X_0 .

Proof of Proposition 60. “ \Rightarrow ”: Assume that \mathcal{G} monotonely generates. Let $s, s' \in S$ be such that $c(s) \leq c(s')$, for every $c \in \mathcal{G}$. This means that

$$(B.5) \quad \text{ev}_{\mathcal{G}}(s) \leq \text{ev}_{\mathcal{G}}(s').$$

By Remark 59 and our assumption there exists a monotone function $F : [0, \infty]^{\mathcal{G}} \rightarrow [0, \infty]$, such that

$$c_s := c^{\leq}(s, \cdot) = F \circ \text{ev}_{\mathcal{G}}.$$

We have

$$\begin{aligned} 1 &\leq c_s(s) && \text{(since } \leq \text{ is reflexive and hence } s \leq s) \\ &= F \circ \text{ev}_{\mathcal{G}}(s) \\ &\leq F \circ \text{ev}_{\mathcal{G}}(s') && \text{(using (B.5) and monotonicity of } F) \\ &= c_s(s'). \end{aligned}$$

Hence \mathcal{G} is almost order-reflecting. This proves “ \Rightarrow ”.

To prove the implication “ \Leftarrow ”, assume that \mathcal{G} is almost order-reflecting. Let $c_0 \in \text{Cap}(S)$.

Claim 1. *For every pair of points $s, s' \in S$, satisfying $\text{ev}_{\mathcal{G}}(s) \leq \text{ev}_{\mathcal{G}}(s')$, we have $c_0(s) \leq c_0(s')$.*

Proof. Since $c(s) \leq c(s')$, for every $c \in \mathcal{G}$, by assumption, we have $c_s(s') \geq 1$. Let $a_0 \in (0, 1)$. It follows that there exists $a \in [a_0, \infty)$, such that $as \leq s'$. It follows that

$$a_0 c_0(s) \leq a c_0(s) = c_0(as) \leq c_0(s').$$

Since this holds for every $a_0 \in (0, 1)$, it follows that $c_0(s) \leq c_0(s')$. This proves Claim 1. □

We define $f : \text{im}(\text{ev}_{\mathcal{G}}) \rightarrow [0, \infty]$ by setting $f(x) := c_0(s)$, where s is an arbitrary point in $\text{ev}_{\mathcal{G}}^{-1}(x) \subseteq S$. By Claim 1 this function is well-defined, i.e., it does not depend on the choice of s . It satisfies

$$(B.6) \quad f \circ \text{ev}_{\mathcal{G}} = c_0.$$

It follows from this equality and Claim 1 that f is monotone. By Remark 61(i,iii) and equality (B.6) the monotone function F of f is a monotone function on $[0, \infty]^{\mathcal{G}}$ that satisfies $F \circ \text{ev}_{\mathcal{G}} = c_0$. This proves “ \Leftarrow ” and completes the proof of Proposition 60. □

Proof of Theorem 29. We equip the set of ellipsoids in (V, ω) with the pre-order $E \leq E'$ iff there exists a symplectic embedding of E into E' . By Theorem 1.1 in D. McDuff’s article [11] the condition $c_j^V(E) \leq c_j^V(E')$, for all

$j \in \mathbb{N}_0$, implies that E symplectically embeds into E' . It follows that the set of all c_j^V (with $j \in \mathbb{N}_0$) is almost order-reflecting. Hence by Proposition 60 this set monotonely generates. This proves Theorem 29. \square

References

- [1] V. I. Arnold and B. A. Khesin, *Topological Methods in Hydrodynamics*, Springer-Verlag, Berlin, Heidelberg (1998), ISBN 0-387-94947-x.
- [2] M. Brown, *Locally flat embeddings of topological manifolds*, in *Topology of 3-manifolds and related topics* (Proc. The Univ. of Georgia Institute, 1961), 83–91, Prentice-Hall, Englewood Cliffs, N.J. (1962).
- [3] K. Cieliebak, H. Hofer, J. Latschev, and F. Schlenk, *Quantitative symplectic geometry*, in *Dynamics, ergodic theory, and geometry*, Vol. 54 of *Math. Sci. Res. Inst. Publ.*, 1–44, Cambridge Univ. Press, Cambridge (2007).
- [4] R. Connelly, *A new proof of Brown's collaring theorem*, Proc. Amer. Math. Soc. **27** (1971) 180–182.
- [5] I. Ekeland and H. Hofer, *Symplectic Topology and Hamiltonian Dynamics.*, Mathematische Zeitschrift **200** (1988/89), no. 3, 355–378.
- [6] ———, *Symplectic Topology and Hamiltonian Dynamics II.*, Mathematische Zeitschrift **203** (1990), no. 4, 553–568.
- [7] M. W. Hirsch, *Differential topology*, Vol. 33 of *Graduate Texts in Mathematics*, Springer-Verlag, New York (1994), ISBN 0-387-90148-5. Corrected reprint of the 1976 original.
- [8] M. Hutchings, *Quantitative embedded contact homology*, J. Differential Geom. **88** (2011), no. 2, 231–266.
- [9] A. S. Kechris, *Classical descriptive set theory*, Vol. 156 of *Graduate Texts in Mathematics*, Springer-Verlag, New York (1995), ISBN 0-387-94374-9.
- [10] D. McDuff, *Symplectic embeddings of 4-dimensional ellipsoids*, J. Topol. **2** (2009), no. 1, 1–22.
- [11] ———, *The Hofer conjecture on embedding symplectic ellipsoids*, J. Differential Geom. **88** (2011), no. 3, 519–532.

- [12] D. McDuff and D. Salamon, *Introduction to symplectic topology*, Oxford Mathematical Monographs, The Clarendon Press, Oxford University Press, New York, second edition (1998), ISBN 0-19-850451-9.
- [13] F. Schlenk, *Embedding problems in symplectic geometry*, Vol. 40 of *De Gruyter Expositions in Mathematics*, Walter de Gruyter GmbH & Co. KG, Berlin (2005), ISBN 3-11-017876-1.
- [14] ———, *Symplectic embedding problems, old and new*, *Bull. Amer. Math. Soc. (N.S.)* **55** (2018), no. 2, 139–182.
- [15] J. Swoboda and F. Ziltener, *Coisotropic displacement and small subsets of a symplectic manifold*, *Mathematische Zeitschrift* **271** (2012), no. 1, 415–445.
- [16] T. Tao, *An Introduction to Measure Theory* Vol. 126, Amer Mathematical Society (2011).
- [17] K. Zehmisch and F. Ziltener, *Discontinuous symplectic capacities*, *J. Fixed Point Theory Appl.* **14** (2013), no. 1, 299–307.

LABORATOIRE DE MATHÉMATIQUES D'ORSAY, UNIVERSITÉ PARIS-SACLAY
BÂTIMENT 307, RUE MICHEL MAGAT, 91400 ORSAY, FRANCE
E-mail address: dusanjoksimovic123@gmail.com

MATHEMATICAL INSTITUTE, UTRECHT UNIVERSITY
BUDAPESTLAAN 6, 3584 CD UTRECHT, THE NETHERLANDS
E-mail address: f.ziltener@uu.nl

RECEIVED AUGUST 20, 2021

ACCEPTED NOVEMBER 13, 2021