

MATHEMATICAL RESEARCH LETTERS

CONTENTS

Volume 20, Number 5

September 2013

Jeffrey D. Achter , <i>On the abelian fivefolds attached to cubic surfaces</i>	805
Andre Chatzistamatiou and Sinan Ünver , <i>On p-adic periods for mixed Tate motives over a number field</i>	825
Bing-Long Chen, Guoyi Xu and Zhuhong Zhang , <i>Local pinching estimates in 3-dim Ricci flow</i>	845
M. Colțoiu, K. Diederich and C. Joița , <i>On complex spaces with prescribed singularities</i>	857
Anand Deopurkar and Anand Patel , <i>Sharp slope bounds for sweeping families of trigonal curves</i>	869
Tara S. Holm and Ana Rita Pires , <i>The topology of toric origami manifolds</i>	885
A.S. Iakovlev and J.-O. Strömberg , <i>Lower bounds for the weak type $(1,1)$ estimate for the maximal function associated to cubes in high dimensions</i> .	907
Rodney Keaton , <i>Level stripping for degree 2 Siegel modular forms</i>	919
Dong Li , <i>On a frequency localized Bernstein inequality and some generalized Poincaré-type inequalities</i>	933
Zhong-Hua Li , <i>Another proof of Zagier's evaluation formula of the multiple zeta values $\zeta(2, \dots, 2, 3, 2, \dots, 2)$</i>	947
Daisuke Matsushita and De-Qi Zhang , <i>Zariski F-decomposition and Lagrangian fibration on hyperkähler manifolds</i>	951
Jennifer Park , <i>A universal first-order formula defining the ring of integers in a number field</i>	961
Yasha Savelyev , <i>Proof of the index conjecture in Hofer geometry</i>	981
Gabor Wiese , <i>An application of Maeda's conjecture to the inverse Galois problem</i>	985