

The algebra of derivations of quasi-modular forms from mirror symmetry*

MURAD ALIM, VADYM KURYLENKO, AND MARTIN VOGRIN

Abstract: We study moduli spaces of mirror non-compact Calabi-Yau threefolds enhanced with choices of differential forms. The differential forms are elements of the middle dimensional cohomology whose variation is described by a variation of mixed Hodge structures which is equipped with a flat Gauss-Manin connection. We construct graded differential rings of special functions on these moduli spaces and show that they contain rings of quasi-modular forms. We show that the algebra of derivations of quasi-modular forms can be obtained from the Gauss-Manin connection contracted with vector fields on the enhanced moduli spaces. We provide examples for this construction given by the mirrors of the canonical bundles of \mathbb{P}^2 and \mathbb{F}_2 .

Keywords: Calabi-Yau, modular forms, mixed Hodge structure, Gauss-Manin connection, mirror symmetry.

1. Introduction

The study of mirror symmetry, which was discovered in the context of string theory, has had a remarkable impact on mathematics putting forward new connections between different research areas such as symplectic and algebraic geometry. In particular the field of enumerative geometry has benefited greatly from the study of mirror symmetry starting from the predictions of the numbers of rational curves of arbitrary degree on the quintic from the study of variation of Hodge structure of the mirror family [11], see also [15, 51] and references therein.

A particularly fruitful arena of studies has been the interplay of modularity and geometry in the study of generating functions appearing in the

Received December 29, 2021.

2010 Mathematics Subject Classification: Primary 14D07, 14J15, 14J32; secondary 14J33.

*Research supported by DFG Emmy-Noether grant on “Building blocks of physical theories from the geometry of quantization and BPS states”, number AL 1407/2-1.

enumerative geometry of families of Calabi-Yau (CY) manifolds. The quasi-modular forms studied by Kaneko and Zagier [28] appear for instance explicitly in the study of higher genus mirror symmetry of families of elliptic curves [18]. The appearance of quasi-modular forms has moreover been a recurrent theme in the study of higher genus Gromov-Witten invariants for families of CY threefolds. The modularity of Gromov-Witten generating functions of CY families is related to the appearance of elliptic curves or K3 surfaces in the geometries. This can be either explicitly as factors of product spaces as for instance in [9], see also [37] and references therein as well as [47] for a recent treatment. Alternatively the CY families are elliptic or K3 fibrations as in e. g. [26, 25, 33, 5, 32]. In the case of mirror symmetry of non-compact CY families the modularity can be traced to the appearance of mirror curves, a locus where the mirror families of canonical bundles over Fano manifolds degenerate. The latter modularity has been extensively studied, see e. g. [27, 1, 6, 14], as well as [22] and references therein.

For higher genus mirror symmetry of generic families of CY threefolds a set of special functions on the moduli spaces which form a differential ring, i. e. are closed under derivations, was put forward for the mirror quintic by Yamaguchi and Yau [52] and generalized to arbitrary families in [3]. It was shown that the generating functions of higher genus Gromov-Witten invariants are polynomial in these special functions. An isomorphic differential ring of special functions for the mirror quintic was independently put forward by Movasati [39] (see also [41]),¹ working on the moduli space of the mirror quintic enhanced with choices of differential forms. The motivation of Movasati's construction came from the study of the geometric realization of modular forms which we briefly outline in the following.

In [38], Movasati defined the moduli space \mathbb{T} of pairs $(E, (\omega_1, \omega_2))$, where E is an elliptic curve and (ω_1, ω_2) is a basis of $H^1(E, \mathbb{C})$ such that ω_1 is holomorphic and the intersection pairing of ω_1 and ω_2 is 1. It was observed that the ring of regular function $\mathcal{O}_{\mathbb{T}}$ is closely related to the ring of quasi-modular forms and can be equipped with a differential structure coming from a special vector field on \mathbb{T} , equivalent to the Ramanujan differential relations between the Eisenstein series

$$\partial_{\tau} E_2 = \frac{E_2^2 - E_4}{12}, \quad \partial_{\tau} E_4 = \frac{E_4 E_2 - E_6}{3}, \quad \partial_{\tau} E_6 = \frac{E_6 E_2 - E_4^2}{2},$$

where $\partial_{\tau} := \frac{1}{2\pi i} \frac{\partial}{\partial \tau}$. Moreover, two other natural differential structures on $\mathcal{O}_{\mathbb{T}}$ were identified in [40], such that all together they constitute an $\mathfrak{sl}_2(\mathbb{C})$ Lie

¹The equivalence of the two constructions was shown in [4] and generalized to other families.

algebra. This Lie algebra is a realization of the $\mathfrak{sl}_2(\mathbb{C})$ triple of derivations acting on the ring of quasi-modular forms, see for instance [53].

A similar construction can be carried out for more general varieties. Consider a pair (X, ω) , where X is a compact Calabi-Yau variety of dimension n of fixed topological type, i.e. X is diffeomorphic to some fixed variety X_0 , and ω is a basis of $H^n(X, \mathbb{C})$, respecting the Hodge filtration \mathcal{F}^\bullet (see Definition 9) and such that the intersection pairing in this basis takes the form of a fixed constant matrix. Let \mathbb{T} be a moduli space of such pairs. Analogously to the case of elliptic curves, there is a set of distinguished differential structures on the ring of regular functions $\mathcal{O}_{\mathbb{T}}$. They are defined by the special (modular) vector fields R_i on \mathbb{T} , defined by

$$(1.1) \quad \nabla_{R_i} \omega_t = \begin{pmatrix} 0_{f_n, f_n} & *_{f_n, f_{n-1}} & 0_{f_n, f_{n-2}} & \cdots & 0_{f_n, f_0} \\ 0_{f_{n-1}, f_n} & 0_{f_{n-1}, f_{n-1}} & *_{f_{n-1}, f_{n-2}} & \cdots & 0_{f_{n-1}, f_0} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0_{f_1, f_n} & 0_{f_1, f_{n-1}} & 0_{f_1, f_{n-2}} & \cdots & *_{f_1, f_0} \\ 0_{f_0, f_n} & 0_{f_0, f_{n-1}} & 0_{f_0, f_{n-2}} & \cdots & 0_{f_0, f_0} \end{pmatrix} \omega_t,$$

where ω_t is a frame of $\mathcal{H}^n(X/\mathbb{T})$ defined in the section 3.2, $f_k = \text{rk } \mathcal{F}^k / \mathcal{F}^{k+1}$ and $*_{a,b}$ is an $a \times b$ matrix with entries in $\mathcal{O}_{\mathbb{T}}$.

The construction of moduli spaces, together with the modular vector fields on them was carried out for a number of compact families: compact CY threefolds in [39, 4, 2], Dwork families in [44, 46], genus 2 hyperelliptic curves in [12], K3 surfaces in [2, 46], elliptic K3 surfaces in [19, 7] and abelian varieties in [20]. The program of constructing moduli spaces of varieties enhanced with differential forms and investigating their properties was named Gauss-Manin Connection in Disguise (GMCD), see [42].

Besides the (modular) vector field, there are other distinguished vector fields inducing differential structures on $\mathcal{O}_{\mathbb{T}}$. Let \mathbb{G} be the following algebraic group

$$\mathbb{G} = \{g \in \text{GL}(b_n, \mathbb{C}) \mid g \text{ preserves filtration and } g\Phi g^{\text{tr}} = \Phi\},$$

where $b_n = \dim H^n(X, \mathbb{C})$ is the n -th Betti number, Φ is a constant skew-symmetric matrix, and let $\text{Lie}(\mathbb{G})$ denote its Lie algebra. The special vector fields R_g on \mathbb{T} can be defined as

$$\nabla_{R_g} \omega_t = g\omega_t, \quad g \in \text{Lie}(\mathbb{G}).$$

Any vector field on \mathbb{T} can be viewed as a derivation on $\mathcal{O}_{\mathbb{T}}$. We refer to the algebra of derivations coming from the vector fields R_g and the modular

vector fields R_i as the Gauss–Manin Lie algebra.² It is a generalization of the algebra of derivations on the ring of quasi-modular forms to more general rings associated to CY varieties. The algebra of derivations for elliptic curves was characterized in the framework of GMCD in [40]. It was computed for compact CY threefolds in [4], for Dwork families in [44, 46] and for elliptic K3 surfaces in [7]. In [23] the framework of GMCD was studied in the case of elliptically fibered CY manifolds of various dimensions. In particular the authors of [23] identify an $\mathfrak{sl}_2(\mathbb{C})$ Lie subalgebra corresponding to the algebra of derivations of quasi-modular forms, which in their case is directly related to the modularity of the elliptic fiber. In the present work, we identify an $\mathfrak{sl}_2(\mathbb{C})$ Lie subalgebra of the Gauss–Manin Lie algebra which is more indirect and which is related to the modular structure of the mirror curves appearing in the study of mirror symmetry of non-compact CY threefolds.

This work is concerned with the study of the local structure of moduli spaces \mathbb{T} for non-compact CY threefolds mirror to toric CY threefolds. These are of the form

$$\mathcal{X} = \left\{ uv + \sum_{i=0}^{s-1} a_i y_i = 0 \mid u, v \in \mathbb{C}, y_i \in \mathbb{C}^*, \prod_{i=0}^{s-1} y_i^{Q_k^i} = 1 \right\},$$

where Q_k^i is the charge matrix defining the toric CY, s is the number of 1-dimensional cones in the corresponding fan and k runs from 1 to $s - 3$. The middle cohomology of such varieties does not carry a pure Hodge structure, but instead – a mixed Hodge structure. We construct locally a moduli space \mathbb{T} of non-compact CY varieties enhanced with differential forms. By (not necessarily compact) CY threefold enhanced with differential forms we refer to the pair (X, ω) , where X is a CY threefold and ω is a basis of its middle cohomology, respecting both the Hodge filtration F^\bullet and the weight filtration W_\bullet on $H^3(X, \mathbb{C})$. We further require that the pairing on $\mathrm{Gr}_i^W = W_i/W_{i-1}$ takes the form of some fixed constant matrix Φ_i for all i . As for compact families, the ring of regular functions $\mathcal{O}_{\mathbb{T}}$ can be equipped with a set of derivations coming from the modular vector fields (1.1), as well as the derivations corresponding to elements in $\mathrm{Lie}(\mathbf{G})$, giving the full Gauss–Manin Lie algebra.

The program is applied to two cases of interest, the mirror families to local \mathbb{P}^2 and local Hirzebruch surface \mathbb{F}_2 , as an illustration. The developed techniques are universal and can be generalized to other non-compact CY varieties, as well as variations of mixed Hodge structure associated to different geometries. For the mirrors of local \mathbb{P}^2 and \mathbb{F}_2 , the local structure of the

²Note that it has also been called AMSY Lie algebra in [46, 45].

moduli spaces can partially be characterized by the inclusion of differential rings of the mirror curves into the differential ring of the non-compact CY threefold family. The following structural theorem gives the isomorphism of differential rings in the case of the mirror of local \mathbb{P}^2 :

Theorem 1. *There is an isomorphism between the ring of regular functions $\mathcal{O}_{\mathbb{T}}$ for the mirror of local \mathbb{P}^2 equipped with the differential structure coming from a modular vector field \mathbf{R} on \mathbb{T} and the ring of quasi-modular forms on $\Gamma_0(3)$ with the derivation ∂_{τ} .*

For the mirror of local \mathbb{F}_2 we prove the following theorem, relating the differential ring of the family of non-compact CY threefold to the differential ring of the associated mirror curve:

Theorem 2. *The ring of regular functions $\mathcal{O}_{\mathbb{T}}$ for the mirror of local \mathbb{F}_2 equipped with the differential structures coming from modular vector fields \mathbf{R}_1 and \mathbf{R}_2 contains as a differential sub-ring the ring of quasi-modular forms on $\Gamma_0(2)$ with the derivation ∂_{τ} .*

Moreover, the Gauss-Manin Lie algebra is computed in both cases and the following theorem characterize its relation to the Gauss-Manin Lie algebra of the associated mirror curve:

Theorem 3. *There is an $\mathfrak{sl}_2(\mathbb{C})$ Lie subalgebra of the Gauss-Manin Lie algebra \mathfrak{G} for mirrors of local \mathbb{P}^2 and local \mathbb{F}_2 .*

Organization of the paper

This paper is organized as follows: In the second section we briefly review quasi-modular forms and construct the algebra of derivations on it. In the third section we review the theory of enhanced CY varieties and their moduli spaces. We recall in detail the construction for elliptic curves. In the fourth section we review the construction of variation of mixed Hodge structure for families of non-compact CY threefolds and propose a definition of enhanced CY varieties for non-compact situations, extending the GMCD program to the case of mixed Hodge structures. At the end of the article, we apply the construction to two families, mirror to non-compact CY threefolds and prove Theorems 1, 2 and 3.

2. Modular forms and differential operators

2.1. Basic definitions

In this section we review some basic definitions of quasi-modular forms and their associated algebra of derivations, see [28, 17, 53] for more background.

Let $\mathbb{H} = \{\tau \in \mathbb{C} \mid \text{Im } \tau > 0\}$ be the upper half plane and $\overline{\mathbb{H}} = \mathbb{H} \cup \mathbb{Q} \cup \{i\infty\}$ be the extended upper half plane, which is closed under the natural $\text{SL}_2(\mathbb{Z})$ action

$$\tau \rightarrow \gamma.\tau = \frac{a\tau + b}{c\tau + d}, \quad \text{for} \quad \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \gamma \in \text{SL}_2(\mathbb{Z}).$$

Let N be a positive integer, then the *principal congruence subgroup of level N* is

$$\Gamma(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}_2(\mathbb{Z}) : \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \pmod{N} \right\}.$$

A subgroup Γ of $\text{SL}_2(\mathbb{Z})$ is called a *congruence subgroup* if $\Gamma(N) \subseteq \Gamma$. For example,

$$(2.1) \quad \Gamma_0(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}_2(\mathbb{Z}) : \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \pmod{N} \right\}.$$

Definition 1. A *multiplier system* is a map $v : \text{SL}_2(\mathbb{Z}) \rightarrow \mathbb{C}^\times$, such that if we denote

$$j_{v,k}(\gamma, z) := v(\gamma)(cz + d)^k$$

for $\gamma \in \text{SL}_2(\mathbb{Z})$, then

$$j_{v,k}(\gamma_1\gamma_2, z) = j_{v,k}(\gamma_1, z)j_{v,k}(\gamma_2, z)$$

and

$$j_{v,k}(-\gamma, z) = (-1)^k j_{v,k}(\gamma, z).$$

Definition 2. A *modular form of weight $k \in \mathbb{Z}$* with respect to a congruence subgroup Γ and a multiplier system v is a function $f : \mathbb{H} \rightarrow \mathbb{P}^1$ such that the following conditions hold:

1. $f(\gamma.\tau) = v(\gamma)(c\tau + d)^k f(\tau)$, $\forall \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$, where $(c\tau + d)^k$ is called the automorphy factor,
2. f is holomorphic on \mathbb{H} ,
3. f is holomorphic at the cusps in the sense that the function

$$(c\tau + d)^{-k} f(\gamma.\tau)$$

is holomorphic at $\tau = i\infty$ for any $\gamma \in \text{PSL}_2(\mathbb{Z}) = \text{SL}_2(\mathbb{Z})/\{\pm I\}$.

The ring of modular forms for Γ and v is denoted by $\mathcal{M}(\Gamma, v)$. It is naturally graded by the weight and finitely generated. If the multiplier system v is trivial we simply omit it from notation. For example,

$$\mathcal{M}(\mathrm{SL}_2(\mathbb{Z})) = \bigoplus_{k \in 2\mathbb{Z}} \mathcal{M}_k(\mathrm{SL}_2(\mathbb{Z})) \cong \mathbb{C}[E_4, E_6]$$

is finitely generated by the Eisenstein series E_4 and E_6 defined by

$$E_{2k}(\tau) = 1 + \frac{2}{\zeta(1-2k)} \sum_{n=1}^{\infty} \frac{n^{2k-1}q^n}{1-q^n}, \quad \text{for } q = e^{2\pi i\tau},$$

where ζ is the Riemann ζ -function.

Definition 3. A *quasi-modular form* on Γ of weight $k \in \mathbb{Z}$ and depth $p \in \mathbb{N}$ is a holomorphic function $f : \mathbb{H} \rightarrow \mathbb{P}^1$, holomorphic at the cusps, such that there exist $p + 1$ holomorphic functions f_0, \dots, f_p , such that

$$(c\tau + d)^{-k} f(\gamma.\tau) = \sum_{i=0}^p f_i(\tau) \left(\frac{c}{c\tau + d} \right)^i, \quad \forall \gamma \in \Gamma.$$

If a multiplier system is involved, one just has to multiply the right-hand side by $v(\gamma)$.

Example 1. The Eisenstein series E_2 is a quasi-modular form of depth 1 and weight 2 for $\Gamma = \mathrm{SL}_2(\mathbb{Z})$

$$E_2(\gamma.\tau) = (c\tau + d)^2 E_2(\tau) + \frac{12c}{2\pi i} (c\tau + d).$$

The ring of quasi-modular forms for $\mathrm{SL}_2(\mathbb{Z})$ is finitely generated

$$\widetilde{\mathcal{M}}(\mathrm{SL}_2(\mathbb{Z})) \cong \mathbb{C}[E_2, E_4, E_6].$$

2.2. $\mathfrak{sl}_2(\mathbb{C})$ Lie algebra of derivations

Unlike the ring of modular forms, the ring of quasi-modular forms is closed under the action of $\partial_\tau := \frac{1}{2\pi i} \frac{\partial}{\partial \tau}$. For example, in the case of $\Gamma = \mathrm{SL}_2(\mathbb{Z})$ we have the following relations

$$(2.2) \quad \partial_\tau E_2 = \frac{E_2^2 - E_4}{12}, \quad \partial_\tau E_4 = \frac{E_4 E_2 - E_6}{3}, \quad \partial_\tau E_6 = \frac{E_6 E_2 - E_4^2}{2},$$

called the Ramanujan relations.

In addition to the derivation ∂_τ we can define two more derivations on $\widetilde{\mathcal{M}}(\mathrm{SL}_2(\mathbb{Z}))$, as it was done in [53, 54]. Due to the fact that $\widetilde{\mathcal{M}}(\mathrm{SL}_2(\mathbb{Z})) \cong \mathbb{C}[E_2, E_4, E_6]$, we can write any quasi-modular form f of weight k as a polynomial in E_2

$$f = P_f(E_2) = \sum_i a_n E_2^n,$$

with $a_n \in \mathcal{M}(\mathrm{SL}_2(\mathbb{Z}))$ of weight $k - 2n$. Define

$$\mathfrak{d}f = -12P'_f(E_2).$$

The other operator simply returns the weight of the quasi-modular form

$$W(f) = kf.$$

Proposition 1. *The operators ∂_τ , \mathfrak{d} and W defined above form an $\mathfrak{sl}_2(\mathbb{C})$ Lie algebra*

$$(2.3) \quad [W, \partial_\tau] = 2\partial_\tau, \quad [W, \mathfrak{d}] = -2\mathfrak{d}, \quad [\partial_\tau, \mathfrak{d}] = W.$$

It is possible to construct this algebra of derivations for the other congruence subgroups as well (see section 5.3 in [53]).

Remark 4. There exists a similar Lie algebra of derivations in the case of Jacobi modular forms, see [48].

2.3. Differential rings of quasi-modular forms

In this section we review the differential rings of quasi-modular forms described in [36] (see also [6]) for the congruence subgroups $\Gamma_0(N)$ with $N = 1^*, 3, 2, 3$. For suitable multiplier systems on $\Gamma_0(N)$ (see loc.cit.) consider the following weight one modular forms:

N	A	B	C
1^*	$E_4(\tau)^{1/4}$	$\left(\frac{E_4(\tau)^{3/2} + E_6(\tau)}{2}\right)^{1/6}$	$\left(\frac{E_4(\tau)^{3/2} - E_6(\tau)}{2}\right)^{1/6}$
2	$\frac{(64\eta(2\tau)^{24} + \eta(\tau)^{24})^{1/4}}{\eta(\tau)^2 \eta(2\tau)^2}$	$\frac{\eta(\tau)^4}{\eta(2\tau)^2}$	$2^{3/2} \frac{\eta(2\tau)^4}{\eta(\tau)^2}$
3	$\frac{(27\eta(3\tau)^{12} + \eta(\tau)^{12})^{1/3}}{\eta(\tau)\eta(3\tau)}$	$\frac{\eta(\tau)^3}{\eta(3\tau)}$	$3 \frac{\eta(3\tau)^3}{\eta(\tau)}$

³Here 1^* corresponds to $\Gamma_0(1)^*$ – the unique normal index 2 subgroup of $\Gamma(1)$.

where $\eta(\tau)$ denotes the Dedekind η -function

$$\eta(\tau) = \left(\frac{E_4(\tau)^3 - E_6(\tau)^2}{1728} \right)^{\frac{1}{24}}.$$

Although some of these functions are not single-valued, their squares are. Moreover, the modular forms B and C can be written in terms of A and a weight 0 modular form α – the Hauptmodul. The following holds:

$$B = (1 - \alpha)^{1/r}A, \quad C = \alpha^{1/r}A,$$

where $r = 6$ for $N = 1^*$, $r = 4$ for $N = 2$ and $r = 3$ for $N = 3$.

Define also the analogue of the Eisenstein series E_2 as

$$(2.4) \quad E = \partial_\tau \log B^r C^r.$$

Definition 4. By a differential ring of quasi-modular forms on $\Gamma_0(N)$ we would mean a polynomial ring modulo the above relations:

$$R_N = \mathbb{C}[\alpha, A, B, C, E]/(B^r - (1 - \alpha)A^r, C^r - \alpha A^r)$$

with differential relations given by

$$(2.5) \quad \begin{aligned} \partial_\tau \alpha &= \alpha(1 - \alpha)A^2, \\ \partial_\tau A &= \frac{1}{2r}A \left(E + \frac{C^r - B^r}{A^{r-2}} \right), \\ \partial_\tau B &= \frac{1}{2r}B(E - A^2), \\ \partial_\tau C &= \frac{1}{2r}C(E + A^2), \\ \partial_\tau E &= \frac{1}{2r}(E^2 - A^4). \end{aligned}$$

The other derivations on the ring are given by

$$(2.6) \quad H : f \mapsto kf,$$

and

$$(2.7) \quad F : f \mapsto -2r\partial_E P_f(E),$$

for a weight k quasi-modular form $f \in \widetilde{\mathcal{M}}(\Gamma_0(N))$, where $P_f(E)$ is viewed as a formal expansion of f in E , as in the section 2. The Lie algebra of derivations is $\mathfrak{sl}_2(\mathbb{C})$.

3. Moduli space of enhanced CY varieties

3.1. Hodge structures

Definition 5. A *pure \mathbb{Z} -Hodge structure* of weight $n \in \mathbb{Z}$ is a pair $(H_{\mathbb{Z}}, F^{\bullet})$, where $H_{\mathbb{Z}}$ is a lattice, i. e. a finitely generated free abelian group, and F^{\bullet} is a decreasing (Hodge) filtration on $H_{\mathbb{C}} = H_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{C}$

$$0 \subset F^n \subset \dots \subset F^0 = H_{\mathbb{C}},$$

such that $F^p \oplus \overline{F^{n-p+1}} \simeq H_{\mathbb{C}}$.

Definition 6. A *polarization* of a \mathbb{Z} -Hodge structure $(H_{\mathbb{Z}}, F^{\bullet})$ is a bilinear form

$$\langle -, - \rangle : H_{\mathbb{Z}} \otimes H_{\mathbb{Z}} \rightarrow \mathbb{Z},$$

such that:

- It is symmetric if n is even and skew-symmetric if n is odd,
- The orthogonal complement of F^p is F^{n-p+1} ,
- For $\alpha, \beta \in H_{\mathbb{C}}$ the hermitian form $\langle C\alpha, \overline{\beta} \rangle$ is positive definite, where C is the so-called Weil operator which acts on $H^{p,q} := F^p \cap \overline{F^q}$ by multiplication with i^{p-q} .

A Hodge structure is called *polarizable* if it admits a polarization.

Definition 7. A *variation of Hodge structure* on a complex variety \mathbb{B} is a pair $(\mathcal{H}_{\mathbb{Z}}, \mathcal{F}^{\bullet})$ such that

- $\mathcal{H}_{\mathbb{Z}}$ is a locally constant sheaf of finitely generated \mathbb{Z} -modules on \mathbb{B} ,
- \mathcal{F}^{\bullet} is a finite decreasing filtration on $\mathcal{H} = \mathcal{H}_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathcal{O}_{\mathbb{B}}$ by holomorphic subbundles,

such that

- For each $b \in \mathbb{B}$ the fibers \mathcal{F}_b^{\bullet} form a Hodge filtration on $(\mathcal{H}_{\mathbb{Z}})_b \otimes_{\mathbb{Z}} \mathbb{C}$,
- The Gauss-Manin connection $\nabla : \mathcal{H} \rightarrow \mathcal{H} \otimes \Omega_{\mathbb{B}}^1$ defined by

$$\nabla(s \otimes f) = s \otimes df,$$

for $s \in \mathcal{H}_{\mathbb{Z}}$ and $f \in \mathcal{O}_{\mathbb{B}}$, satisfies the Griffiths' transversality condition

$$\nabla \mathcal{F}^p \subset \mathcal{F}^{p-1} \otimes_{\mathcal{O}_{\mathbb{B}}} \Omega_{\mathbb{B}}^1.$$

A variation of Hodge structure is said to be *polarizable* if there exists a ∇ -flat bilinear pairing

$$\langle -, - \rangle : \mathcal{H}_{\mathbb{Z}} \otimes \mathcal{H}_{\mathbb{Z}} \rightarrow \mathbb{Z},$$

such that it polarizes $(\mathcal{H}_{\mathbb{Z}})_b$ for each $b \in \mathbb{B}$.

Definition 8. A *Calabi-Yau variety* is a non-singular quasi-projective variety X , such that the canonical sheaf ω_X is trivial and $H^i(X, \mathcal{O}_X) = 0$ for $0 < i < \dim X$.

Example 2. Let $\pi : \mathcal{X} \rightarrow \mathbb{B}$ be a family of compact CY varieties of dimension n . The middle cohomology of a general fibre $X_b = \pi^{-1}(b)$, $b \in \mathbb{B}$ carries a Hodge structure $(H^n(X_b, \mathbb{Z}), F^\bullet(X_b))$, naturally polarized by

$$(3.1) \quad \langle \alpha, \beta \rangle = (-1)^{\frac{n(n-1)}{2}} \int_{X_b} \alpha \wedge \beta \quad \alpha, \beta \in H^n(X_b, \mathbb{Z}),$$

and there is a polarized variation of Hodge structure $(\mathcal{H}_{\mathbb{Z}}^n(\mathcal{X}/\mathbb{B}), \mathcal{F}^\bullet)$, where $\mathcal{H}_{\mathbb{Z}}^n(\mathcal{X}/\mathbb{B}) = R^n \pi_* \mathbb{Z}$, and \mathcal{F}^\bullet are unique vector subbundles of $\mathcal{H}^n(\mathcal{X}/\mathbb{B}) = \mathcal{H}_{\mathbb{Z}}^n(\mathcal{X}/\mathbb{B}) \otimes_{\mathbb{Z}} \mathcal{O}_{\mathbb{B}}$ such that $\mathcal{F}_b^\bullet = F^\bullet(X_b)$. The variation of Hodge structure is naturally polarized by (3.1).

3.2. Enhanced families

Definition 9. An *enhanced compact CY variety* of dimension n is a pair (X, ω) , where X is a compact CY variety of dimension n and

$$\omega = (\omega_1, \dots, \omega_i, \dots, \omega_{b_n})^{\text{tr}}, \quad b_n = \dim H^n(X, \mathbb{C}),$$

is a basis of $H^n(X, \mathbb{C})$ compatible with the Hodge filtration, i. e.

$$(\omega_1, \dots, \omega_{\dim F^k})$$

spans a basis of F^k , and such that

$$\langle \omega, \omega \rangle = \Phi,$$

for a constant non-degenerate matrix Φ . We call a basis ω satisfying the above conditions *compatible*.

In this paper we focus on the local structure of the moduli space \mathbb{T} rather than its global structure which requires a more careful study. We will denote by \mathcal{X}/\mathbb{T} the associated family of enhanced CY varieties over it. For construction of the moduli space \mathbb{T} in some specific examples we refer for example to [43]. Consider a family \mathcal{X}/\mathbb{B} of smooth CY varieties as in Example 2. Around any point $b \in \mathbb{B}$ we can choose a Zariski open neighborhood $V_b \subset \mathbb{B}$ and a global frame ω_b of the vector bundle $\mathcal{H}^n(\mathcal{X}/\mathbb{B})|_{V_b}$, such that at every point b' of V_b the basis $\omega_b(b')$ is compatible. All other frames with this property can be constructed by a transformation of ω_b by a non-singular lower block-triangular $b_n \times b_n$ matrix

$$(3.2) \quad S = \begin{pmatrix} *_{f_n, f_n} & 0_{f_n, f_{n-1}} & 0_{f_n, f_{n-2}} & \cdots & 0_{f_n, f_0} \\ *_{f_{n-1}, f_n} & *_{f_{n-1}, f_{n-1}} & 0_{f_{n-1}, f_{n-2}} & \cdots & 0_{f_{n-1}, f_0} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ *_{f_1, f_n} & *_{f_1, f_{n-1}} & *_{f_1, f_{n-2}} & \cdots & 0_{f_1, f_0} \\ *_{f_0, f_n} & *_{f_0, f_{n-1}} & *_{f_0, f_{n-2}} & \cdots & *_{f_0, f_0} \end{pmatrix},$$

where $f_k = \text{rk } \mathcal{F}^k / \mathcal{F}^{k+1}$ and $*_{a,b}$ is an $a \times b$ matrix, satisfying

$$(3.3) \quad S\Phi S^{\text{tr}} = \Phi.$$

We denote the corresponding frame by $\omega_t = S\omega_b$. Define U to be the underlying complex variety of

$$\text{Spec} \left(\mathbb{C} \left[s_{ij}^{\text{ind}}, \frac{1}{\det S} \right] \right),$$

where s_{ij}^{ind} are the algebraically independent entries of the matrix S , then a patch $\tilde{\mathbb{T}}$ of the moduli space \mathbb{T} can be constructed as the fiber product

$$\tilde{\mathbb{T}} = V_b \times_{\mathbb{C}} U.$$

Since we are going to work only with structures that do not require a construction of the global moduli space \mathbb{T} , we choose an analytic neighborhood V_0 inside of V_{b_0} and an analytic neighborhood $U_0 \subseteq U$ of the point $S = \text{Id}$. Further in this paper we abuse notation and by \mathbb{T} we mean

$$\mathbb{T} = V_0 \times_{\mathbb{C}} U_0.$$

On the algebraic de Rham cohomology $\mathcal{H}^n(\mathcal{X}/\mathbb{T})$ there exists an algebraic Gauss-Manin connection [29]

$$(3.4) \quad \nabla : \mathcal{H}^n(\mathcal{X}/\mathbb{T}) \rightarrow \mathcal{H}^n(\mathcal{X}/\mathbb{T}) \otimes_{\mathcal{O}_{\mathbb{T}}} \Omega_{\mathbb{T}}^1,$$

where \mathcal{O}_T is the ring of regular functions and Ω_T^1 is an \mathcal{O}_T -module of differential 1-forms on T . Using the local description of T , we can calculate the Gauss-Manin connection (3.4) from the Gauss-Manin connection on $\mathcal{H}^n(\mathcal{X}/\mathbb{B})$. Let

$$\nabla\omega_b = A\omega_b,$$

with an $\Omega_{\mathbb{B}}^1$ -valued matrix A , then in the basis ω_t we have

$$(3.5) \quad \nabla\omega_t = (dS + SA)S^{-1}\omega_t.$$

3.3. Gauss-Manin Lie algebra

It was shown in [40] that in the case of elliptic curves the ring of regular functions \mathcal{O}_T equipped with a differential structure coming from a certain vector field R on T is isomorphic, as a differential ring, to the ring of quasi-modular forms of [28] with derivation ∂_τ . In other words, the Ramanujan differential equations (2.2) are equivalent to the existence of a (modular) vector field R on T . There exists a similar algebraic structure on the rings of regular functions \mathcal{O}_T for more general varieties. It can be formalized by introducing special (modular) vector fields on T .

Definition 10. A modular vector field R is a rational vector field on T such that

$$(3.6) \quad \nabla_R\omega_t = \begin{pmatrix} 0_{f_n, f_n} & *_{f_n, f_{n-1}} & 0_{f_n, f_{n-2}} & \cdots & 0_{f_n, f_0} \\ 0_{f_{n-1}, f_n} & 0_{f_{n-1}, f_{n-1}} & *_{f_{n-1}, f_{n-2}} & \cdots & 0_{f_{n-1}, f_0} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0_{f_1, f_n} & 0_{f_1, f_{n-1}} & 0_{f_1, f_{n-2}} & \cdots & *_{f_1, f_0} \\ 0_{f_0, f_n} & 0_{f_0, f_{n-1}} & 0_{f_0, f_{n-2}} & \cdots & 0_{f_0, f_0} \end{pmatrix} \omega_t,$$

where $f_k = \text{rk } \mathcal{F}^k / \mathcal{F}^{k+1}$ and $*_{a,b}$ is an $a \times b$ matrix with entries in \mathcal{O}_T .

Moreover, it is possible to define a set of vector fields, corresponding to derivations on \mathcal{O}_T , generalizing (2.3).

Let G be the algebraic group consisting of all matrices S defined above,

$$G = \{g \in \text{GL}(b_n, \mathbb{C}) \mid g \text{ is of the form (3.2) and } g\Phi g^{\text{tr}} = \Phi\}.$$

Theorem 5. [4] For every element $g \in \text{Lie}(G)$ there is a unique vector field R_g on T such that

$$\nabla_{R_g}\omega_t = g\omega_t.$$

Definition 11. The Gauss-Manin Lie algebra \mathfrak{G} is an \mathcal{O}_T -module generated by the vector fields R_g and the modular vector fields.

3.4. Elliptic curve

We review the case of a family of elliptic curves following the work of [40]. Consider an affine variety in the weighted projective space $\mathbb{P}(1, 2, 3)$ given by

$$(3.7) \quad \left\{ W = x_1^6 + x_2^3 + x_3^2 - z^{-1/6}x_1x_2x_3 = 0 \right\}, \quad z \in \mathbb{P}^1 \setminus \left\{ 0, \frac{1}{432}, \infty \right\},$$

with x_i being the homogeneous coordinates on $\mathbb{P}(1, 2, 3)$. There is an action of the finite group $\mathbb{Z}_2 \times \mathbb{Z}_3$ on it given by

$$\mathbb{Z}_2 : (x_1, x_2, x_3) \rightarrow (\mu_2x_1, x_2, \mu_2x_3), \quad \mathbb{Z}_3 : (x_1, x_2, x_3) \rightarrow (\mu_3^2x_1, \mu_3x_2, x_3),$$

where μ_i is an i -th root of unity. A resolution of the quotient family gives rise to a family of smooth elliptic curves with a general fiber E_z . The monodromy group of this family is $\Gamma_0(1)^*$ – the unique index two subgroup of $\mathrm{SL}_2(\mathbb{Z})$. The periods of the holomorphic differential

$$\Omega = \mathrm{Res}_{W=0} \left(\frac{-x_1dx_2 \wedge dx_3 + 2x_2dx_1 \wedge dx_3 - 3x_3dx_1 \wedge dx_2}{W} \right),$$

are annihilated by the following Picard-Fuchs operator

$$\theta^2 - 12z(6\theta + 5)(6\theta + 1), \quad \theta = z \frac{d}{dz}.$$

Fix a basis

$$\omega = (\Omega, (1 - 432z)\nabla_\theta\Omega)^{\mathrm{tr}},$$

of $H^1(E_z, \mathbb{C})$. It respects the Hodge filtration, i. e. $\Omega \in F^1, (1 - 432z)\nabla_\theta\Omega \in F^0$ and the pairing is

$$\int_{E_z} \Omega \wedge (1 - 432z)\nabla_\theta\Omega = -1.$$

In other words, the pairing matrix is

$$\Phi = \langle \omega_i, w_j \rangle = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix},$$

in this basis. The transformations of the basis preserving filtration condition are of the form

$$S = \begin{pmatrix} s_{11} & 0 \\ s_{21} & s_{22} \end{pmatrix}.$$

The pairing condition (3.3) implies

$$S = \begin{pmatrix} s_{22}^{-1} & 0 \\ s_{21} & s_{22} \end{pmatrix}.$$

Therefore the moduli space \mathbb{T} of enhanced elliptic curves is three-dimensional and we use a coordinate patch $\{z, s_{21}, s_{22}\}$.

Proposition 2. *There is a unique vector field R on \mathbb{T} , such that*

$$\nabla_R \omega_t = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \omega_t.$$

It is given by

$$R = (1 - 432z)z s_{22}^2 \frac{\partial}{\partial z} - s_{21} s_{22}^2 \frac{\partial}{\partial s_{22}} - 60z(1 - 432z) s_{22}^3 \frac{\partial}{\partial s_{21}}.$$

Proof. Existence and uniqueness follow from a direct computation. □

We consider the group

$$G = \left\{ S = \begin{pmatrix} s_{22}^{-1} & 0 \\ s_{21} & s_{22} \end{pmatrix} \mid s_{22} \in \mathbb{C}^*, s_{21} \in \mathbb{C} \right\}.$$

Its Lie algebra is easily computable and has two generators

$$\mathfrak{g}_{22} = \left(\frac{\partial}{\partial s_{22}} S \right) S^{-1} \Big|_{S=\text{Id}} = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix},$$

$$\mathfrak{g}_{21} = \left(\frac{\partial}{\partial s_{21}} S \right) S^{-1} \Big|_{S=\text{Id}} = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}.$$

To find the corresponding vector fields, we look for solutions of the system

$$\left(i_{1*} \nabla \frac{\partial}{\partial z} + i_{2*} \nabla \frac{\partial}{\partial s_{22}} + i_{3*} \nabla \frac{\partial}{\partial s_{21}} \right) \omega_t = \mathfrak{g}_{2*} \omega_t,$$

of four linear equations for three variables i_{1*}, i_{2*}, i_{3*} . Note, that if there is a solution it is unique. We find the following:

$$R_{\mathfrak{g}_{22}} = s_{22} \frac{\partial}{\partial s_{22}} + s_{21} \frac{\partial}{\partial s_{21}}, \quad R_{\mathfrak{g}_{21}} = \frac{1}{s_{22}} \frac{\partial}{\partial s_{21}}.$$

The vector field $R_{\mathfrak{g}_{22}}$ is the so-called radial vector field.

Proposition 3. *The algebra generated by the vector fields $R, R_{\mathfrak{g}_{22}}, R_{\mathfrak{g}_{21}}$ is isomorphic to the $\mathfrak{sl}_2(\mathbb{C})$ Lie algebra.*

Proof. The proof follows from the explicit evaluation of the commutation relations. Explicitly, we find

$$[R, R_{\mathfrak{g}_{21}}] = R_{\mathfrak{g}_{22}}, \quad [R, R_{\mathfrak{g}_{22}}] = -2R, \quad [R_{\mathfrak{g}_{21}}, R_{\mathfrak{g}_{22}}] = 2R_{\mathfrak{g}_{21}}. \quad \square$$

The construction of this section leads to version of a theorem of [40] for the family (3.7).

Theorem 6. *There is a ring isomorphism between the ring of regular functions \mathcal{O}_T equipped with differential structures coming from the vector fields $R, R_{\mathfrak{g}_{22}}, R_{\mathfrak{g}_{21}}$ and the differential ring R_{1^*} with the differential structures described in the section 2.3.*

Proof. The isomorphism is given by sending (z, s_{22}, s_{21}) to

$$\alpha = 432z, \quad A = s_{22}, \quad E = (1 - 864z)s_{22}^2 - 12s_{21}s_{22}.$$

Then modular vector field R is equivalent to the differential relations from 2.3. Moreover, the action of $R_{\mathfrak{g}_{22}}$ and $R_{\mathfrak{g}_{21}}$ is equivalent to the action of W and \mathfrak{d} respectively. \square

4. Non-compact CY varieties and mixed Hodge structure

4.1. Local CY and mirror families

In this work we consider toric CY threefolds which are given by the total space of a bundle over a surface or a curve, we call these local CY threefolds. Toric varieties can be described by a fan Σ , but substantial amount of information about it is encapsulated by a charge matrix Q_k^i that encodes the linear relations between one dimensional cones $\Sigma(1)$ of the fan Σ , i. e. $\sum_{i=0}^{s-1} Q_k^i v_i = 0$ for $v_i \in \Sigma(1)$ and $s = |\Sigma(1)|$. The mirror families of such varieties are known to have the following form, see e. g. [31, 30, 13, 24]:

$$(4.1) \quad \mathcal{X} = \left\{ uv + F_{\mathbf{a}}(y_i) = uv + \sum_{i=0}^{s-1} a_i y_i = 0 \mid u, v \in \mathbb{C}, y_i \in \mathbb{C}^*, \prod_{i=0}^{s-1} y_i^{Q_k^i} = 1 \right\}.$$

It is also common to set as a mirror family a family of curves

$$(4.2) \quad \Sigma = \left\{ F_{\mathbf{a}}(y_i) = \sum_{i=0}^{s-1} a_i y_i = 0 \mid y_i \in \mathbb{C}^*, \prod_{i=0}^{s-1} y_i^{Q_k^i} = 1 \right\}.$$

We are interested in a special class of local CY varieties, namely canonical bundles over del Pezzo surfaces. In this case all the information can be encoded in a 2-dimensional reflexive polytope Δ . In particular, the set of the one dimensional cones v_i of the fan Σ corresponds to the set $A(\Delta)$ of the integral points m_i of the polytope Δ , thus the coordinates y_i can be parametrized by

$$y_i = t^{m_i} := t_1^{m_{i,1}} t_2^{m_{i,2}}, \quad t_1, t_2 \in \mathbb{C}^*,$$

for $m_i = (m_{i,1}, m_{i,2}) \in A(\Delta)$. Hence the information about the general fibre $X_{\mathbf{a}}$ of \mathcal{X} is encoded in the polynomial

$$F_{\mathbf{a}}(t_1, t_2) = \sum_{m_i \in A(\Delta)} a_i t^{m_i}.$$

The parameters a_i redundantly describe the deformations of complex structure on $X_{\mathbf{a}}$. The GIT quotient of the space of parameters a_i by the natural torus action induced by the action $F(t_1, t_2) \rightarrow \lambda_0 F(\lambda_1 t_1, \lambda_2 t_2)$ gives the complex structure moduli space \mathbf{B} , for which the torus-invariant local coordinates can be chosen as $z_k = (-1)^{Q_k^0} \prod_i a_i^{Q_k^i}$.

The holomorphic 3-form on $X_{\mathbf{a}}$ is given by the residue of the top form on the ambient space

$$\Omega = \text{Res} \left[\frac{1}{F_{\mathbf{a}}(t_1, t_2) + uv} \frac{dt_1}{t_1} \wedge \frac{dt_2}{t_2} \wedge du \wedge dv \right],$$

where Res is the residue map

$$\text{Res} : H^4(\mathbb{C}^2 \times (\mathbb{C}^*)^2 \setminus X_{\mathbf{a}}) \rightarrow H^3(X_{\mathbf{a}}),$$

which sends an $\omega \in H^4(\mathbb{C}^2 \times (\mathbb{C}^*)^2 \setminus X_{\mathbf{a}})$ to $\int_{\gamma} \omega$, with $\gamma \in H_4(\mathbb{C}^2 \times (\mathbb{C}^*)^2 \setminus X_{\mathbf{a}})$ being a cycle around $uv + F_{\mathbf{a}}(t_1, t_2) = 0$. The periods of the holomorphic 3-form satisfy Picard-Fuchs equations, which are obtained from a GKZ system [21] of differential equations

$$\left[\prod_{i:Q_k^i>0} \left(\frac{\partial}{\partial a_i} \right)^{Q_k^i} - \prod_{i:Q_k^i<0} \left(\frac{\partial}{\partial a_i} \right)^{-Q_k^i} \right] \int_{\gamma} \Omega = 0, \quad \gamma \in H_3(X_{\mathbf{a}}, \mathbb{Z}).$$

4.2. Mixed Hodge structure

In contrast to the compact CY varieties, cohomology groups of non-compact CY varieties are naturally endowed with a mixed Hodge structure introduced by [16].

Definition 12. A *mixed Hodge structure* $(H_{\mathbb{Z}}, W_{\bullet}, F^{\bullet})$ is a \mathbb{Z} -module $H_{\mathbb{Z}}$ together with an increasing (weight) filtration W_{\bullet} on $H_{\mathbb{Q}} = H_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{Q}$ and a decreasing (Hodge) filtration F^{\bullet} , which defines a pure \mathbb{Q} -Hodge structure of weight i on the graded piece $\text{Gr}_i^W = W_i/W_{i-1}$.

A mixed Hodge structure $(H_{\mathbb{Z}}, W_{\bullet}, F^{\bullet})$ is called *graded-polarizable* if the induced Hodge structure on Gr_i^W is polarizable for all i .

Definition 13. A *variation of mixed Hodge structure* on a complex variety B is a triple $(\mathcal{H}_{\mathbb{Z}}, \mathcal{W}_{\bullet}, \mathcal{F}^{\bullet})$, where

1. $\mathcal{H}_{\mathbb{Z}}$ is a locally constant sheaf of finitely generated \mathbb{Z} -modules on B ,
2. \mathcal{W}_{\bullet} is an increasing filtration of $\mathcal{H}_{\mathbb{Q}} = \mathcal{H}_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{Q}$ by locally constant subsheaves,
3. \mathcal{F}^{\bullet} is a decreasing filtration on $\mathcal{H} = \mathcal{H}_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathcal{O}_B$ by holomorphic subbundles, satisfying Griffiths' transversality,

such that for each $b \in B$ the filtrations \mathcal{F}_b^{\bullet} and $\mathcal{W}_{\bullet,b}$ define a mixed Hodge structure. A variation of mixed Hodge structure is called *graded-polarizable* if there exists a pairing $\langle -, - \rangle$, flat with respect to the Gauss-Manin connection ∇ on each $\text{Gr}_i^{\mathcal{W}} = \mathcal{W}_i/\mathcal{W}_{i-1}$.

A mixed Hodge structure on the middle cohomology $H^3(X_{\mathbf{a}}, \mathbb{C})$ for the families (4.1) was described in [8, 35] (see also [50]). Define the following ring:

$$\mathbf{S}_{\Delta} = \bigoplus_{k \geq 0} \mathbf{S}_{\Delta}^k, \quad \mathbf{S}_{\Delta}^k = \bigoplus_{m \in A(\Delta(k))} \mathbb{C} t_0^k t^m,$$

where t_0 is an additional parameter and

$$\Delta(k) := \left\{ m \in \mathbb{Z}^2 \subseteq \mathbb{R}^2 \mid \frac{m}{k} \in \Delta \right\}$$

for $k \geq 1$, $\Delta(0) := \{0\} \subset \mathbb{R}^2$. The grading is given by $\deg(t_0^k t^m) = k$. Define also the following differential operators on \mathbf{S}_{Δ}

$$\mathcal{D}_0 := \theta_{t_0} + t_0 F_{\mathbf{a}}, \quad \mathcal{D}_i := \theta_{t_i} + t_0 \theta_{t_i} F_{\mathbf{a}}, \quad i = 1, 2,$$

and a \mathbb{C} -vector space

$$\mathcal{R}_{F_{\mathbf{a}}} := \mathbf{S}_{\Delta} / \left(\sum_{i=0}^{s-1} \mathcal{D}_i \mathbf{S}_{\Delta} \right).$$

There is a decreasing filtration on $\mathcal{R}_{F_{\mathbf{a}}}$

$$0 \subset \mathcal{E}^0 \subset \mathcal{E}^{-1} \subset \mathcal{E}^{-2} \subset \dots,$$

with \mathcal{E}^{-k} being a subspace generated by monomials of degree $\leq k$.

There is also an increasing filtration. Define $I_\Delta^{(j)}$ to be the homogeneous ideals in \mathbf{S}_Δ generated as \mathbb{C} -subspaces by all monomials $t_0^k t^m$ ($k \geq 1$) with $m \in \Delta(k)$ which does not belong to any face of codimension j . Since everything belongs to the codimension 0 face, we get $I_\Delta^{(0)} = 0$. Everything that belongs to the face of codimension 2 also belongs to some face of codimension 1, therefore $I_\Delta^{(1)} \subset I_\Delta^{(2)}$. There are no faces of codimension 3, thus $I_\Delta^{(3)} = \bigoplus_{k \geq 1} \mathbf{S}_\Delta^k$, which contains the last two, and set $I_\Delta^{(4)} = \mathbf{S}_\Delta$. These form an increasing filtration on \mathbf{S}_Δ :

$$0 = I_\Delta^{(0)} \subset I_\Delta^{(1)} \subset I_\Delta^{(2)} \subset I_\Delta^{(3)} \subset I_\Delta^{(4)} = \mathbf{S}_\Delta,$$

which defines under the quotient an increasing filtration on $\mathcal{R}_{F_\mathbf{a}}$:

$$0 = \mathcal{I}_0 \subset \mathcal{I}_1 \subset \mathcal{I}_2 \subset \mathcal{I}_3 \subset \mathcal{I}_4 = \mathcal{R}_{F_\mathbf{a}},$$

where \mathcal{I}_j is the image of $I_\Delta^{(j)}$ in $\mathcal{R}_{F_\mathbf{a}}$.

Let us consider $\mathbf{S}_\Delta[\mathbf{a}] = \mathbf{S}_\Delta \otimes \mathbb{C}[\mathbf{a}]$ and

$$\mathcal{R}_{F_\mathbf{a}}[\mathbf{a}] := \mathbf{S}_\Delta[\mathbf{a}] / \left(\sum_{i=0}^n \mathcal{D}_i \mathbf{S}_\Delta[\mathbf{a}] \right).$$

On $\mathbf{S}_\Delta[\mathbf{a}]$ we can define the following differential operators:

$$\mathcal{D}_{a_i} := \frac{\partial}{\partial a_i} + t_0 t^{m_i}.$$

Since they commute with \mathcal{D}_i , they descend to $\mathcal{R}_{F_\mathbf{a}}[\mathbf{a}]$.

Theorem 7 ([35]). *There is an isomorphism*

$$r : \mathcal{R}_{F_\mathbf{a}} \cong H^3(X_\mathbf{a}, \mathbb{C}).$$

Explicitly for every \mathbf{a} it is given by

$$r(t_0^k t^m) = \text{Res}_{X_\mathbf{a}} \left(\frac{(-1)^k k! \cdot t^m}{(F_\mathbf{a}(t_1, t_2) + uv)^{k+1}} \frac{dt_1}{t_1} \wedge \frac{dt_2}{t_2} \wedge du \wedge dv \right).$$

Therefore, as a consequence, the filtrations $\mathcal{E}_\bullet, \mathcal{I}_\bullet$ define a mixed Hodge structure on $H^3(X_\mathbf{a}, \mathbb{C})$, namely $F^{6-k} = r(\mathcal{E}^{-k})$ and $W_3 = r(\mathcal{I}_1)$, $W_4 = W_5 = r(\mathcal{I}_3)$, $W_6 = r(\mathcal{I}_4)$. Moreover the derivations \mathcal{D}_{a_i} correspond to the Gauss-Manin connection $\nabla_{\frac{\partial}{\partial a_i}}$.

Remark 8 ([35]). The weight 3 filtration space W_3 is in fact just the first cohomology of the corresponding curve $\overline{\Sigma}_{\mathbf{a}}$, which is the compactification of a fibre of (4.2) at \mathbf{a} :

$$W_3H^3(X_{\mathbf{a}}) \cong H^1(\overline{\Sigma}_{\mathbf{a}}).$$

For the families (4.1) there is an inclusion $W_3H^3(X_{\mathbf{a}}) \cong H^1(\overline{\Sigma}_{\mathbf{a}}) \subset H^3(X_{\mathbf{a}})$. We define a polarization $\langle -, - \rangle$ to be equal to the intersection pairing on $W_3H^3(X_{\mathbf{a}})$ and 0 otherwise. It was shown in [35] that this pairing is flat on Gr_i^W with respect to the Gauss-Manin connection. Since in the case of local del Pezzo all the mirror curves are elliptic curves (see [13]), in a suitable basis the pairing matrix is

$$\begin{pmatrix} 0_{(b_n-2) \times (b_n-2)} & 0_{(b_n-2) \times 2} \\ 0_{2 \times (b_n-2)} & \begin{matrix} 0 & -1 \\ 1 & 0 \end{matrix} \end{pmatrix},$$

where $0_{a \times b}$ is an $a \times b$ block of zeroes.

4.3. Enhanced non-compact CY varieties

An important property of the intersection pairing in the case of compact CY varieties is that it is flat with respect to the Gauss-Manin connection. In the case of variation of mixed Hodge structure we no longer have this property, but we have flatness of the pairing on the graded components $\text{Gr}_i^W = \mathcal{W}_i/\mathcal{W}_{i-1}$. This fact motivates the following definition.

Definition 14. An *enhanced CY variety* of dimension n is a pair (X, ω) , where X is a CY variety of dimension n and

$$\omega = (\omega_1, \dots, \omega_i, \dots, \omega_{b_n})^{\text{tr}}, \quad b_n = \dim H^n(X, \mathbb{C}),$$

is a basis of $H^n(X, \mathbb{C})$ such that:

- It respects the Hodge filtration, i. e.

$$(\omega_1, \dots, \omega_{\dim F^k})$$

spans F^k ;

- it respects the weight filtration, i. e.

$$(\omega_{b_n - \dim W_{k+1}}, \dots, \omega_{b_n})$$

spans W_k ;

- the pairing $\langle -, - \rangle$ on Gr_i^W takes the form of a constant matrix Φ_i in this basis for all i .

The construction of the moduli space of enhanced varieties, as well as the Gauss-Manin Lie algebra generalize to this setting. The difference is the additional constraints imposed on the entries of S by the second condition above. It sets some of the lower-diagonal blocks to zero, in a similar fashion as the Hodge filtration condition sets the upper-diagonal blocks of S to zero. Moreover, the pairing condition of the Definition 9 is imposed on each graded component of the weight filtration separately.

5. Examples from mirror symmetry

The construction of section 4 is applied to two families of non-compact CY threefolds, mirrors to local \mathbb{P}^2 and local \mathbb{F}_2 . We construct the rings of regular functions on \mathbb{T} and the Gauss-Manin Lie algebra and prove Theorems 1, 2 and 3.

5.1. Local \mathbb{P}^2

5.1.1. Setup The first example we study is the total space of the canonical bundle over the projective plane \mathbb{P}^2 . It is defined by the toric charge vectors $Q = (-3, 1, 1, 1)$ and the mirror family is given by

$$\mathcal{X} = \left\{ uv + a_0y_0 + a_1y_1 + a_2y_2 + a_3y_3 = 0 \mid u, v \in \mathbb{C}, y_k \in \mathbb{C}^*, \frac{y_1y_2y_3}{y_0^3} = 1 \right\}.$$

We have

$$F_{\mathbf{a}}(t_1, t_2) = a_0 + a_1t_1 + a_2t_2 + \frac{a_3}{t_1t_2}.$$

The torus invariant coordinate reads

$$z = -\frac{a_1a_2a_3}{a_0^3},$$

and the Picard-Fuchs operator is

$$\mathcal{L} = (\theta^2 - 3z(3\theta + 1)(3\theta + 2))\theta, \quad \theta = z \frac{d}{dz}.$$

The middle cohomology can be identified as

$$H^3(X_z) \cong \mathcal{R}_{F_{\mathbf{a}}} = \mathbb{C}1 \oplus \mathbb{C}t_0 \oplus \mathbb{C}t_0^2,$$

by the theorem 7, where we denote the general fibre of \mathcal{X} by X_z to emphasize the dependence on z . The mixed Hodge structure described in the previous section is

$$0 = W_2 \subset W_3 = W_4 = W_5 = \mathbb{C}t_0 \oplus \mathbb{C}t_0^2 \subset W_6 = \mathcal{R}_{F_a},$$

$$0 = F^4 \subset F^3 = \mathbb{C}1 \subset F^2 = \mathbb{C}1 \oplus \mathbb{C}t_0 \subset F^1 = F^0 = \mathcal{R}_{F_a}.$$

We can generate it with the help of \mathcal{D}_{a_0} :

$$\mathcal{D}_{a_0}(1) = \left(\frac{\partial}{\partial a_0} + t_0 \right) (1) = t_0, \quad \mathcal{D}_{a_0}(t_0) = t_0^2.$$

There is a patch of the moduli space \mathbb{B} , where

$$\theta_{a_0} = -3\theta = -3z \frac{d}{dz} = -3\theta,$$

holds. Thus we can take $(\Omega, \nabla_\theta \Omega, \nabla_\theta^2 \Omega)$ as a basis of $H^3(X_z)$ that satisfies first two conditions from the Definition 14. The only non-zero pairing is between $\nabla_\theta \Omega$ and $\nabla_\theta^2 \Omega$. It is the Yukawa coupling Y_{111} for local \mathbb{P}^2 , namely

$$\langle \nabla_\theta^2 \Omega, \nabla_\theta \Omega \rangle = Y_{111} = -\frac{1}{3(1 - 27z)}.$$

In the basis

$$\omega = \left(\Omega, \nabla_\theta \Omega, Y_{111}^{-1} \nabla_\theta^2 \Omega \right),$$

the pairing has the form

$$\begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}.$$

5.1.2. Moduli space \mathbb{T} Let us fix $\Phi_3 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ and $\Phi_6 = \begin{pmatrix} 0 \end{pmatrix}$. We saw in the previous subsection that ω is a basis satisfying the conditions of the Definition 14, therefore we can construct (locally) the moduli space \mathbb{T} by considering the complex structure modulus z and algebraically independent entries of the matrix S . The requirement on S to preserve the Hodge filtration restricts it to the lower-diagonal form

$$S = \begin{pmatrix} s_{11} & 0 & 0 \\ s_{21} & s_{22} & 0 \\ s_{31} & s_{32} & s_{33} \end{pmatrix}.$$

The second condition – preservation of the weight filtration, sets $s_{21} = s_{31} = 0$. Next, we have to satisfy the condition that

$$\langle -, - \rangle|_{\text{Gr}_3^W} = \Phi_3, \quad \langle -, - \rangle|_{\text{Gr}_6^W} = \Phi_6,$$

note that Gr_4^W and Gr_5^W are empty. The second condition is empty, and the first one implies $s_{22} = s_{33}^{-1}$, thus we have

$$S = \begin{pmatrix} 1 & 0 & 0 \\ 0 & s_{33}^{-1} & 0 \\ 0 & s_{32} & s_{33} \end{pmatrix}.$$

The element s_{11} corresponds to the normalization of the holomorphic 3-form. One can check that it is constant with respect to the modular vector field and we set it to be 1 for simplicity.

5.1.3. Modular vector field and Gauss-Manin Lie algebra Modular vector fields have the form

$$\nabla_{\mathbf{R}}\omega_{\mathbf{t}} = \begin{pmatrix} 0 & * & 0 \\ 0 & 0 & * \\ 0 & 0 & 0 \end{pmatrix} \omega_{\mathbf{t}}.$$

We make a choice of normalization inspired by [6]

$$(5.1) \quad \nabla_{\mathbf{R}}\omega_{\mathbf{t}} = \begin{pmatrix} 0 & Y^{-1}(\mathbf{t}) & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} \omega_{\mathbf{t}}.$$

The Gauss-Manin connection matrix in the basis ω reads

$$\nabla_{\theta}\omega = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & Y_{111} \\ 0 & 18z & 0 \end{pmatrix} \omega = A_{\theta}\omega.$$

Let

$$\mathbf{R} = i_z \frac{\partial}{\partial z} + i_{32} \frac{\partial}{\partial s_{32}} + i_{33} \frac{\partial}{\partial s_{33}},$$

then,

$$(5.2) \quad \nabla_{\mathbf{R}}\omega_{\mathbf{t}} = \left[\frac{i_z}{z} (SA_{\theta}S^{-1}) + i_{32} \frac{\partial S}{\partial s_{32}} S^{-1} + i_{33} \frac{\partial S}{\partial s_{33}} S^{-1} \right] \omega_{\mathbf{t}}.$$

Proposition 4. *There exists a unique modular vector field R on T . It is given by*

$$R = \frac{zs_{33}^2}{Y_{111}} \frac{\partial}{\partial z} - s_{32}s_{33}^2 \frac{\partial}{\partial s_{33}} + \frac{18zs_{33}^3}{Y_{111}} \frac{\partial}{\partial s_{32}}.$$

Proof. Using non-degeneracy of the Gauss-Manin connection we equate the right hand sides of the equations (5.1) and (5.2). In such a way one obtains a system of linear equations that has a unique solution for (i_z, i_{32}, i_{33}) . \square

Introduce $\dot{f} = df(R)$ for $f \in \mathcal{O}_T$, then the existence of R is equivalent to the following differential structure on the ring \mathcal{O}_T :

$$\begin{cases} \dot{z} = zs_{33}^2 Y_{111}^{-1}, \\ \dot{s}_{32} = 18zs_{33}^3 Y_{111}^{-1}, \\ \dot{s}_{33} = -s_{32}s_{33}^2. \end{cases}$$

Theorem 1. *There is an isomorphism between the ring of regular functions \mathcal{O}_T equipped with the differential structure R and the ring R_3 of quasi-modular forms on $\Gamma_0(3)$ with the derivation ∂_τ .*

Proof. Define

$$A = \sqrt{-3}s_{33}, \quad \alpha = 27z,$$

$$B = (1 - 27z)^{1/3}A, \quad C = 3z^{1/3}A, \quad E = d\log(B^3C^3)(R),$$

Then we can rewrite the differential ring as

$$\begin{aligned} \dot{\alpha} &= \alpha(1 - \alpha)A^2, \\ \dot{A} &= \frac{1}{6}(AE + C^3 - B^3), \\ \dot{B} &= \frac{1}{6}B(E - A^2), \\ \dot{C} &= \frac{1}{6}C(E + A^2), \\ \dot{E} &= \frac{1}{6}(E^2 - A^4), \end{aligned}$$

which is the differential ring R_3 of quasi-modular forms on $\Gamma_0(3)$ in 2.3. \square

5.1.4. Gauss-Manin Lie algebra

Theorem 3 (Part 1). *The Gauss-Manin Lie algebra for local \mathbb{P}^2 is spanned by*

$$\begin{aligned}
 R &= \frac{zs_{33}^2}{Y_{111}} \frac{\partial}{\partial z} - s_{32}s_{33}^2 \frac{\partial}{\partial s_{33}} + \frac{18zs_{33}^3}{Y_{111}} \frac{\partial}{\partial s_{32}}, \\
 R_{\mathfrak{g}_{33}} &= s_{33} \frac{\partial}{\partial s_{33}} + s_{32} \frac{\partial}{\partial s_{32}}, \\
 R_{\mathfrak{g}_{32}} &= s_{33} \frac{\partial}{\partial s_{32}},
 \end{aligned}$$

and it is isomorphic to $\mathfrak{sl}_2(\mathbb{C})$.

Proof. $\text{Lie}(\mathbb{G})$ is generated by

$$\begin{aligned}
 \mathfrak{g}_{33} &= \left(\frac{\partial}{\partial s_{33}} S \right) S^{-1} \Big|_{S=\text{Id}} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \\
 \mathfrak{g}_{21} &= \left(\frac{\partial}{\partial s_{32}} S \right) S^{-1} \Big|_{S=\text{Id}} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}.
 \end{aligned}$$

The corresponding vector fields are exactly $R_{\mathfrak{g}_{33}}$ and $R_{\mathfrak{g}_{32}}$. It can be checked explicitly that commutators are

$$[R, R_{\mathfrak{g}_{32}}] = R_{\mathfrak{g}_{33}}, \quad [R, R_{\mathfrak{g}_{33}}] = -2R, \quad [R_{\mathfrak{g}_{32}}, R_{\mathfrak{g}_{33}}] = 2R_{\mathfrak{g}_{32}}. \quad \square$$

5.2. Local \mathbb{F}_2

The canonical bundle over the second Hirzebruch surface $K_{\mathbb{F}_2}$ is defined by the toric charge vectors $Q_1 = (-2, 1, 0, 1, 0)$ and $Q_2 = (0, 0, 1, -2, 1)$ and the mirror family is given by

$$\mathcal{X} = \left\{ uv + \sum_{i=0}^4 a_i y_i = 0 \mid u, v \in \mathbb{C}, y_k \in \mathbb{C}^*, \frac{y_1 y_3}{y_0^2} = 1, \frac{y_2 y_4}{y_3^2} = 1 \right\}.$$

We have

$$F_{\mathbf{a}}(t_1, t_2) = a_0 + a_1 t_1 + a_2 t_2 + \frac{a_3}{t_1} + \frac{a_4}{t_1^2 t_2}.$$

The torus-invariant coordinates are

$$z_1 = \frac{a_1 a_3}{a_0^2}, \quad z_2 = \frac{a_2 a_4}{a_3^2},$$

and the Picard-Fuchs operators read

$$\begin{aligned} \mathcal{L}_1 &= \theta_1(\theta_1 - 2\theta_2) - 2z_1(2\theta_1 + 1)\theta_1, \\ \mathcal{L}_2 &= \theta_2^2 - z_2(\theta_1 - 2\theta_2 - 1)(\theta_1 - 2\theta_2), \quad \theta_i = z_i \frac{d}{dz_i}. \end{aligned}$$

The middle cohomology can be written as

$$H^3(X_z) \cong \mathcal{R}_{F_a} = \mathbb{C}1 \oplus \mathbb{C}t_0 \oplus \mathbb{C}\frac{t_0}{t_1} \oplus \mathbb{C}t_0^2.$$

The mixed Hodge structure is

$$0 = W_2 \subset W_3 = \mathbb{C}t_0 \oplus \mathbb{C}t_0^2 \subset W_4 = W_5 = W_3 \oplus \mathbb{C}\frac{t_0}{t_1} \subset W_6 = \mathcal{R}_{F_a},$$

$$0 = F^4 \subset F^3 = \mathbb{C}1 \subset F^2 = \mathbb{C}1 \oplus \mathbb{C}t_0 \oplus \mathbb{C}\frac{t_0}{t_1} \subset F^1 = F^0 = \mathcal{R}_{F_a}.$$

Using the Gauss-Manin connection we obtain

$$\mathcal{D}_{a_0}(1) = t_0, \quad \mathcal{D}_{a_0}(t_0) = t_0^2, \quad \mathcal{D}_{a_3}(1) = \frac{t_0}{t_1}.$$

On the moduli space \mathbf{B} there is a patch where

$$a_0 \frac{\partial}{\partial a_0} = -2\theta_1, \quad a_3 \frac{\partial}{\partial a_3} = \theta_1 - 2\theta_2,$$

therefore in the basis

$$\left(\Omega, \nabla_{(\theta_1 - 2\theta_2)}\Omega, \nabla_{\theta_1}\Omega, \nabla_{\theta_1}^2\Omega \right),$$

the pairing has the form

$$\begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -Y_{111} \\ 0 & 0 & Y_{111} & 0 \end{pmatrix},$$

where Y_{111} is the Yukawa coupling

$$Y_{111} = \frac{1}{(1 - 4z_1)^2 - 64z_1^2 z_2}.$$

5.2.1. Moduli space Let us fix

$$\Phi_3 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad \Phi_4 = (0), \quad \Phi_6 = (0).$$

In addition to the complex structure moduli z_1 and z_2 we need to identify the independent elements of the matrix S . The Hodge filtration condition restricts it to the lower-block diagonal form

$$S = \begin{pmatrix} s_{11} & 0 & 0 & 0 \\ s_{21} & s_{22} & s_{23} & 0 \\ s_{31} & s_{32} & s_{33} & 0 \\ s_{41} & s_{42} & s_{43} & s_{44} \end{pmatrix}.$$

The weight filtration condition gives

$$S = \begin{pmatrix} s_{11} & 0 & 0 & 0 \\ 0 & s_{22} & s_{23} & 0 \\ 0 & 0 & s_{33} & 0 \\ 0 & 0 & s_{43} & s_{44} \end{pmatrix}.$$

The conditions on the pairing on Gr_4^W and Gr_6^W are empty and the corresponding condition on Gr_3^W gives us the final form of S :

$$(5.3) \quad S = \begin{pmatrix} s_{11} & 0 & 0 & 0 \\ 0 & s_{22} & s_{23} & 0 \\ 0 & 0 & s_{44}^{-1} & 0 \\ 0 & 0 & s_{43} & s_{44} \end{pmatrix}.$$

Moreover, we again set $s_{11} = 1$.

5.2.2. Modular vector fields We define modular vector fields by

$$(5.4) \quad \nabla_{R_1} \omega_t = \begin{pmatrix} 0 & Y_1^{-1}(t) & Y_2^{-1}(t) & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix} \omega_t,$$

and

$$(5.5) \quad \nabla_{R_2} \omega_t = \begin{pmatrix} 0 & Y_3^{-1}(t) & Y_4^{-1}(t) & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \omega_t,$$

for some $Y_i(t) \in \mathcal{O}_T$.

Theorem 9. *There exist unique modular vector fields $R_1, R_2 \in \mathfrak{X}(T)$ satisfying (5.4) and (5.5).*

The proof is analogous to the proof of Proposition 4.

The existence of the modular vector fields is equivalent to two differential structures on \mathcal{O}_T . By defining

$$\partial_{\tau_1} f := df(R_1) \quad \text{and} \quad \partial_{\tau_2} f := df(R_2)$$

for $f \in \mathcal{O}_T$, we can write

$$(5.6) \quad \begin{aligned} \partial_{\tau_1} z_1 &= z_1 s_{44}^2 s_{22}^{-1} (s_{22} + (1 - 4z_1)(1 - 4z_2)s_{23} - 4z_1(1 + 4z_2)s_{22}), \\ \partial_{\tau_1} z_2 &= 2z_2(-1 + 4z_2)s_{44}^2 s_{22}^{-1} (s_{23} + 4z_1 s_{22}), \\ \partial_{\tau_1} s_{22} &= 4z_2 s_{44}^2 (s_{23} + 4z_1 s_{22}), \\ \partial_{\tau_1} s_{23} &= 2z_1 s_{44}^2 s_{22}^{-1} ((s_{22} + s_{23})^2 - 4z_2 s_{23}^2), \\ \partial_{\tau_1} s_{43} &= -2z_1 s_{44}^2 s_{22}^{-1} ((-1 + 4z_2)s_{23}(-s_{43} + 4z_2 s_{44}) \\ &\quad + s_{22}(s_{43} - 4z_2 s_{43} + 4z_2(1 - 10z_1 + 8z_1 z_2)s_{44})), \\ \partial_{\tau_1} s_{44} &= -s_{44}^2 s_{22}^{-1} (2z_1(-1 + 4z_2)s_{23}s_{44} + s_{22}(s_{43} + (-1 + 4z_2)s_{44})), \\ \partial_{\tau_2} z_1 &= z_1(1 - 4z_1)(1 - 4z_2)s_{44} s_{22}^{-1}, \\ \partial_{\tau_2} z_2 &= 2z_2(1 - 4z_2)s_{44} s_{22}^{-1}, \\ \partial_{\tau_2} s_{22} &= -4z_2 s_{44}, \\ \partial_{\tau_2} s_{23} &= s_{43} + 2z_1 s_{44} + 2z_1 s_{23} s_{44} s_{22}^{-1} + 8z_1 z_2 s_{44}(1 - s_{23} s_{22}^{-1}), \\ \partial_{\tau_1} s_{43} &= 2z_1(1 - 4z_2)s_{44} s_{22}^{-1}(s_{43} - 4z_2 s_{44}), \\ \partial_{\tau_2} s_{44} &= 2z_1(-1 + 4z_2)s_{44}^2 s_{22}^{-1}. \end{aligned}$$

Theorem 2. *The differential ring $(\mathcal{O}_T, \partial_{\tau_1}, \partial_{\tau_2})$ contains as differential subring the ring R_2 of quasi-modular forms on $\Gamma_0(2)$ (see 2.3).*

Proof. By introducing a new variable

$$u = \frac{64z_1^2 z_2}{(1 - z_1)^2},$$

and

$$A = \sqrt{2(1 - 4z_1)}s_{44}, \quad B = (1 - u)^{1/4}A, \quad C = u^{1/4}A,$$

$$E = \partial_{\tau_1} \log(B^4 C^4) = \frac{2 - 8s_{33}s_{43} - 8z_1(1 + 8z_2)}{s_{33}^2},$$

the differential ring relations can be rewritten as

$$\begin{aligned} \partial_{\tau_1} u &= u(1 - u)A^2, \\ \partial_{\tau_1} A &= \frac{1}{8}A\left(E + \frac{C^4 - B^4}{A^2}\right), \\ \partial_{\tau_1} B &= \frac{1}{8}B(E - A^2), \\ \partial_{\tau_1} C &= \frac{1}{8}C(E + A^2), \\ \partial_{\tau_1} E &= \frac{1}{8}(E^2 - A^4), \end{aligned}$$

which is the differential ring R_2 . Furthermore, one can check that all of the functions u, A, B, C, E are independent of τ_2 . □

5.2.3. Gauss-Manin Lie algebra

Theorem 3 (Part 2). *The Gauss-Manin Lie algebra for local \mathbb{F}_2 is isomorphic to the semi-direct product $\mathfrak{L}_V \rtimes \mathfrak{sl}_2(\mathbb{C})$, where \mathfrak{L}_V denotes the Lie algebra of type V , as classified in [10], corresponding to an ideal of \mathfrak{G} generated by \mathbf{R}_2 , $\mathbf{R}_{\mathfrak{g}_{22}}$ and $\mathbf{R}_{\mathfrak{g}_{23}}$.*

Proof. In this case $\text{Lie}(\mathbf{G})$ is generated by

$$\mathfrak{g}_{22} = \left(\frac{\partial}{\partial s_{22}} S \right) S^{-1} \Big|_{S=\text{Id}} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

$$\mathfrak{g}_{23} = \left(\frac{\partial}{\partial s_{23}} S \right) S^{-1} \Big|_{S=\text{Id}} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

$$\mathfrak{g}_{43} = \left(\frac{\partial}{\partial s_{43}} S \right) S^{-1} \Big|_{S=\text{Id}} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix},$$

$$\mathfrak{g}_{44} = \left(\frac{\partial}{\partial s_{44}} S \right) S^{-1} \Big|_{S=\text{Id}} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

The associated vector fields $R_{\mathfrak{g}_{ij}}$ on T defined by $\nabla_{R_{\mathfrak{g}_{ij}}} \omega_t = \mathfrak{g}_{ij} \omega_t$ are

$$R_{\mathfrak{g}_{22}} = s_{22} \frac{\partial}{\partial s_{22}} + s_{23} \frac{\partial}{\partial s_{23}},$$

$$R_{\mathfrak{g}_{23}} = s_{44}^{-1} \frac{\partial}{\partial s_{23}},$$

$$R_{\mathfrak{g}_{43}} = s_{44}^{-1} \frac{\partial}{\partial s_{43}},$$

$$R_{\mathfrak{g}_{44}} = s_{44} \frac{\partial}{\partial s_{44}} + s_{43} \frac{\partial}{\partial s_{43}}.$$

Together with R_1 and R_2 these vector fields form the Gauss-Manin Lie algebra \mathfrak{G} and the commutation relations are the following:

$$\begin{aligned} [R_1, R_{\mathfrak{g}_{44}}] &= -2R_1, & [R_1, R_{\mathfrak{g}_{43}}] &= R_{\mathfrak{g}_{44}}, & [R_{\mathfrak{g}_{43}}, R_{\mathfrak{g}_{44}}] &= 2R_{\mathfrak{g}_{43}}, \\ [R_1, R_2] &= 0, & [R_1, R_{\mathfrak{g}_{23}}] &= R_2, & [R_1, R_{\mathfrak{g}_{22}}] &= 0, \\ [R_{\mathfrak{g}_{43}}, R_2] &= R_{\mathfrak{g}_{23}}, & [R_{\mathfrak{g}_{43}}, R_{\mathfrak{g}_{23}}] &= 0, & [R_{\mathfrak{g}_{43}}, R_{\mathfrak{g}_{22}}] &= 0, \\ [R_{\mathfrak{g}_{44}}, R_2] &= R_2, & [R_{\mathfrak{g}_{44}}, R_{\mathfrak{g}_{23}}] &= -R_{\mathfrak{g}_{23}}, & [R_{\mathfrak{g}_{44}}, R_{\mathfrak{g}_{22}}] &= 0, \\ [R_2, R_{\mathfrak{g}_{23}}] &= 0, & [R_2, R_{\mathfrak{g}_{22}}] &= R_2, & [R_{\mathfrak{g}_{23}}, R_{\mathfrak{g}_{22}}] &= R_{\mathfrak{g}_{23}}. \end{aligned}$$

The first line above is the $\mathfrak{sl}_2(\mathbb{C})$ Lie algebra corresponding to the quasi-modular forms. The last line is the 3-dimensional algebra \mathfrak{L}_V according to the Bianchi’s classification and it is an ideal in the Gauss-Manin Lie algebra \mathfrak{G} . Therefore, we have

$$\mathfrak{G} \cong \mathfrak{L}_V \rtimes \mathfrak{sl}_2(\mathbb{C}). \quad \square$$

6. Conclusion

In this article we studied moduli spaces of mirror non-compact CY threefolds enhanced with choices of differential forms in their middle cohomology. Certain vector fields contracted with the Gauss-Manin connection give rise in

this context to a Lie-algebra which was put forward in [4] for arbitrary CY threefolds, generalizing an earlier similar construction for the elliptic curve [40] and the mirror quintic [39].

In particular the resulting Lie algebra in the case of elliptic curves [40] is isomorphic to the $\mathfrak{sl}_2(\mathbb{C})$ Lie algebra of derivations associated to quasi-modular forms. The appearance of this Lie algebra as a Lie subalgebra of the full Gauss–Manin Lie algebra for families of CY threefolds is often connected to the appearance of classical quasi-modular forms. An $\mathfrak{sl}_2(\mathbb{C})$ Lie subalgebra was for example studied in the context of elliptically fibered CY manifolds [23], where the modularity is originating from the elliptic fibers. In our work we explored a different source of classical modular objects⁴ which appear in the moduli spaces of CY geometries, namely the case of the mirror families of non-compact CY threefolds. The appearance of classical modular forms in this context is related to the mirror curve.

We focused on the cases where the mirror curves of the mirror geometries are of genus one and studied the enhanced moduli spaces and the Gauss–Manin connection of the variation of mixed Hodge structure. We identified the $\mathfrak{sl}_2(\mathbb{C})$ Lie subalgebra corresponding to the algebra of derivations of quasi-modular forms in the cases of the mirrors of local \mathbb{P}^2 and local \mathbb{F}_2 . We note that the mirror geometry of local \mathbb{F}_2 has two parameters, although the moduli space is only one-dimensional, this was already addressed, for instance, in [1] and is related to the fact that \mathbb{F}_2 has two distinct curve classes but only one non-trivial four-cycle. On the level of the Gauss–Manin Lie algebra we note that the additional parameter gives rise to an additional vector field, enlarging the $\mathfrak{sl}_2(\mathbb{C})$ algebra of derivations of the quasi-modular forms in this case. It would be interesting to find an interpretation in the context of modular forms for this enhancement.

It is natural to expect that the study of the analogous moduli spaces of enhanced mirror CY families where the mirror curve is of genus higher than one would give rise to algebraic structures associated to modular objects associated to higher genus curves. The GMCD program for certain genus two curves was recently studied in [12] giving rise to differential rings of Siegel modular forms and their associated algebraic structures. Differential rings attached to genus two curves were studied in [34, 49] in the context of mirror symmetry, it would be interesting to recover the corresponding Gauss–Manin Lie algebras in these cases.

⁴We use the terminology classical modular objects to refer to modular structures which have been widely studied, such as quasi-modular forms, Siegel and Jacobi forms. For generic compact CY manifolds the analogues of the modular structures are referred to as CY modular forms in [42] and are in general not expressible in terms of classical modular forms.

Acknowledgements

The authors would like to thank Florian Beck for helpful discussions. M.V. would like to thank IMPA, where part of this work was carried out, for hospitality. This research is supported by DFG Emmy-Noether grant on “Building blocks of physical theories from the geometry of quantization and BPS states”, number AL 1407/2-1.

References

- [1] MINA AGANAGIC, VINCENT BOUCHARD, and ALBRECHT KLEMM. Topological strings and (almost) modular forms. *Comm. Math. Phys.*, **277**(3):771–819, 2008. [MR2365453](#)
- [2] MURAD ALIM. Algebraic structure of tt^* equations for Calabi-Yau sigma models. *Comm. Math. Phys.*, **353**(3):963–1009, 2017. [MR3652481](#)
- [3] MURAD ALIM and JEAN D. LÄNGE. Polynomial structure of the (open) topological string partition function. *J. High Energy Phys.*, (10):045, 13, 2007. [MR2357933](#)
- [4] MURAD ALIM, HOSSEIN MOVASATI, EMANUEL SCHEIDEGGER, and SHING-TUNG YAU. Gauss-Manin connection in disguise: Calabi-Yau threefolds. *Comm. Math. Phys.*, **344**(3):889–914, 2016. [MR3508164](#)
- [5] MURAD ALIM and EMANUEL SCHEIDEGGER. Topological strings on elliptic fibrations. *Comm. Number Theory Phys.*, **8**(4):729–800, 2014. [MR3318389](#)
- [6] MURAD ALIM, EMANUEL SCHEIDEGGER, SHING-TUNG YAU, and JIE ZHOU. Special polynomial rings, quasi modular forms and duality of topological strings. *Adv. Theor. Math. Phys.*, **18**(2):401–467, 2014. [MR3273318](#)
- [7] MURAD ALIM and MARTIN VOGRIN. Gauss-Manin Lie algebra of mirror elliptic K3 surfaces. *arXiv preprint: arXiv:1812.03185*, 2018. [MR4270267](#)
- [8] VICTOR V. BATYREV. Dual polyhedra and mirror symmetry for Calabi-Yau hypersurfaces in toric varieties. *J. Algebraic Geom.*, **3**(3):493–535, 1994. [MR1269718](#)
- [9] MICHAEL BERSHADSKY, SERGIO CECOTTI, HIROSI OOGURI, and CUMRUN VAFA. Kodaira-Spencer theory of gravity and exact results for quantum string amplitudes. *Comm. Math. Phys.*, **165**(2):311–427, 1994. [MR1301851](#)

- [10] LUIGI BIANCHI. On the three-dimensional spaces which admit a continuous group of motions. *Gen. Relativity Gravitation*, **33**(12):2171–2253 (2002), 2001. [MR1900159](#)
- [11] PHILIP CANDELAS, XENIA DE LA OSSA, PAUL S. GREEN, and LINDA PARKES. A pair of Calabi-Yau manifolds as an exactly soluble superconformal theory. *Nucl. Phys.*, **B359**(1):21–74, 1991. [MR1115626](#)
- [12] JIN CAO, HOSSEIN MOVASATI, and SHING-TUNG YAU. Gauss-Manin connection in disguise: genus two curves. *arXiv preprint: arXiv:1910.07624*, 2019. [MR3643758](#)
- [13] TI-MING CHIANG, ALBRECHT KLEMM, SHING-TUNG YAU, and ERIC ZASLOW. Local mirror symmetry: calculations and interpretations. *Adv. Theor. Math. Phys.*, **3**(3):495–565, 1999. [MR1797015](#)
- [14] TOM COATES and HIROSHI IRITANI. Gromov-Witten invariants of local \mathbb{P}^2 and modular forms. *arXiv preprint arXiv:1804.03292*, 2018. [MR4301054](#)
- [15] DAVID A. COX and SHELDON KATZ. *Mirror symmetry and algebraic geometry*, volume 68 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1999. [MR1677117](#)
- [16] PIERRE DELIGNE. Théorie de Hodge. II. *Inst. Hautes Études Sci. Publ. Math.*, (**40**):5–57, 1971. [MR0498551](#)
- [17] FRED DIAMOND and JERRY SHURMAN. *A first course in modular forms*, volume 228 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2005. [MR2112196](#)
- [18] ROBBERT DIJKGRAAF. Mirror symmetry and elliptic curves. In *The moduli space of curves (Texel Island, 1994)*, volume 129 of *Progr. Math.*, pages 149–163. Birkhäuser Boston, Boston, MA, 1995. [MR1363055](#)
- [19] CHARLES F. DORAN, ANDREW HARDER, HOSSEIN MOVASATI, and URSULA WHITCHER. Humbert surfaces and the moduli of lattice polarized K3 surfaces. In *String-Math 2014*, volume 93 of *Proc. Sympos. Pure Math.*, pages 109–140. Amer. Math. Soc., Providence, RI, 2016. [MR3524237](#)
- [20] TIAGO J. FONSECA. Higher Ramanujan equations and periods of abelian varieties. *arXiv preprint: arXiv:1807.11044*, 2018.
- [21] ISRAEL M. GEL'FAND, ANDREI V. ZELEVINSKIĬ, and MIKHAIL M. KAPRANOV. Hypergeometric functions and toric varieties. *Funktsional. Anal. i Prilozhen.*, **23**(2):12–26, 1989. [MR1011353](#)

- [22] BABAK HAGHIGHAT. Mirror symmetry and modularity. *arXiv preprint arXiv:1712.00601*, 2017.
- [23] BABAK HAGHIGHAT, HOSSEIN MOVASATI, and SHING-TUNG YAU. Calabi-Yau modular forms in limit: elliptic fibrations. *Comm. Number Theory Phys.*, **11**(4):879–912, 2017. [MR3733768](#)
- [24] SHINOBU HOSONO. Local mirror symmetry and type IIA monodromy of Calabi-Yau manifolds. *Adv. Theor. Math. Phys.*, **4**(2):335–376, 2000. [MR1838444](#)
- [25] SHINOBU HOSONO. Counting BPS states via holomorphic anomaly equations. In *Calabi-Yau varieties and mirror symmetry (Toronto, ON, 2001)*, volume 38 of *Fields Inst. Commun.*, pages 57–86. Amer. Math. Soc., Providence, RI, 2003. [MR2019147](#)
- [26] SHINOBU HOSONO, MASA-HIKO SAITO, and ATSUSHI TAKAHASHI. Holomorphic anomaly equation and BPS state counting of rational elliptic surface. *Adv. Theor. Math. Phys.*, **3**(1):177–208, 1999. [MR1704198](#)
- [27] MIN-XIN HUANG and ALBRECHT KLEMM. Holomorphic anomaly in gauge theories and matrix models. *J. High Energy Phys.*, **(9)**:054, 33, 2007. [MR2342377](#)
- [28] MASANOBU KANEKO and Don Zagier. A generalized Jacobi theta function and quasimodular forms. In *The moduli space of curves (Texel Island, 1994)*, volume 129 of *Progr. Math.*, pages 165–172. Birkhäuser Boston, Boston, MA, 1995. [MR1363056](#)
- [29] NICHOLAS M. KATZ and TADAO ODA. On the differentiation of de Rham cohomology classes with respect to parameters. *J. Math. Kyoto Univ.*, **8**:199–213, 1968. [MR0237510](#)
- [30] SHELDON KATZ, PETER MAYR, and CUMRUN VAFA. Mirror symmetry and exact solution of 4D $N = 2$ gauge theories. I. *Adv. Theor. Math. Phys.*, **1**(1):53–114, 1997. [MR1489296](#)
- [31] ALBRECHT KLEMM, WOLFGANG LERCHE, PETER MAYR, CUMRUN VAFA, and NICHOLAS WARNER. Self-dual strings and $N = 2$ supersymmetric field theory. *Nucl. Phys. B*, **477**(3):746–764, 1996. [MR1413475](#)
- [32] ALBRECHT KLEMM, JAN MANSCHOT, and THOMAS WOTSCHKE. Quantum geometry of elliptic Calabi-Yau manifolds. *Comm. Number Theory Phys.*, **6**(4):849–917, 2012. [MR3068410](#)

- [33] ALBRECHT KLEMM and MARCOS MARIÑO. Counting BPS states on the Enriques Calabi-Yau. *Comm. Math. Phys.*, **280**(1):27–76, 2008. [MR2391190](#)
- [34] ALBRECHT KLEMM, MAXIMILIAN PORETSCHKIN, THORSTEN SCHIMANNEK, and MARTIN WESTERHOLT-RAUM. Direct integration for mirror curves of genus two and an almost meromorphic Siegel modular form. *arXiv preprint arXiv:1502.00557*, 2015. [MR3636672](#)
- [35] YUKIKO KONISHI and SATOSHI MINABE. Local B-model and mixed Hodge structure. *Adv. Theor. Math. Phys.*, **14**(4):1089–1145, 2010. [MR2821394](#)
- [36] ROBERT S. MAIER. Nonlinear differential equations satisfied by certain classical modular forms. *Manuscr. Math.*, **134**(1-2):1–42, 2011. [MR2745252](#)
- [37] MARCOS MARIÑO. Gromov-Witten invariants and modular forms. 2003.
- [38] HOSSEIN MOVASATI. On differential modular forms and some analytic relations between Eisenstein series. *Ramanujan J.*, **17**(1):53–76, 2008. [MR2439525](#)
- [39] HOSSEIN MOVASATI. Eisenstein type series for Calabi–Yau varieties. *Nucl. Phys.*, **B847**(2):460–484, 2011. [MR2774983](#)
- [40] HOSSEIN MOVASATI. Quasi-modular forms attached to elliptic curves, I. *Ann. Math. Blaise Pascal*, **19**(2):307–377, 2012. [MR3025138](#)
- [41] HOSSEIN MOVASATI. Modular-type functions attached to mirror quintic Calabi-Yau varieties. *Math. Z.*, **281**(3-4):907–929, 2015. [MR3421646](#)
- [42] HOSSEIN MOVASATI. *Gauss-Manin connection in disguise: Calabi-Yau modular forms*. Surveys of Modern Mathematics. International Press of Boston, Inc., 2017. [MR3643758](#)
- [43] HOSSEIN MOVASATI. Modular and automorphic forms & beyond. 2020. [MR4375351](#)
- [44] HOSSEIN MOVASATI and YOUNES NIKDELAN. Gauss-Manin connection in disguise: Dwork family. *arXiv preprint: arXiv:1603.09411*, 2016. [MR4310933](#)
- [45] YOUNES NIKDELAN. Rankin-Cohen brackets for Calabi-Yau modular forms. *arXiv preprint: arXiv:1912.12809*, 2019.
- [46] YOUNES NIKDELAN. Modular vector fields attached to Dwork family: $sl_2(\mathbb{C})$ Lie algebra. *Moscow Math. J.*, **20**(1):127–151, 2020. [MR4060315](#)

- [47] GEORG OBERDIECK and RAHUL PANDHARIPANDE. Curve counting on $K3 \times E$, the Igusa cusp form χ_{10} , and descendent integration. In *K3 surfaces and their moduli*, volume 315 of *Progr. Math.*, pages 245–278. Birkhäuser/Springer, [Cham], 2016. [MR3524171](#)
- [48] GEORG OBERDIECK and AARON PIXTON. Gromov-Witten theory of elliptic fibrations: Jacobi forms and holomorphic anomaly equations. *Geom. Topol.*, **23**(3):1415–1489, 2019. [MR3956895](#)
- [49] YONGBIN RUAN, YINGCHUN ZHANG, and JIE ZHOU. Genus two quasi-Siegel modular forms and Gromov-Witten theory of toric Calabi-Yau threefolds. *arXiv preprint arXiv:1911.07204*, 2019.
- [50] JAN STIENSTRA. Resonant hypergeometric systems and mirror symmetry. In *Integrable Systems and Algebraic Geometry (Kobe/Kyoto, 1997)*, pages 412–452. World Sci. Publ., River Edge, NJ, 1998. [MR1672077](#)
- [51] CUMRUN VAFA and SHING-TUNG YAU, editors. *Winter School on Mirror Symmetry, Vector Bundles and Lagrangian Submanifolds, volume 23 of AMS/IP Studies in Advanced Mathematics. American Mathematical Society, Providence, RI; International Press, Somerville, MA, 2001.* [MR1876062](#)
- [52] SATOSHI YAMAGUCHI and SHING-TUNG YAU. Topological string partition functions as polynomials. *JHEP*, **07**:047, 2004. [MR2095047](#)
- [53] DON ZAGIER. Elliptic modular forms and their applications. In *The 1-2-3 of Modular Forms, Universitext*, pages 1–103. Springer, Berlin, 2008. [MR2409678](#)
- [54] DON ZAGIER. Partitions, quasimodular forms, and the Bloch-Okounkov theorem. *Ramanujan J.*, **41**(1-3):345–368, 2016. [MR3574638](#)

Murad Alim
Fachbereich Mathematik
University of Hamburg
Bundesstr. 55, 20146
Hamburg
Germany
E-mail: murad.alim@uni-hamburg.de

Vadym Kurylenko
Geometry and Mathematical Physics
SISSA
Via Bonomea 265, 34136
Trieste
Italy
E-mail: vkurylen@sissa.it

Martin Vognin
Institut für Physik
Johannes Gutenberg-Universität Mainz
Staudingerweg 9, 55128
Mainz
Germany
E-mail: mvognin@uni-mainz.de