

Interplay between nonlinear potential theory and fully nonlinear elliptic PDEs

F. REESE HARVEY AND KEVIN R. PAYNE*

Dedicated to Blaine Lawson on the occasion of his 80th birthday

Abstract: We discuss one of the many topics that illustrate the interaction of Blaine Lawson’s deep geometric and analytic insights. The first author is extremely grateful to have had the pleasure of collaborating with Blaine over many enjoyable years. The topic to be discussed concerns the fruitful interplay between *nonlinear potential theory*; that is, the study of subharmonics with respect to a general constraint set in the 2-jet bundle and the study of subsolutions and supersolutions of a nonlinear (degenerate) elliptic PDE. The main results include (but are not limited to) the validity of the comparison principle and the existence and uniqueness to solutions to the relevant Dirichlet problems on domains which are suitably “pseudoconvex”. The methods employed are geometric and flexible as well as being very general on the potential theory side, which is interesting in its own right. Moreover, in many important geometric contexts no natural operator may be present. On the other hand, the potential theoretic approach can yield results on the PDE side in terms of non standard structural conditions on a given differential operator.

Keywords: Subequations, potential theory, fully nonlinear degenerate elliptic PDEs, comparison principles, viscosity solutions, admissibility constraints, monotonicity, duality, fiberegularity.

1 Introduction

2974

1.1 Potential theories: from calibrations to subequations

2976

Received March 25, 2022.

2010 Mathematics Subject Classification: Primary 35B51, 35J60, 35J70, 35D40, 31C45, 35E20.

*Partially supported by the Gruppo Nazionale per l’Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM) and the project: GNAMPA 2022 “Proprietà qualitative e quantitative per EDP non lineari: dai principi del massimo alla teoria di regolarità”.

1.2	Potential theoretic results suggested by complex analysis	2979
1.3	The Dirichlet problem	2981
1.4	Boundary pseudoconvexity	2986
1.5	Some examples of PDEs	2987
2	Fundamental aspects of nonlinear potential theory	2995
2.1	Subequations, subharmonics and duality	2996
2.2	Monotonicity	2998
2.3	Fiberegularity	3001
2.4	Subharmonic addition for quasiconvex functions	3002
2.5	A “tool kit” for \mathcal{F} -subharmonics (subsolutions)	3004
3	Comparison by monotonicity, duality and fiberegularity	3006
3.1	Statement and history of the general result	3006
3.2	Outline of the proof	3007
4	The correspondence principle	3009
	References	3013

1. Introduction

Our main aim is to give a partial survey of a research endeavor which was initiated in a trio of papers of Harvey and Lawson [22], [23] and [24] published in 2009 that has grown into a wide ranging investigation with many interesting and important avenues still to pursue. For simplicity of the exposition and in order to make the discussion more accessible to analysts, we will focus on the Euclidean setting of open subsets X of \mathbb{R}^n , although X could also be a Riemannian manifold as in [26] and [30], or an almost complex manifold as in [35]. We will emphasize the fruitful interplay between *nonlinear potential theory*; that is, the study of the family of \mathcal{F} -subharmonics with respect to a given *subequation (constraint set)*

$$(1.1) \quad \mathcal{F} \subset \mathcal{J}^2(X) := X \times \mathcal{J}^2 := X \times \mathbb{R} \times \mathbb{R}^n \times \mathcal{S}(n), \quad X \subset \mathbb{R}^n$$

and the study of solutions/subsolutions/supersolutions of a given *fully nonlinear (elliptic) PDE*

$$(1.2) \quad F(x, J_x^2 u) := F(x, u(x), Du(x), D^2 u(x)) = 0, \quad x \in X \subset \mathbb{R}^n$$

determined by a given *operator* $F \in C(\mathcal{J}^2(X))$. The equation (1.2) will also be written more succinctly as

$$(1.3) \quad F(J^2 u) = 0 \quad \text{on } X.$$

Here \mathcal{J}^2 is the vector space of 2-jets. We will use the notation $J_x^2 u$ for the second order Taylor development of u indifferently with respect to the differentiability of u . This interplay has been developed in detail in [12], [13], [11] and [14].

Given the fully nonlinear setting, one cannot expect solutions to be regular in general, and distribution theory is generally available for convex subequations or equations in divergence form. Hence all notions are to be interpreted pointwise in the *viscosity sense* that will be recalled in Definition 2.4 (see [34] for the equivalence of the distributional approach and the viscosity approach in the convex case).

There is a satisfying unification that comes from a potential theoretic (pluripotential theoretic) viewpoint as it includes classical (Laplacian) subharmonics, convex and quasiconvex functions as well as new geometric potential theories (some of which are useful for theoretical physics) as well as an immense universe of first and second order potential theories determined by classes of (degenerate) elliptic operators.

We now describe the main motivating principles. There are many opportunities for cross-fertilization and synergy between the potential theory and the operator theory. First, the conditions imposed on a constraint set \mathcal{F} correspond to and encode structural conditions on the operator F ; for instance, a convex constraint set \mathcal{F} corresponds to a concave operator F . Second, the subequation \mathcal{F} “frees” a given PDE from any particular form of F (many different F correspond to the same \mathcal{F}); this is an important point in the work of Krylov [52] on the general notion of ellipticity. Moreover, \mathcal{F} “liberates” the user from needing an operator F to apply nonlinear elliptic potential theory. Third, “forgetting” about the operator leads to interesting questions that at first glance might not seem important for operator theory and provides a “machine” for formulating new conjectures and theorems. For instance, taking one’s cue from known results in pluripotential theory or convex analysis, one is led to seek generalizations in other potential theoretic

situations as well. Some examples of this will be discussed in Subsection 1.2 below. In this way, one can find “welcome surprises” in the operator theory. Fourth, along with a rich abundance of geometrically motivated potential theories, there are many new PDEs to discover. For example, as will be discussed in Subsection 1.1, while every calibrated geometry has an underlying potential theory, known “natural” smooth operators are “rare gems”. Nevertheless, for any given subequation \mathcal{F} , one can construct several “non smooth” operators. One good example is the construction of the so-called *canonical operator* associated to a given subequation. This is a canonical construction that is scattered out, first in [26, Remark 14.11] and [25, Examples 3.4 and 3.5] and then better explained in [43, subsection on canonical operators in section 6] and [11, Proposition 11.17]. It includes the *truncated Laplacians*, among the many examples, so it might be referred to as the “canonical eigenvalue operator construction”. These operators are discussed a bit further in Example 1.16. Another good example is the *signed distance operator*

$$(1.4) \quad F(x, J) := \begin{cases} \text{dist}(J, \partial\mathcal{F}_x) & J \in \mathcal{F}_x \\ -\text{dist}(J, \partial\mathcal{F}_x) & J \in \mathcal{J}^2 \setminus \mathcal{F}_x \end{cases},$$

where

$$(1.5) \quad \mathcal{F}_x := \{J \in \mathcal{J}^2 : (x, J) \in \mathcal{F}\}$$

is the *fiber of \mathcal{F} over $x \in X$* . The operator (1.4) was studied in the pure second order case in Theorem 3.2 of [52]. Finally, in the rare cases when a natural operator F is known for a fixed \mathcal{F} -potential theory, the operator F will have much to say about the potential theory; for example, by taking derivatives of the equation.

Having stated the main aims and philosophical motivations, we proceed to describe the origins and objectives, along with key concepts, nice features, some results and significant examples which illustrate the theory. We begin with discussion of the origins of the investigation which led to a hierarchy of potential theories.

1.1. Potential theories: from calibrations to subequations

The story begins in calibrated geometry. The geometric side of calibrated geometry was developed to emphasize the *calibrated submanifolds* which are those submanifolds for which the *calibration* restricts to be the volume form. Said infinitesimally, a calibration ϕ of degree p restricts to be a function on

the Grassmannian of oriented p -planes where it attains a maximum value of one on the subset $G(\phi)$ of p -planes calibrated by ϕ . In turn, a submanifold is calibrated by ϕ if its tangent planes are calibrated by ϕ .

The basic example (beyond calibrated geodesics), going back to Wirtinger in the last century, is the Kähler/symplectic form on $\mathbb{C}^n = \mathbb{R}^{2n}$. Here the calibrated submanifolds are simply the complex curves in \mathbb{C}^n . This geometric example, which has an analytic side involving a rich and well developed *potential theory* (or more precisely a *pluripotential theory*), then cries out for a pluripotential theory for other calibrations, providing the impetus to search for general ϕ -potential theories.

The ϕ -subharmonic (or ϕ -plurisubharmonic) functions u are easy to define for smooth functions (see [22]). One simply requires that the restriction of u to ϕ -submanifolds is classically subharmonic (with respect to the Laplacian Δ). This imposes a constraint on the second derivative (Hessian matrix) D^2u of u at each point. This constraint condition is that D^2u restricts to have trace zero on any p -plane calibrated by ϕ . More precisely, by identifying a p -plane W with the orthogonal projection P_W onto W and by using the natural inner product on the space $\mathcal{S}(n)$ of second derivatives (i.e. the symmetric matrices), the second derivative constraint set determined by $G(\phi)$ is just the *polar cone*

$$(1.6) \quad G(\phi)^\circ := \{A \in \mathcal{S}(n) : \langle A, P_W \rangle := \text{tr}(A|_W) \geq 0, \quad \forall W \in G(\phi)\}$$

of the set $G(\phi)$. This C^∞ potential theory suffices for many purposes (see [22]) where it was noticed, as a (big) surprise, that the calibration plays a minor role subordinate to the set $G(\phi)$ of distinguished/calibrated p -planes, including the case of G being the full set of p -planes (see [24], [27], [28] and [31]), thus extending the realm of ϕ -potential theories to G -potential theories.

In fact, replacing $G(\phi)$ by any closed set $G \subset \mathcal{S}(n)$ and then defining “ G -submanifolds” and “ G -plurisubharmonic functions” as above, it is still possible to obtain a robust G -pluripotential theory. Again, the constraint condition on the second derivatives of a function u at each $x \in X$

$$(1.7) \quad \langle D^2u(x), P_W \rangle := \text{tr} \left(D^2u(x)|_W \right) \geq 0, \quad \forall W \in G; \quad \text{i.e., } D^2u(x) \in G^\circ,$$

defines G -plurisubharmonicity of u on X . These functions u are characterized by their restrictions to G -submanifolds M that are also minimal; u is subharmonic with respect to the induced Laplacian on M (see [33]). Therefore it is justified to state that

G -pluripotential theory is the correct pluripotential theory for the geometry of minimal G -submanifolds.

Note that if $G := G(\phi)$ with ϕ a calibration, each G -submanifold is automatically actually absolutely volume minimizing (see [21]) and hence a minimal submanifold. Potential theory based on a closed subset G of the p -Grassmanian will be referred to as the *geometric case*.

This unifies many known cases and includes lots of new interesting cases. First, as our basic motivation, this includes classical pluripotential theory in complex analysis by taking G to be the Grassmannian of complex lines in \mathbb{C}^n as a subset of real 2-planes. Second, taking G to be the full Grassmannian $\mathbb{G}(1, \mathbb{R}^n)$ of real lines in \mathbb{R}^n we include convex function theory, providing more precision to the well-known parallel between pseudoconvexity and convexity. Third, this leads to new surprising examples not related to calibrations. As an example, for each degree p , let G be the full Grassmannian of p -planes in \mathbb{R}^n . One obtains a p -pluripotential theory associated with the geometry of all minimal submanifolds of dimension p . Another noteworthy new example is Lagrangian pluripotential theory, defined by taking $G = \text{LAG}$ to be the set of Lagrangian n -planes in \mathbb{C}^n . This is the appropriate potential theory for minimal Lagrangian submanifolds [41]. These are the two cases where new natural polynomial operators were discovered. First, the *p -fold sum operator* whose domain is the subequation $\mathbb{G}(p, \mathbb{R}^n)^\circ$ (see [24, (10.12)]) and second the *Lagrangian Monge-Ampère operator* (see [24, (10.11)] and [41]).

For the next level of generality, one can focus entirely on the second derivative constraint set $\mathcal{F} \subset \mathcal{S}(n)$, which in the geometric case is given by the polar $\mathcal{F} = G^\circ$ of G . There is surprising simplicity here as well. Other than \mathcal{F} being closed, a single condition on \mathcal{F} described below, which is called *positivity* (P), is needed. This condition (P) ensures that the notion of \mathcal{F} -subharmonicity for upper semicontinuous u agrees with the definition $D^2u(x) \in \mathcal{F}$ for C^2 functions. Such sets \mathcal{F} are called *subequations in $\mathcal{S}(n)$* . This is the pure second order constant coefficient case. This condition, besides providing the weakest possible condition ensuring *coherence* between the two definitions of \mathcal{F} -subharmonicity should also be viewed as the weakest possible form of ellipticity.

As with classical potential theory (for the Laplacian), the regularity of a general G -subharmonic function should only be required to be upper semicontinuous. This extension, although carried out with *Dirichlet duality* and *subaffine functions* in [24], is equivalent to a viscosity theory formulation (see Remark 4.9 of [24]). The viscosity approach is more direct and can be used easily for potential theories for subequation constraint sets with variable coefficients and dependence on all the jet variables (as will be discussed in the next paragraph). However, the (Dirichlet) duality continues to be important. It clarifies the notion of superharmonics and leads to straightforward proofs

of the comparison principle. This is examined extensively in the constant coefficient setting in [11].

As a final step in the delineation of a hierarchy of potential theories, it is natural to consider the extension from a pure second order constraint on second derivatives in the Euclidian setting of open subsets $X \subset \mathbb{R}^n$ to the case of X being a Riemannian manifold [26] or an almost complex manifold [35]. In local coordinates, this constraint will have variable coefficients and may depend on the full Taylor development up to second order (not just second derivatives). Returning to the Euclidian setting, this suggests considering subequation constraint sets $\mathcal{F} \subset \mathcal{J}^2(X) = X \times \mathcal{J}^2$ and their associated potential theories. The needed axioms for a robust potential theory are given in Definition 2.1 Briefly stated, one adds two additional axioms (negativity (N) and topological stability (T)) to the positivity (P) and requiring that \mathcal{F} be closed. The interplay between nonlinear potential theory and fully nonlinear elliptic PDEs most naturally takes place at this level of the hierarchy. This will be discussed further, beginning in Subsection 1.3.

An important part of the story is that in studying this hierarchy of possible levels of potential theories; ϕ -subharmonics, G -subharmonics and finally \mathcal{F} -subharmonics, a distinguished (differential) operator F is missing from the picture. This absence of an operator has advantages (and disadvantages) which have been noted above, and will be amplified below.

1.2. Potential theoretic results suggested by complex analysis

A surprising number of results in complex analysis (of several variables) can be established in greater generality in nonlinear potential theory. We mention seven such topics that were suggested by results in pluripotential theory. Most will be stated in the pure second order case, but they provide impetus for investigating them at all levels of the potential theory hierarchy. Here we are sketchy, leaving many definitions to the references. Hence, rather than giving formal statements of theorems, we will recall the main results informally.

1) The Andreotti-Frankel Theorem for subequations $\mathcal{F} \subset \mathcal{S}(n)$: (see [27] for details). First, \mathcal{F} -convex domains can be defined. Then the notion of \mathcal{F} -free submanifolds extends that of totally real submanifolds and one has the following result.

An \mathcal{F} -convex domain has the homotopy type of a CW-complex of dimension less than or equal to the maximal dimension of an \mathcal{F} -free subspace.

This maximal dimension is easy to calculate in the multitude of examples.

2) The Levi Problem in p -convex geometry: In [31] this problem is solved, including the case of non-integer p . One has

Local p -convexity implies global p -convexity:

3) Boundary pseudoconvexity: For bounded domains Ω with smooth boundary the following local to global result was established in Theorem 5.12 of [26] for *cone subequations* $\mathcal{F} \subset \mathcal{S}(n)$:

If $\partial\Omega$ is strictly \mathcal{F} -pseudoconvex at each point, then $\partial\Omega$ has a smooth strictly \mathcal{F} -subharmonic defining function.

For subequations $\mathcal{F} \subset \mathcal{S}(n)$ which are not cones, boundary convexity for \mathcal{F} is governed by the asymptotic behavior of \mathcal{F} at infinity. This asymptotic behavior is captured by a new subequation $\vec{\mathcal{F}}$ which is a cone, so that the above can be applied. See also Corollary 11.8 of [26] (and an incomplete discussion after the proof) for the general case.

If strictness is dropped, then boundary pseudoconvexity does not imply existence of a global plurisubharmonic defining function in complex analysis. On the other hand, Forsternič recently proved that this is the case if $G = \mathbb{G}(p, \mathbb{R}^n)$ and $\mathcal{F} = G^\circ$ (see [18]). This is particularly interesting as it runs counter to our point made here that generally speaking several complex variables is usually the source of results for the other potential theories.

4) \mathcal{F} -pluriharmonics for subequations $\mathcal{F} \subset \mathcal{S}(n)$: These functions are the analogue of the real part of holomorphic functions. They are defined by requiring that the second derivative belongs to the largest linear subspace of \mathcal{F} , referred to as the *edge* of \mathcal{F} (see [44]). In particular, in Theorem 9.3 of [44] conditions on \mathcal{F} are found that ensure that the family of functions that can be written locally as the maximum of a finite number of \mathcal{F} -pluriharmonics suffices for solving the \mathcal{F} -Dirichlet problem via the Perron process.

5) Removable singularities for subequations \mathcal{F} on manifolds: Pluri-potential methods for proving removable singularity theorems in several complex variables can be extended to \mathcal{F} -pluripotential theory and used to prove

Removable singularity theorems in \mathcal{F} -potential theory.

See [32] for details.

6) Tangents to subharmonics: Kieselmann's theory of tangents to plurisubharmonic functions in complex analysis extends to \mathcal{F} -plurisubharmonic

functions if \mathcal{F} is in a broad family of convex cones. The tangents to an \mathcal{F} -plurisubharmonic function u can be used to study the singularities of u . See [42] and [40] for details. It is shown that

Tangents always exist and are maximal functions.

Maximal functions can be thought of as \mathcal{F} -subharmonic functions with certain singularities allowed. The strongest form of uniqueness of tangents is when *strong uniqueness* holds; that is, the tangent of an \mathcal{F} -plurisubharmonic function equals the density times the Riesz kernel. It is shown that, except for \mathcal{P} (where it is false)

Strong uniqueness of tangents holds if \mathcal{F} is $\mathcal{O}(n)$ invariant.

This strong uniqueness fails in all three of the basic cases $\mathcal{F} = \mathcal{P}$, $\mathcal{F} = \mathcal{P}_{\mathbb{C}}$ (the case studied by Kieselman) and $\mathcal{F} = \mathcal{P}_{\mathbb{H}}$ (the quaternionic case).

7) A Bombieri-Hörmander-Siu type structure theorem: The result for the sets of high density for a plurisubharmonic function in complex analysis has a weakened version which extends to \mathcal{F} -subharmonic functions for many convex $\mathcal{F} \subset \mathcal{S}(n)$, concluding that

Strong uniqueness of tangents implies that sets of high density are discrete.

See section 14 of [42] for details.

Recently, Chu [10] dramatically improved this result by showing that the singular set of an \mathcal{F} -subharmonic function stratifies, and proving each stratum is a rectifiable set.

1.3. The Dirichlet problem

Much can be said about the interplay between potential theory and operator theory by studying the Dirichlet problem. We will focus here on the Euclidian (coordinate chart) setting. We assume that X is an open subset in \mathbb{R}^n and consider bounded domains $\Omega \subset\subset X$ with smooth (i.e. C^2) boundaries and boundary data functions $\varphi \in C(\partial\Omega)$. One can state the standard Dirichlet problem in a vague form as:

(DP) – Vague Formulation: Find a function $h \in C(\overline{\Omega})$ which satisfies:

- 1) h is a *solution* on Ω , and
- 2) $h|_{\partial\Omega} = \varphi$.

In order to be precise as to the meaning of solution in 1), one needs to start with either a subequation constraint set $\mathcal{F} \subset \mathcal{J}^2(X)$ or an operator F on the 2-jets $\mathcal{J}^2(X)$. We want to make both choices and then bring them together.

The Dirichlet problem involves three basic questions: **uniqueness**, **existence** and **regularity**. Uniqueness follows from the *comparison principle*

$$(1.8) \quad (\textit{Comparison}) \quad u \leq w \text{ on } \partial\Omega \Rightarrow u \leq w \text{ on } \Omega,$$

for every pair of subharmonics/superharmonics for \mathcal{F} (or for every pair of subsolutions/supersolutions to the equation $F(\mathcal{J}^2u) = 0$). Existence is established by *Perron's method*. The candidate solution is defined pointwise as the upper envelope

$$(1.9) \quad u(x) := \sup_{w \in \mathfrak{F}} w(x), \quad x \in \Omega,$$

of the Perron family \mathfrak{F} of subsolutions w with $w|_{\partial\Omega} \leq \varphi$. For comparison, roughly speaking, the size, but not the shape of the domain Ω can be of importance. By contrast, for existence one has a dichotomy between subequations \mathcal{F} where existence holds for all domains (with $\partial\Omega$ smooth), and subequations \mathcal{F} with an interesting distinguished boundary geometry of \mathcal{F} -*pseudoconvexity* required for existence (see Subsection 1.4 below). It is important to have a condition on the boundary $\partial\Omega$ which is a local (geometrical) requirement. Finally, the vast and important regularity question will, in essence, not be treated here.

We now begin to describe the main ingredients in the two approaches, potential theory and operator theory. The potential theoretic formulation starts with a constraint set $\mathcal{F} \subset \mathcal{J}^2(X)$, while the operator theoretic formulation starts with an operator F whose domain is a subset $\mathcal{G} \subset \mathcal{J}^2(X)$ ($\mathcal{G} \equiv \mathcal{J}^2(X)$ is allowed and, in fact, is frequently required in the literature). Additional conditions must be imposed in either case.

(DP) – Potential Theoretic Formulation: Find $h \in C(\bar{\Omega})$ which satisfies:

- 1a) h is \mathcal{F} -subharmonic on Ω (i.e. $J_x^{2,+}h \subset \mathcal{F}_x$ for each $x \in \Omega$).
- 1b) $-h$ is $\tilde{\mathcal{F}}$ -subharmonic on Ω (i.e. $J_x^{2,+}(-h) \subset \tilde{\mathcal{F}}_x$ for each $x \in \Omega$). Equivalently, we will say that h is \mathcal{F} -superharmonic on Ω .
- 2) $h|_{\partial\Omega} = \varphi$.

Here and below, Int denotes the interior of a set, $\mathcal{F}_x := \{J \in \mathcal{J}^2 : (x, J) \in \mathcal{F}\}$ is the *fiber of \mathcal{F} over x* , and $\tilde{\mathcal{F}}$ is the *dual* of \mathcal{F} (see Definition 2.2). The space $J_x^{2,+}h$ of *upper test jets of h at x* are defined in (2.6) (also see Definition 2.4). One can easily see that the definition 1b) is equivalent to saying that the lower test jets of h satisfy $J_x^{2,-}h \subset (\text{Int } \mathcal{F}_x)^c$, for each $x \in \Omega$. Additional conditions placed on the constraint set \mathcal{F} are made precise in Definition 2.1. They are summarized by saying that \mathcal{F} is a **subequation**.

The operator theoretic formulation of (DP), although standard, requires some explanation including the imposition of some conditions on the operator F and its domain \mathcal{G} , which we give now for the sake of completeness.

Definition 1.1 (Proper elliptic operators). An operator $F \in C(\mathcal{G})$ where either

$$\mathcal{G} = \mathcal{J}^2(X) \quad (\text{unconstrained case})$$

or

$$\mathcal{G} \subsetneq \mathcal{J}^2(X) \text{ is a subequation constraint set} \quad (\text{constrained case}).$$

is said to be *proper elliptic* if for each $x \in X$ and each $(r, p, A) \in \mathcal{G}_x$ one has

$$(1.10) \quad F(x, r, p, A) \leq F(x, r + s, p, A + P) \quad \forall s \leq 0 \text{ in } \mathbb{R}, \forall P \geq 0 \text{ in } \mathcal{S}(n).$$

The pair (F, \mathcal{G}) will be called a *proper elliptic¹ (operator-subequation) pair*.

The minimal monotonicity (1.10) of the operator F parallels the minimal monotonicity properties (P) and (N) for subequations \mathcal{F} . It is needed for *coherence* and eliminates obvious counterexamples for comparison. This is explained for subequations after Definition 2.4. A given operator F must often be restricted to a suitable background constraint domain $\mathcal{G} \subset \mathcal{J}^2(X)$ in order to have this minimal monotonicity (the constrained case). The historical example clarifying the need for imposing a constraint is the Monge-Ampère operator

$$(1.11) \quad F(D^2u) = \det(D^2u),$$

where one restricts the operator’s domain to be the convexity subequation $\mathcal{G} = \mathcal{P} := \{A \in \mathcal{S}(n) : A \geq 0\}$. The scope of the constrained case is perhaps

¹Such operators are often referred to as *proper operators* (starting from [15]). We prefer to maintain the term “elliptic” to emphasise the importance of the *degenerate ellipticity* (\mathcal{P} -monotonicity in A) in the theory.

best illustrated by the more general *Gårding-Dirichlet operators*. See Example 1.12, with (1.11) the fundamental case. These polynomial operators F of degree m have an ordered sequence of *Gårding eigenvalues* $\Lambda_1(A) \leq \dots \leq \Lambda_m(A)$ which determine **branches** of the equation $F(J^2u) = 0$. The notion of branches well illustrates the interplay between potential theory and operator theory and will be discussed in Example 1.15.

The unconstrained case, in which F is proper elliptic on all of $\mathcal{J}^2(X)$ is the case usually treated in the literature and is perhaps best illustrated by the canonical operators mentioned above.

We now recall the precise notion of solutions in the operator theoretic formulation of the Dirichlet problem. The definitions again make use of *upper/lower test jets*.

Definition 1.2 (Admissible viscosity solutions). Given $F \in C(\mathcal{G})$ with $\mathcal{G} \subset \mathcal{J}^2(X)$ a subequation on an open subset $X \subset \mathbb{R}^n$:

- (a) a function $u \in \text{USC}(X)$ is said to be an (\mathcal{G} -admissible) *viscosity subsolution of $F(J^2u) = 0$ on X* if for every $x \in X$ one has

$$(1.12) \quad J \in J_x^{2,+}u \Rightarrow J \in \mathcal{G}_x \quad \text{and} \quad F(x, J) \geq 0;$$

- (b) a function $u \in \text{LSC}(\Omega)$ is said to be an (\mathcal{G} -admissible) *viscosity supersolution of $F(J^2u) = 0$ on X* if for every $x \in X$ one has

$$(1.13) \quad J \in J_x^{2,-}u \Rightarrow \text{either } [J \in \mathcal{G}_x \text{ and } F(x, J) \leq 0] \text{ or } J \notin \mathcal{G}_x.$$

A function $u \in C(\Omega)$ is an (\mathcal{G} -admissible viscosity) *solution of $F(J^2u) = 0$ on X* if both (a) and (b) hold.

In the unconstrained case where $\mathcal{G} \equiv \mathcal{J}^2(X)$, the definitions are standard. In the constrained case where $\mathcal{G} \subsetneq \mathcal{J}^2(X)$, the definitions give a systematic way of doing of what is sometimes done in an ad-hoc way (see [48] for operators of Monge-Ampère type and [55] for prescribed curvature equations). Note that (1.12) says that the subsolution u is also \mathcal{G} -subharmonic and that (1.13) is equivalent to saying that $F(x, J) \leq 0$ for the lower test jets which lie in the constraint \mathcal{G}_x .

(DP)' – Operator Theoretic Formulation: Find $h \in C(\overline{\Omega})$ which satisfies:

- 1a)' h is a (\mathcal{G} -admissible) *subsolution of $F(J^2h) = 0$ on Ω* (Definition 1.2(a)).
- 1b)' h is a (\mathcal{G} -admissible) *supersolution of $F(J^2h) = 0$ on Ω* (Definition 1.2(b)).
- 2) $h|_{\partial\Omega} = \varphi$.

We now discuss the equivalence of the potential theoretic and operator theoretic formulations of the Dirichlet problem; that is, the equivalence of (DP) for a given subequation \mathcal{F} and (DP)' for a given (proper elliptic) operator-subequation pair (F, \mathcal{G}) . By the definitions, the equivalence of 1a) and 1a)' is the same as the following equivalence: for each $x \in \Omega$ one has

$$J_x^{2,+}h \subset \mathcal{F}_x \Leftrightarrow \text{both } J_x^{2,+}h \subset \mathcal{G}_x \text{ and } F(x, J) \geq 0 \text{ for each } J \in J_x^{2,+}h.$$

This holds if and only if one has the **correspondence relation**

$$(1.14) \quad \mathcal{F} = \{(x, J) \in \mathcal{G} : F(x, J) \geq 0\}.$$

In addition, the equivalence of 1b) and 1b)' is the same as the following equivalence: for each $x \in \Omega$ one has

$$(1.15) \quad J_x^{2,+}(-h) \subset \tilde{\mathcal{F}}_x \Leftrightarrow J \notin \mathcal{G}_x \text{ or } [J \in \mathcal{G}_x \text{ and } F(x, J) \leq 0], \forall J \in J_x^{2,-}h.$$

Using duality (2.2) and $J_x^{2,+}(-h) = -J_x^{2,-}h$ one can see that that the equivalence (1.15) holds if and only if one has **compatibility**

$$(1.16) \quad \text{Int } \mathcal{F} = \{(x, J) \in \mathcal{G} : F(x, J) > 0\},$$

which for subequations \mathcal{F} defined by (1.14) is equivalent to

$$(1.17) \quad \partial\mathcal{F} = \{(x, J) \in \mathcal{F} : F(x, J) = 0\}.$$

The pair of equivalences 1a) \Leftrightarrow 1a)' and 1b) \Leftrightarrow 1b)' is referred to as the **correspondence principle** and will be discussed further in Section 4. These considerations can be summarized in the following result.

Theorem 1.3 (Correspondence Principle). *Suppose that $F \in C(\mathcal{G})$ is proper elliptic and \mathcal{F} , defined by the correspondence relation (1.14), is a subequation. If compatibility (1.16) is satisfied, then $h \in C(\bar{\Omega})$ satisfies the correspondence principle: 1a) \Leftrightarrow 1a)' and 1b) \Leftrightarrow 1b)'. In conclusion, the two formulations (DP) and (DP)' are equivalent.*

Remark 1.4 (On compatibility). Given one of either a proper elliptic pair (F, \mathcal{G}) or a subequation \mathcal{F} , finding the other so that both the correspondence relation (1.14) and compatibility (1.16) hold can be impossible, easy or in between requiring some work. For example, given any subequation \mathcal{F} the pair $(F, \mathcal{J}^2(X))$ with F the signed distance operator (1.4) will do. Other natural choices of $(F, \mathcal{J}^2(X))$ with \mathcal{F} given, which require some additional work, are

the canonical operators introduced in [43]. In the other direction, in Subsection 1.5 we will present various examples of determining the subequation \mathcal{F} given a proper elliptic pair (F, \mathcal{G}) . In particular, finding \mathcal{F} is easy in Examples 1.6 and 1.7, requires some work in Examples 1.8, 1.9, 1.12 and 1.10, and is impossible in Example 1.18.

1.4. Boundary pseudoconvexity

The potential theoretic approach to the Dirichlet problem (DP) naturally leads to an appropriate notion of pseudoconvexity for $\partial\Omega$ (smooth) required for existence. This is perhaps best illustrated by focusing on the case of a constant coefficient pure second order subequation $\mathcal{F} \subset \mathcal{S}(n)$ which is a cone. The definition, with roots in [7], is given in [24, section 5] with several equivalent formulations.

Definition 1.5 (Boundary pseudoconvexity). A smooth boundary $\partial\Omega$ is said to be *strictly \mathcal{F} -pseudoconvex at $x \in \partial\Omega$* if

$$(1.18) \quad \exists t_0 > 0 \quad \text{such that} \quad A_x + tP_{e(x)} \in \text{Int } \mathcal{F}, \quad \forall t \geq t_0,$$

where A_x denotes the second fundamental form of $\partial\Omega$ at x with respect to the inward pointing unit normal $e(x)$ and $P_{e(x)}$ is orthogonal projection onto the normal line through $e(x)$ (the eigenvalues of A_x are the principal curvatures of $\partial\Omega$ at x).

These (cone) subequations \mathcal{F} divide into two kinds, those with and those without a boundary geometry. By those without a boundary geometry we mean that all boundaries $\partial\Omega$ are strictly \mathcal{F} -pseudoconvex at all points. This is equivalent to requiring that

$$(1.19) \quad \forall A \in \mathcal{S}(n), \forall e \in S^{n-1} \quad \exists t_0 > 0 \quad \text{s.t.} \quad A_x + tP_e \in \text{Int } \mathcal{F}, \quad \forall t \geq t_0,$$

Taking $A = 0$ implies

$$(1.20) \quad P_e \in \text{Int } \mathcal{F} \quad \text{for every } e \in S^{n-1}.$$

Conversely, if (1.20) holds then $P_e + \varepsilon A \in \text{Int } \mathcal{F}$ for $\varepsilon > 0$ small, which is equivalent to (1.19). This proves that \mathcal{F} has no boundary (geometric) restriction for existence for the (DP) if and only if \mathcal{F} is strictly elliptic, since (1.20) is one of the ways of defining strict ellipticity. This proves that

the strictly elliptic potential theories are exactly the ones without a boundary pseudoconvexity geometry.

It is easy to see that none of the geometric potential theories $\mathcal{F} = G^\circ$, G a closed subset of $\mathbb{G}(n, \mathbb{R}^n)$, satisfy (1.20); i.e. none are strictly elliptic, so each has a boundary geometry. This geometry has the following nice description

$\partial\Omega$ is strictly $\mathcal{F} = G^\circ$ -pseudoconvex at $x \in \partial\Omega \iff$ the restriction of the second fundamental form A_x to any k -plane $W \in G$ which is also tangential, i.e. $W \subset T_x\partial\Omega$, has strictly positive trace.

1.5. Some examples of PDEs

The potential theory approach to treating nonlinear PDEs is well illustrated by many examples of operators (and classes of operators). We mention a few here.

Example 1.6 (Perturbed Monge-Ampère). With fixed $M \in C(\Omega, \mathcal{S}(n))$ and $f \in C(\Omega)$ non-negative, consider

$$(1.21) \quad \det(D^2u + M(x)) = f(x), \quad x \in \Omega \subset \subset \mathbb{R}^n.$$

This is an important test example of Krylov [51, Example 8.2.4] for probabilistic and analytic methods. It is also noteworthy because it fails to satisfy the standard viscosity structural conditions for comparison as given in Crandall-Ishii-Lions [15, condition (3.14)] unless M is the square of a Lipschitz continuous matrix valued function. In [12], comparison is proved for general continuous M (along with the existence of a unique continuous solution of the Dirichlet problem on strictly convex domains). The potential theoretic proof, makes use of the compatible subequation whose fibers are defined by

$$\mathcal{F}_x := \{A \in \mathcal{S}(n) : A + M(x) \geq 0 \text{ and } F(x, A) := \det(A + M(x)) - f(x) \geq 0\}.$$

This was done with the introduction and application of the notion of (Hausdorff) continuity of the fiber map

$$\Theta : \Omega \rightarrow \wp(\mathcal{S}(n)) \text{ defined by } \Theta(x) := \mathcal{F}_x, \quad \forall x \in \Omega.$$

This is a representative example of the “constrained case” in which operators F come with domains; that is, F must be restricted to \mathcal{G} defined by its fibers $\mathcal{G}_x := \{A \in \mathcal{S}(n) : A + M(x) \geq 0\}$.

Example 1.7 (Special Lagrangian potential equation). With *phase function* $\theta \in C(\Omega, I)$ where $I = (-n\pi/2, n\pi/2)$ consider

$$(1.22) \quad G(D^2u) := \sum_{k=1}^n \arctan(\lambda_k(D^2u)) = \theta(x), \quad x \in \Omega \subset \subset \mathbb{R}^n.$$

The geometric interpretation of this equation is that the graph of the gradient of u will have Lagrangian phase θ (see [21]). Comparison for constant phases, as well as existence/uniqueness for the Dirichlet problem via Perron’s method, was proven in [24]. For non-constant phases, comparison is difficult and only recently completely settled. The operator G is particularly difficult to analyze if the non-constant phase θ assumes a *special phase value* $\theta_k = (n - 2k)\pi/2, k = 1, \dots, n - 1$. Comparison was proven in [17] if h has range in the first/last intervals I_k determined by θ_k . This is the “relatively easy” case where G is concave/convex. Comparison was proven in [13] for phases taking values in any phase interval

$$(1.23) \quad I_k = (\theta_{k-1}, \theta_k), \quad k = 1, \dots, n.$$

There the key was to establish the fiber regularity of the fiber map

$$\Theta(x) := \{A \in \mathcal{S}(n) : F(x, A) := G(A) - h(x) \geq 0\},$$

which is false across the special phase values. Recently, counterexamples to comparison have been given in [6] when the phase takes on a critical value. Combining comparison with the appropriate pseudoconvexity assumption on Ω yields existence/uniqueness for the Dirichlet problem for phases taking values in the intervals (1.23), as shown in [46] (including a study of the needed pseudoconvexity). This leaves open the interesting and unresolved question as to whether uniqueness holds for the Dirichlet problem if the phase takes on a critical value. Example 1.7 is a representative example of the unconstrained case (also pure second order) where the operator G is increasing on all of $\mathcal{S}(n)$.

Example 1.8 (Eigenvalue equation for k -Hessian operators). With $k = 1, \dots, n$ and $\mu \in \mathbb{R}$ fixed, consider

$$(1.24) \quad S_k(D^2u) + \mu u|u|^{k-1} = 0, \quad x \in \Omega \subset \subset \mathbb{R}^n,$$

where for $A \in \mathcal{S}(n)$ the k -Hessian operator is defined by

$$S_k(A) := \sigma_k(\boldsymbol{\lambda}(A)) = \sigma_k(\lambda_1(A), \dots, \lambda_n(A)) = \sum_{1 \leq i_1 < \dots < i_k \leq n} \lambda_{i_1}(A) \cdots \lambda_{i_k}(A).$$

Since the equation is k -homogeneous, one can search for eigen-directions (rays) u that solve (1.24) for an eigenvalue μ . The operator S_k is degenerate elliptic (increasing in A) when restricted to the closed cone

$$\Sigma_k := \{A \in \mathcal{S}(A) : \boldsymbol{\lambda}(A) \in \bar{\Gamma}_k\} \text{ with } \Gamma_k := \{\boldsymbol{\lambda} \in \mathbb{R}^n : \sigma_j(\boldsymbol{\lambda}) > 0, j = 1, \dots, k\},$$

the (closed) *Gårding cone* associated to σ_k (k th elementary symmetric polynomial). Notice that S_k interpolates between $S_1(D^2u) = \text{tr}(D^2u) = \Delta u$ and $S_n(D^2u) = \det(D^2u)$. One uses Σ_k as a background (subequation cone) constraint set; that is, one looks for k -convex subharmonics and uses k -convex lower test functions for supersolutions. The interesting case concerns $u \leq 0$ and $\mu > 0$ where the equation has the “wrong” monotonicity in u . In [5], a maximum principle characterization of a *generalized principal eigenvalue* in the sense of Berestycki-Nirenberg-Varadhan [3] is proven as well as the existence of a corresponding eigenfunction vanishing on the boundary. An important step in the proof is to prove an a priori Hölder estimate, which is needed for compactness in an iterative scheme for the construction of the eigenfunction. The proof shows that the theory of admissibility constraints extends in a natural way the technique pioneered by Ishii-Lions [48] in the unconstrained case.

Example 1.9 (Hyperbolic affine sphere equation). With $X \subset \mathbb{R}^n$ open and $f \in C(X)$ non-negative consider the following equation on X

$$(1.25) \quad [-u]^{n+2} \det(D^2u) = f.$$

The geometric interpretation of the equation emerges by setting $h := -f$ so that the equation becomes

$$(1.26) \quad \det(D^2u) = (h/u)^{n+2},$$

which for u convex and neqative describes the graphing function of a *hyperbolic affine sphere with (constant) mean curvature $h \leq 0$* as discussed in Cheng-Yau [9]. Comparison for the equation (1.25) was established in [13]. This is another representative (gradient-free) example of the constrained case, where (F, \mathcal{G}) with

$$F(x, r, A) := (-r)^{n+2} \det A - f(x) \quad \text{and} \quad \mathcal{G} = \mathcal{Q} := \mathcal{N} \times \mathcal{P}.$$

Example 1.10 (Optimal transport equations). With $X \subset \mathbb{R}^n$ open, $f \in C(X)$ non-negative and $g \in C(\mathbb{R}^n)$ non-negative, consider the following equation on X

$$(1.27) \quad g(Du) \det(D^2u) = f.$$

The functions f and g represent the source and target densities respectively which should have the same mass (L^1 -norm) (see [16] and [56] for more details).

Comparison has been shown for constant target densities g in [13]. For non-constant g , one requires the additional monotonicity property of **directionality**; that is, there exists a closed convex cone $\mathcal{D} \subset \mathbb{R}^n$ with non-empty interior and vertex at the origin for which

$$(1.28) \quad g(p + q) \geq g(p) \text{ for each } p, q \in \mathcal{D}.$$

Examples of g with directionality include

$$g(p) = -p_n \text{ with } \mathcal{D} = \{(p', p_n) \in \mathbb{R}^n : p_n \geq 0\}$$

and for $k \in \{1, \dots, n\}$

$$g(p) = \prod_{j=1}^k p_j \text{ with } D = \{(p_1, \dots, p_n) \in \mathbb{R}^n : p_j \geq 0 \text{ for each } j = 1, \dots, k\}$$

For target densities g with directionality, comparison has been shown for constant source densities f in [11] and for non-constant f in [14].

Remark 1.11. Examples 1.9 and 1.10 have a product structure

$$F(x, u, Du, D^2u) = g(x, u)h(x, Du)G(x, D^2u) - f(x).$$

This structure helps with the correspondence principle. One illustration of this is provided by considering the following pair of constant coefficient gradient-free operators

$$F(r, A) := -r \det(A) \quad \text{and} \quad G(r, A) := -r + \det(A).$$

The first operator is proper elliptic when restricted to the subequation $\mathcal{Q} = \mathcal{N} \times \mathcal{P}$ and with $\mathcal{F} := \{(r, A) \in \mathcal{Q} : F(r, A) \geq 0\}$ one has the compatibility

$$\partial\mathcal{F} = \{(r, A) \in \mathcal{F} : F(r, A) = 0\}$$

and hence the correspondence principle. On the other hand, while G is also proper elliptic when restricted to \mathcal{Q} (or even $\mathbb{R} \times \mathcal{P}$), the boundary of

$$\mathcal{G} := \{(r, A) \in \mathcal{Q} : G(r, A) \geq 0\}$$

includes $\mathcal{N} \times \{0\}$, so that all negative C^2 affine functions will be \mathcal{G} -harmonic but the operator G is not zero on them. Thus the correspondence principle fails here.

Next we discuss perhaps what is perhaps the most interesting and important class of examples. They illustrate why the constrained case is required.

Example 1.12 (Gårding-Dirichlet operators). These nonlinear operators are obtained from Gårding’s beautiful theory of hyperbolic polynomials [19]. We briefly give the precise definition and enumerate a few examples. See [25] and [24], [26], [30] and [11] for an extensive discussion.

Definition 1.13 (Hyperbolic polynomials). A homogeneous real polynomial F of degree m on $\mathcal{S}(n)$ is I -hyperbolic if $F(I) > 0$ and for all $A \in \mathcal{S}(n)$ the one variable polynomial $F(sI + A)$ has all m roots real.

In keeping with the example $F(A) := \det A$, it is useful to focus on the negatives of the roots of $F(sI + A)$, which are called *Gårding I -eigenvalues* and are denoted by $\Lambda_1(A), \dots, \Lambda_m(A)$. Hence

$$(1.29) \quad F(sI + A) = F(A) \prod_{j=1}^m (s + \Lambda_j(A)) \quad \text{and} \quad F(I) = F(I) \prod_{j=1}^m \Lambda_j(A).$$

The *open Gårding cone*

$$(1.30) \quad \Gamma := \{A \in \mathcal{S}(n) : \Lambda_j(A) > 0, j = 1, \dots, m\}$$

is a convex cone, which, along with F , has many nice properties.

Definition 1.14 (Gårding-Dirichlet operator²). An I -hyperbolic polynomial operator F of degree m on $\mathcal{S}(n)$ is called a *Gårding-Dirichlet operator* if $\mathcal{P} \subset \bar{\Gamma}$; that is, if

$$(1.31) \quad A \geq 0 \quad \Rightarrow \quad \Lambda_j(A) \geq 0, \quad j = 1, \dots, m.$$

In this case

$$(1.32) \quad (F, \mathcal{F} := \bar{\Gamma}) \text{ is a compatible operator-subequation pair.}$$

Also note that by (1.32) $\Lambda_{\min}(A)$ is the canonical operator for the Gårding subequation $\mathcal{F} := \bar{\Gamma}$ since

$$\Lambda_j(A + tI) = \Lambda_j(A) + t, \quad j = 1, \dots, m.$$

²Perhaps they should be called *Gårding-Monge-Ampère operators* instead.

Here are some important examples. Let $\lambda_1(A), \dots, \lambda_n(A)$ denote the standard eigenvalues of $A \in \mathcal{S}(n)$. Of course, the Monge-Ampère operator

$$F(A) = \det A = \prod_{j=1}^n \lambda_j(A)$$

is the prototype. We leave it to the reader to see that these examples are polynomials. Perhaps the most interesting are those for which $\mathcal{F} := \bar{\Gamma}$ is geometrically defined; i.e. the cone subequation $\mathcal{F} := \bar{\Gamma}$ equals the polar G° of a closed subset G of one the Grassmanians. Two new examples of this special nature are as follows.

1. The p -fold sum operator ([24, p. 39] and [29, Proposition 7.11]):

This is the operator

$$(1.33) \quad F(A) := \prod_{i_1 < \dots < i_p} (\lambda_{i_1}(A) + \dots + \lambda_{i_p}(A)), \quad \text{where } p = 1, \dots, n.$$

The degree of F is $\binom{n}{p}$ and the closed Gårding cone is $\bar{\Gamma} = \mathbb{G}(p, \mathbb{R}^n)^\circ$. The canonical operator for $\bar{\Gamma}$ is the sum of the first (smallest) p standard eigenvalues; i.e. the p -th truncated Laplacian (see [24] and [4]).

2. The Lagrangian Monge-Ampère operator ([41, section 5]):

This is the operator

$$(1.34) \quad F(A) := \prod \left(\frac{1}{2} \operatorname{tr} A \pm \mu_1 \pm \dots \pm \mu_n \right).$$

Here $A \in \mathcal{S}(2n)$ is a real symmetric form on $\mathbb{R}^{2n} = \mathbb{C}^n$, and $\pm\mu_1 \dots \pm\mu_n$ are the eigenvalues of the skew Hermitian part of A . The 2^n Gårding eigenvalues are the factors $\frac{1}{2} \operatorname{tr} A \pm \mu_1 \pm \dots \pm \mu_n$. The Gårding subequation $\bar{\Gamma}$ is geometric. It is the polar of $G := \text{LAG} \subset \mathbb{G}_{\mathbb{R}}(n, \mathbb{C}^n)$, the set of Lagrangian n -planes. The plurisubharmonic functions; i.e. the $\bar{\Gamma}$ -subharmonic functions are those upper semicontinuous functions that restrict to be Δ -subharmonic on Lagrangian affine planes in \mathbb{C}^n .

A classical example, which is not geometric for $k \neq 1, n$ is the k -Hessian operator $S_k(D^2u)$ discussed in Example 1.8. Two more new non geometric examples, which have similarities with one another, are the following.

3. The δ -uniformly elliptic operator ([38, Appendix B]): This is the operator

$$(1.35) \quad F(A) := \det(A + \delta(\operatorname{tr} A)I) \text{ with } \delta > 0,$$

The Gårding eigenvalues of F are

$$\Lambda_j(A) = \lambda_j(A) + \delta \operatorname{tr}(A), \quad j = 1, \dots, n.$$

Hence the canonical operator for the Gårding subequation $\bar{\Gamma}$ is

$$\Lambda_{\min}(A) = \lambda_{\min}(A) + \delta \operatorname{tr} A.$$

4. The Pucci-Gårding-Monge-Ampère operator ([30, section 4.5], [38, Appendix B], and in particular [43, Example 6.10]): Fix $0 < \lambda < \Lambda$ and consider the the “cube” in eigenvalue space

$$\mathcal{C}_{\lambda,\Lambda} := \{A \in \mathcal{S}(n) : \lambda I \leq A \leq \Lambda I\} \subset \mathcal{P} \subset \mathcal{S}(n).$$

Its polar $\mathcal{P}_{\lambda,\Lambda}^- := \mathcal{C}_{\lambda,\Lambda}^\circ$ contains $\mathcal{P}^\circ = \mathcal{P}$ and hence is a subequation called the *Pucci* or *Pucci-Gårding cone*. The cone on $\mathcal{C}_{\lambda,\Lambda}$, denoted by $\operatorname{Cone}(\mathcal{C}_{\lambda,\Lambda})$, has a finite set of extreme rays through a subset S of the vertices of $\mathcal{C}_{\lambda,\Lambda}$. The *Pucci-Gårding-Monge-Ampère operator* $F_{\lambda,\Lambda}$ is defined to be the product of the linear functionals in the set S . The factors are the Gårding I -eigenvalues of $F_{\lambda,\Lambda}$. The closed Gårding cone is the Pucci cone $\mathcal{P}_{\lambda,\Lambda}^-$. Its canonical operator is the minimal eigenvalue, which is easily seen to be

$$\Lambda_{\min}(A) = \lambda \operatorname{tr} A^+ + \Lambda \operatorname{tr} A^- \geq 0,$$

where $A = A^+ + A^-$ is the decomposition of A into positive and negative parts, and hence

$$\mathcal{P}_{\lambda,\Lambda}^- := \{A \in \mathcal{S}(n) : \lambda \operatorname{tr} A^+ + \Lambda \operatorname{tr} A^- \geq 0\}.$$

Note that the Pucci-Gårding operator $F_{\lambda,\Lambda}$ has degree $|S|$.

The operator S_k and the operators 1., 3., and 4. above which involve the real eigenvalues of $A \in \mathcal{S}(n)$ have complex and quaternionic analogues that are also Gårding-Dirichlet operators. See [25, section 5], [24, section 10] and [26, section 15] for more details.

The next family of examples provides a good illustration of the interplay between potential theory and operator theory.

Example 1.15 (Branches). The potential theory/subequation approach provides a direct way of extending the Dirichlet problem (DP) for the Monge-Ampère operator to the other *branches* Λ_k of $\det(D^2u) = 0$, where, except for $k = 1$, there is no natural smooth operator F defining the solutions (or Λ_k -subharmonics). The branch $\Lambda_k \subset \mathcal{S}(n)$ for $k = 1, \dots, n$ is the subequation defined by

$$(1.36) \quad \Lambda_k := \{A \in \mathcal{S}(n) : \lambda_k(A) \geq 0\},$$

where $\lambda_1(A) \leq \dots \leq \lambda_n(A)$ are the ordered eigenvalues of $A \in \mathcal{S}(n)$. Despite the fact that $\det(D^2u)$ is not a proper elliptic operator on Λ_k , the (DP) for Λ_k -harmonics is meaningful. Existence and uniqueness for all boundary functions $\varphi \in C(\partial\Omega)$ under the appropriate geometrical conditions on $\partial\Omega$ was established in [24]. This theorem extends to branches of a more general Gårding-Dirichlet operator $F = \mathfrak{g}$ of degree m defined by

$$\Lambda_k^{\mathfrak{g}} := \{A \in \mathcal{S}(n) : \Lambda_k^{\mathfrak{g}}(A) \geq 0\}, \quad k = 1, \dots, m.$$

These branches have a so-called *canonical operator*, which we discuss next. Canonical operators well illustrate the unconstrained case.

Example 1.16 (Canonical operators). For clarity we focus on the pure second order constant coefficient subequation case $\mathcal{F} \subset \mathcal{S}(n)$. See the list of references before the formula (1.4) above for more information. A *canonical operator* for \mathcal{F} is by definition a function $F \in C(\mathcal{S}(n))$ with the following two properties:

$$(1.37) \quad F(A + P) \geq F(A), \quad \forall A \in \mathcal{S}(n) \text{ and } P \geq 0$$

and for some constant $c > 0$

$$(1.38) \quad F(A + tI) = F(A) + ct, \quad \forall A \in \mathcal{S}(n) \text{ and } t \in \mathbb{R}.$$

Proposition 1.17 (Existence and uniqueness of canonical operators). *Given a subequation $\mathcal{F} \subset \mathcal{S}(n)$, there exists a unique canonical operator F with $\mathcal{F} = \{A \in \mathcal{S}(n) : F(A) \geq 0\}$.*

The proof can be summarized succinctly by defining F by requiring

$$A_0 + F(A_0)I \in \partial\mathcal{F} \quad \text{for all } A_0 \perp I \text{ (i.e. } \text{tr } A_0 = 0)$$

and then extending F to all $A = A_0 + tI \in \mathcal{S}(n), A_0 \perp I$ by formula (1.38).

The canonical operator F for some of the examples above are as follows. One has $\lambda_1(A)$ for the convexity subequation \mathcal{P} , $\lambda_1(A) + \dots + \lambda_p(A)$ for the p -fold subequation defined by (1.7) with $G = G(p, \mathbb{R}^n)$, $\frac{1}{2} \operatorname{tr} A - \mu_1 - \dots - \mu_n$ for the Lagrangian subequation defined by (1.7) with $G = \text{LAG}$. As mentioned above, the Gårding subequation $\mathcal{F} = \bar{\Gamma}$ has canonical operator $\Lambda_{\min}(A)$, the minimal Gårding eigenvalue operator. The k -th branch has canonical operator the k -th Gårding eigenvalue operator. The construction of a canonical operator extends to subequations $\mathcal{F} \subset \mathcal{J}^2(X)$ if there is sufficient monotonicity (see section 11.4 of [11]).

We conclude this subsection with one last example. As we have indicated, a general principle is that comparison holds with sufficient monotonicity. With insufficient monotonicity comparison can fail even in the constant coefficient case and even on arbitrarily small balls.

Example 1.18 (Comparison fails). Consider the operators $F, G \in C(\mathbb{R}^n \times \mathcal{S}(n), \mathbb{R})$ defined by

$$F(p, A) := \lambda_{\min}(M(p, A)) \quad \text{and} \quad G(p, A) := \lambda_{\max}(M(p, A))$$

with $M = M_\alpha$ is the $\mathcal{S}(n)$ -valued function defined for $\alpha \in (1, +\infty)$ by

$$M(p, A) := A + |p|^{\frac{\alpha-1}{n}}(P_{p^\perp} + \alpha P_p) \quad \text{if } p \neq 0 \quad \text{and} \quad M(0, A) := A.$$

where for $p \neq 0$, P_p, P_{p^\perp} are the projections onto the subspaces $[p], [p]^\perp$; that is,

$$P_p = \frac{1}{|p|^2} p \otimes p \quad \text{and} \quad P_{p^\perp} = I - P_p.$$

These operators are studied in [11]. Existence for the Dirichlet problem holds on all balls (with continuous Dirichlet data). The comparison principle, maximum principle and uniqueness of solutions fail on all arbitrarily small balls about every point. A partial explanation of these failures is that the *maximal monotonicity cones* for the associated (compatible) subequations \mathcal{F}, \mathcal{G} are $\mathcal{M} := \{0\} \times \mathcal{P} \subset \mathbb{R}^n \times \mathcal{S}(n)$, which have empty interior.

2. Fundamental aspects of nonlinear potential theory

In this section, we give a brief review of some key notions and fundamental results in the theory of \mathcal{F} -subharmonic functions defined by a subequation constraint set \mathcal{F} .

2.1. Subequations, subharmonics and duality

Suppose that X is an open subset of \mathbb{R}^n with 2-jet space denoted by $\mathcal{J}^2(X) = X \times (\mathbb{R} \times \mathbb{R}^n \times \mathcal{S}(n))$. A good definition of a constraint set with a robust potential theory was given in [26] (also for manifolds).

Definition 2.1 (Subequations). A set $\mathcal{F} \subset \mathcal{J}^2(X)$ is called a *subequation* (*constraint set*) if

(P) \mathcal{F} satisfies the *positivity condition* (fiberwise); that is, for each $x \in X$

$$(r, p, A) \in \mathcal{F}_x \Rightarrow (r, p, A + P) \in \mathcal{F}_x, \quad \forall P \geq 0 \text{ in } \mathcal{S}(n).$$

(T) \mathcal{F} satisfies three conditions of *topological stability*:

- (T1) $\mathcal{F} = \overline{\text{Int } \mathcal{F}}$;
- (T2) $\mathcal{F}_x = \overline{\text{Int } (\mathcal{F}_x)}, \quad \forall x \in X$;
- (T3) $(\text{Int } \mathcal{F})_x = \text{Int } (\mathcal{F}_x), \quad \forall x \in X$.

(N) \mathcal{F} satisfies the *negativity condition* (fiberwise); that is, for each $x \in X$

$$(r, p, A) \in \mathcal{F}_x \Rightarrow (r + s, p, A) \in \mathcal{F}_x, \quad \forall s \leq 0 \text{ in } \mathbb{R}.$$

Notice that by property (T1), \mathcal{F} is closed in $\mathcal{J}^2(X)$ and each fiber \mathcal{F}_x is closed in \mathcal{J}^2 by (T2). In addition, the interesting case is when each fiber \mathcal{F}_x is not all of \mathcal{J}^2 , which we almost always assume. Also notice that in the constant coefficient pure second order case $\mathcal{F} \subset \mathcal{S}(n)$, property (N) is automatic and property (T) reduces to (T1) $\mathcal{F} = \overline{\text{Int } \mathcal{F}}$, which is implied by (P) for \mathcal{F} closed. Hence in this case subequations $\mathcal{F} \subset \mathcal{S}(n)$ are closed sets simply satisfying (P).

The conditions (P), (T) and (N) have various (important) implications for the potential theory determined by \mathcal{F} . Some of these will be mentioned below (see the brief discussion following Definition 2.4).

Next is *duality*, a notion first introduced in the pure second order coefficient case in [24].

Definition 2.2 (The dual subequation). For a given subequation $\mathcal{F} \subset \mathcal{J}^2(X)$ the *Dirichlet dual* of \mathcal{F} is the set $\tilde{\mathcal{F}} \subset \mathcal{J}^2(X)$ given by³

$$(2.1) \quad \tilde{\mathcal{F}} := (-\text{Int } \mathcal{F})^c = -(\text{Int } \mathcal{F})^c \quad (\text{relative to } \mathcal{J}^2(X)).$$

³Here and below, c denotes the set theoretic complement of a subset.

With the help of property (T), the dual can be calculated fiberwise

$$(2.2) \quad \tilde{\mathcal{F}}_x := (-\text{Int}(\mathcal{F}_x))^c = -(\text{Int}(\mathcal{F}_x))^c \text{ (relative to } \mathcal{J}^2), \quad \forall x \in X.$$

This is a true duality in the sense that one can show

$$(2.3) \quad \tilde{\tilde{\mathcal{F}}} = \mathcal{F} \quad \text{and} \quad \mathcal{F} \text{ is a subequation} \quad \Rightarrow \quad \tilde{\mathcal{F}} \text{ is a subequation.}$$

Now comes the notion of \mathcal{F} -subharmonicity for a given subequation $\mathcal{F} \subset \mathcal{J}^2(X)$. There are two different natural formulations for differing degrees of regularity. The first is the classical formulation.

Definition 2.3 (Classical or C^2 subharmonics). A function $u \in C^2(X)$ is said to be \mathcal{F} -subharmonic on X if

$$(2.4) \quad J_x^2 u := (u(x), Du(x), D^2u(x)) \in \mathcal{F}_x, \quad \forall x \in X$$

with the accompanying notion of being *strictly* \mathcal{F} -subharmonic if

$$(2.5) \quad J_x^2 u \in \text{Int}(\mathcal{F}_x) = (\text{Int } \mathcal{F})_x, \quad \forall x \in X.$$

For merely upper semicontinuous functions $u \in \text{USC}(X)$ with values in $[-\infty, +\infty)$, one replaces the 2-jet $J_x^2 u$ with the set of C^2 upper test jets

$$(2.6) \quad J_x^{2,+} u := \{J_x^2 \varphi : \varphi \text{ is } C^2 \text{ near } x, u \leq \varphi \text{ near } x \text{ with equality at } x\},$$

thus arriving at the following viscosity formulation.

Definition 2.4 (Semicontinuous subharmonics). A function $u \in \text{USC}(X)$ is said to be \mathcal{F} -subharmonic on X if

$$(2.7) \quad J_x^{2,+} u \subset \mathcal{F}_x, \quad \forall x \in X.$$

We denote by $\mathcal{F}(X)$ the set of all \mathcal{F} -subharmonics on X .

We now recall some of the implications that properties (P), (T) and (N) have on an \mathcal{F} -potential theory. Property (P) is crucial for C^2 -coherence, meaning classical \mathcal{F} -subharmonics are \mathcal{F} -subharmonics in the sense (2.7), since for u which is C^2 near x , one has

$$J_x^{2,+} u = J_x^2 u + (0, 0, \mathcal{P}) \quad \text{where} \quad \mathcal{P} = \{P \in \mathcal{S}(n) : P \geq 0\}.$$

The natural notion of $w \in \text{LSC}(X)$ being \mathcal{F} -superharmonic using lower test jets is

$$(2.8) \quad J_x^{2,-}w \subset (\text{Int}(\mathcal{F}_x))^c, \quad \forall x \in X,$$

which by duality and property (T) is equivalent to $-w \in \text{USC}(X)$ satisfying

$$(2.9) \quad J_x^{2,+}(-w) \subset \tilde{\mathcal{F}}_x, \quad \forall x \in X.$$

That is,

$$(2.10) \quad w \text{ is } \mathcal{F}\text{-superharmonic} \iff -w \text{ is } \tilde{\mathcal{F}}\text{-subharmonic}.$$

Next note that property (T) insures the local existence of strict classical \mathcal{F} -superharmonics at points $x \in X$ for which \mathcal{F}_x is non-empty. One simply takes the quadratic polynomial whose 2-jet at x is $J \in \text{Int}(\mathcal{F}_x)$. Finally, property (N) eliminates obvious counterexamples to comparison. The simplest counterexample is provided by the constraint set $\mathcal{F} \subset \mathcal{J}^2(\mathbb{R})$ in dimension one associated to the equation $u'' - u = 0$.

2.2. Monotonicity

This fundamental notion appears in various guises. It is a useful and unifying concept. One says that a subequation \mathcal{F} is \mathcal{M} -monotone for some subset $\mathcal{M} \subset \mathcal{J}^2(X)$ if

$$(2.11) \quad \mathcal{F}_x + \mathcal{M}_x \subset \mathcal{F}_x \quad \text{for each } x \in X.$$

For simplicity, we will restrict attention to (constant coefficient) *monotonicity cones*; that is, monotonicity sets \mathcal{M} for \mathcal{F} which have constant fibers which are closed cones with vertex at the origin.

First and foremost, the properties (P) and (N) are monotonicity properties. Property (P) for subequations \mathcal{F} corresponds to *degenerate elliptic* operators F and properties (P) and (N) together correspond to *proper elliptic* operators. Note that (P) plus (N) can be expressed as the single monotonicity property

$$(2.12) \quad \mathcal{F}_x + \mathcal{M}_0 \subset \mathcal{F}_x \quad \text{for each } x \in X$$

where

$$(2.13) \quad \mathcal{M}_0 := \mathcal{N} \times \{0\} \times \mathcal{P} \subset \mathcal{J}^2 = \mathbb{R} \times \mathbb{R}^n \times \mathcal{S}(n)$$

with

$$(2.14) \quad \mathcal{N} := \{r \in \mathbb{R} : r \leq 0\} \quad \text{and} \quad \mathcal{P} := \{P \in \mathcal{S}(n) : P \geq 0\}.$$

Hence \mathcal{M}_0 will be referred to as the *minimal monotonicity cone* in \mathcal{J}^2 . However, it is important to remember that $\mathcal{M}_0 \subset \mathcal{J}^2$ is **not** a subequation since it has empty interior so that property (T) fails.

Combined with duality and fiberegularity (defined in Subsection 2.3), one has a very general, flexible and elegant geometrical approach to comparison when a subequation \mathcal{F} admits a constant coefficient monotonicity cone subequation \mathcal{M} . A key ingredient to this approach is the *Subharmonic Addition Theorem*:

$$(2.15) \quad \mathcal{F} + \mathcal{M} \subset \mathcal{F} \quad \Rightarrow \quad \mathcal{F}(X) + \widetilde{\mathcal{F}}(X) \subset \widetilde{\mathcal{M}}(X).$$

This result reduces the comparison principle for \mathcal{F} to the *Zero Maximum Principle* for the constant coefficient dual cone subequation $\widetilde{\mathcal{M}}$; that is, for all $\Omega \subset\subset X$

$$(2.16) \quad (\text{ZMP}) \quad z \leq 0 \text{ on } \partial\Omega \quad \Rightarrow \quad z \leq 0 \text{ on } \partial\Omega,$$

$\forall z \in \text{USC}(\overline{\Omega}) \cap \widetilde{\mathcal{M}}(\Omega)$. This reduction of comparison to (ZMP) will be referred to as the **monotonicity-duality method** and will be discussed in Section 3.

Monotonicity is also used to formulate **reductions** when certain jet variables are “silent” in the subequation constraint \mathcal{F} . for example, one has

$$\begin{aligned} &(\text{pure second order}) \quad \mathcal{F}_x + \mathcal{M}(\mathcal{P}) \subset \mathcal{F}_x : \quad \mathcal{M}(\mathcal{P}) := \mathbb{R} \times \mathbb{R}^n \times \mathcal{P} \\ &(\text{gradient free}) \quad \mathcal{F}_x + \mathcal{M}(\mathcal{N}, \mathcal{P}) \subset \mathcal{F}_x : \quad \mathcal{M}(\mathcal{N}, \mathcal{P}) := \mathcal{N} \times \mathbb{R}^n \times \mathcal{P} \end{aligned}$$

$\mathcal{M}(\mathcal{P})$ and $\mathcal{M}(\mathcal{N}, \mathcal{P})$ are fundamental *constant coefficient (cone) subequations* which can be identified with $\mathcal{P} \subset \mathcal{S}(n)$ and $\mathcal{Q} := \mathcal{N} \times \mathcal{P} \subset \mathbb{R} \times \mathcal{S}(n)$. One can identify \mathcal{F} with subsets of the *reduced jet bundles* $X \times \mathcal{S}(n)$ and $X \times (\mathbb{R} \times \mathcal{S}(n))$, respectively, “forgetting about” the silent jet variables (see Chapter 10 of [11]).

Two important “reduced” examples are worth drawing special attention to.

Example 2.5 (The convexity subequation). The *convexity subequation* is $\mathcal{F} = X \times \mathcal{M}(\mathcal{P})$ and reduces to $X \times \mathcal{P}$ which has constant coefficients (each fiber is \mathcal{P}) and for $u \in \text{USC}(X)$

$$u \text{ is } \mathcal{P}\text{-subharmonic} \quad \Leftrightarrow \quad u \text{ is locally convex}$$

(away from any connected components where $u \equiv -\infty$).

The convexity subequation has its *canonical operator* $F \in C(\mathcal{S}(n), \mathbb{R})$ defined by the minimal eigenvalue $F(A) := \lambda_{\min}(A)$, for which

$$(2.17) \quad \mathcal{P} = \{A \in \mathcal{S}(n) : \lambda_{\min}(A) \geq 0\}.$$

The dual subequation $\tilde{\mathcal{F}}$ has constant fibers given by

$$(2.18) \quad \tilde{\mathcal{P}} = \{A \in \mathcal{S}(n) : \lambda_{\max}(A) \geq 0\}$$

which is the *subaffine subequation*. The set $\tilde{\mathcal{P}}(X)$ of dual subharmonics agrees with $\text{SA}(X)$ the set of *subaffine functions* defined as those functions $u \in \text{USC}(X)$ which satisfy the *subaffine property* (comparison with affine functions): for every $\Omega \subset\subset X$ one has

$$(2.19) \quad u \leq a \text{ on } \partial\Omega \implies u \leq a \text{ on } \Omega, \text{ for every } a \text{ affine.}$$

The fact that $\tilde{\mathcal{P}}(X) = \text{SA}(X)$ is shown in [24]. The subaffine property for u is stronger than the maximum principle for u since constants are affine functions. It has the advantage over the maximum principle of being a local condition on u . This leads to the comparison principle for all pure second order constant coefficient subequations [24] and extends to variable coefficient subequations [13] using a notion of fiberegularity, as will be discussed in Section 3.

Example 2.6 (The convexity-negativity subequation). The constant coefficient gradient-free subequation $\mathcal{F} = X \times \mathcal{M}(\mathcal{N}, \mathcal{P})$ reduces to $X \times \mathcal{Q} \subset X \times (\mathbb{R} \times \mathcal{S}(n))$ whose (constant) fibers are

$$(2.20) \quad \mathcal{Q} = \mathcal{N} \times \mathcal{P} = \{(r, A) \in \mathbb{R} \times \mathcal{S}(n) : r \leq 0 \text{ and } A \geq 0\}.$$

The (reduced) dual subequation has (constant) fibers

$$(2.21) \quad \tilde{\mathcal{Q}} = \{(r, A) \in \mathbb{R} \times \mathcal{S}(n) : r \leq 0 \text{ or } A \in \tilde{\mathcal{P}}\}.$$

The set $\tilde{\mathcal{Q}}(X)$ of dual subharmonics agrees with $\text{SA}^+(X)$, the set of *subaffine plus functions* defined as those functions $u \in \text{USC}(X)$ which satisfy the *subaffine plus property*: for every $\Omega \subset\subset X$ one has

$$(2.22) \quad u \leq a \text{ on } \partial\Omega \implies u \leq a \text{ on } \Omega, \text{ for every } a \text{ affine with } a|_{\bar{\Omega}} \geq 0,$$

from which the (ZMP) for $\tilde{\mathcal{Q}}$ subharmonics follows immediately. The fact that $\tilde{\mathcal{Q}}(X) = \text{SA}^+(X)$ is shown in [11] together with the additional equivalence

$$(2.23) \quad \text{SA}^+(X) := \{u \in \text{USC}(X) : u^+ := \max\{u, 0\} \in \text{SA}(X) = \tilde{\mathcal{P}}(X)\},$$

The validity of the (ZMP) for \mathcal{Q} -subharmonics leads to the comparison principle by the monotonicity-duality method for all gradient free subequations with constant coefficients in [11] and extends to variable coefficient gradient-free subequations in [13], using the notion of fiberegularity.

2.3. Fiberegularity

This fundamental notion can be used to pass from constant coefficient subequations (and operators) to ones with variable coefficients.

Definition 2.7. A subequation $\mathcal{F} \subset \mathcal{J}^2(X)$ is *fiberegular* if the fiber map Θ is (*Hausdorff*) *continuous*; that is, if the set-valued map

$$\Theta : X \rightarrow \mathcal{K}(\mathcal{J}^2) \text{ defined by } \Theta(x) := \mathcal{F}_x, \quad x \in X$$

is continuous when the closed subsets $\mathcal{K}(\mathcal{J}^2)$ of \mathcal{J}^2 are equipped with the *Hausdorff metric*

$$d_{\mathcal{H}}(\Phi, \Psi) := \max \left\{ \sup_{J \in \Phi} \inf_{J' \in \Psi} \|J - J'\|, \sup_{J' \in \Psi} \inf_{J \in \Phi} \|J - J'\| \right\}$$

where

$$\|J\| = \|(r, p, A)\| := \max \left\{ |r|, |p|, \max_{1 \leq k \leq n} |\lambda_k(A)| \right\}$$

is taken as the norm on \mathcal{J}^2 .

This notion was first introduced in [12] in the special case $\mathcal{F} \subset X \times \mathcal{S}(n)$. We will also refer to Θ as a *continuous proper elliptic map* since it takes values in the closed (non-empty and proper) subsets of \mathcal{J}^2 satisfying properties (P) and (N). If \mathcal{F} is \mathcal{M} -monotone for some (constant coefficient) monotonicity cone subequation, fiberegularity has the more useful equivalent formulation: there exists $J_0 \in \text{Int } \mathcal{M}$ such that for each fixed $\Omega \subset\subset X$ and $\eta > 0$ there exists $\delta = \delta(\eta, \Omega)$ such that

$$(2.24) \quad x, y \in \Omega, |x - y| < \delta \implies \Theta(x) + \eta J_0 \subset \Theta(y).$$

Note that property holds for each fixed $J_0 \in \text{Int } \mathcal{M}$ (see [14]) and that in the pure second order and gradient-free cases there is a “canonical” reduced jet $J_0 = I \in \mathcal{S}(n)$ and $J_0 = (-1, I) \in \mathbb{R} \times \mathcal{S}(n)$, respectively. Also note that fiberegularity is uniform on bounded domains as the δ in (2.24) is independent of $x, y \in \Omega$.

Fiberegularity is crucial since it implies the **uniform translation property** for subharmonics: *if $u \in \mathcal{F}(\Omega)$, then there are small C^2 strictly \mathcal{F} -subharmonic perturbations of all small translates of u which belong to $\mathcal{F}(\Omega_\delta)$, where $\Omega_\delta := \{x \in \Omega : d(x, \partial\Omega) > \delta\}$.*

Theorem 2.8 (Uniform translation property for subharmonics). *Suppose that a subequation \mathcal{F} is a fiberegular and \mathcal{M} -monotone on $\Omega \subset\subset \mathbb{R}^n$ for some monotonicity cone subequation \mathcal{M} . Suppose that \mathcal{M} admits a strict approximator; that is, there exists $\psi \in \text{USC}(\overline{\Omega}) \cap C^2(\Omega)$ which is strictly \mathcal{M} -subharmonic on Ω . Given $u \in \mathcal{F}(\Omega)$, for each $\theta > 0$ there exist $\eta = \eta(\psi, \theta) > 0$ and $\delta = \delta(\psi, \theta) > 0$ such that*

$$(2.25) \quad u_{y,\theta} = \tau_y u + \theta\psi \text{ belongs to } \mathcal{F}(\Omega_\delta), \quad \forall y \in B_\delta(0),$$

where $\tau_y u(\cdot) := u(\cdot - y)$.

In the pure second order and gradient-free cases ($\mathcal{F} \subset \Omega \times \mathcal{S}(n)$ and $\mathcal{F} \subset \Omega \times (\mathbb{R} \times \mathcal{S}(n))$), one always has a quadratic strict approximator ψ and the theorem holds for all continuous coefficient \mathcal{F} which are minimally monotone (with $\mathcal{M} = \mathcal{P} \subset \mathcal{S}(n)$ and $\mathcal{M} = \mathcal{Q} = \mathcal{N} \times \mathcal{P} \subset \mathbb{R} \times \mathcal{S}(n)$ respectively) as shown in [12], [13]. The general \mathcal{M} -monotone and fiberegular case is treated in [14]. In this general case, the hypothesis of the existence of a strict approximator ψ creates no additional problems if the objective is to prove comparison. This is because one knows from [11, Theorem 6.2] that the existence of a strict approximator ψ for \mathcal{M} ensures the validity of the (ZMP) for $\widetilde{\mathcal{M}}$, which is needed for our monotonicity duality method. Moreover, the (constant coefficient) monotonicity cone subequations which admit strict approximators are well understood by the study made in [11].

2.4. Subharmonic addition for quasiconvex functions

Many results about \mathcal{F} -subharmonic functions u are more easily proved if one assumes that u is also locally quasiconvex.⁴ Then, one can make use of *quasiconvex approximation* to extend the result to semicontinuous u . Here we discuss some of the main results in this direction. See [53] for an extensive treatment, which borrows heavily from [36] and [37].

⁴We have adopted the term quasiconvex which is consistent with the use of *quasi-plurisubharmonic* function in several complex variables. Quasiconvex functions are sometimes referred to as *semiconvex* functions, although this term is a bit misleading. They are functions whose Hessian (in the viscosity sense) is locally bounded from below.

Definition 2.9. A function $u : C \rightarrow \mathbb{R}$ is λ -quasiconvex on a convex set $C \subset \mathbb{R}^n$ if there exists $\lambda \in \mathbb{R}_+$ such that $u + \frac{\lambda}{2} |\cdot|^2$ is convex on C . A function $u : X \rightarrow \mathbb{R}$ is *locally quasiconvex* on an open set $X \subset \mathbb{R}^n$ if for every $x \in X$, u is λ -quasiconvex on some ball about x for some $\lambda \in \mathbb{R}_+$.

Such functions are twice differentiable for almost every⁵ $x \in X$ by a very easy generalization of Alexandroff’s theorem for convex functions (the addition of a smooth function has no effect on differentiability). This is one of the many properties that quasiconvex functions inherit from convex functions. Quasiconvex functions are used to approximate $u \in \text{USC}(X)$ (bounded from above) by way of the *sup-convolution*, which for each $\varepsilon > 0$ is defined by

$$(2.26) \quad u^\varepsilon(x) := \sup_{y \in X} \left(u(y) - \frac{1}{2\varepsilon} |y - x|^2 \right), \quad x \in X.$$

One has that u^ε is $\frac{1}{\varepsilon}$ -quasiconvex and decreases pointwise to u as $\varepsilon \rightarrow 0^+$. There is an underlying pure second order potential theory for λ -quasiconvex functions on X ; namely with respect to the λ -quasiconvexity subequation

$$(2.27) \quad \mathcal{P}_\lambda := \{A \in \mathcal{S}(n) : A + \lambda I \in \mathcal{P}\}.$$

Two important results follow.

Theorem 2.10 (The Almost Everywhere Theorem). *For locally quasiconvex functions u*

$$J_x^2 u = (u(x), Du(x), D^2 u(x)) \in \mathcal{F}_x \text{ for } \mathcal{L}^n\text{-a.e. } x \in X \iff u \in \mathcal{F}(X).$$

This result is proven [37]. The main point in the proof is to control the measure of the set of *upper contact points* near x if u is quasiconvex. This control comes from either of two results obtained independently by Slodkowski [54] and Jensen [49]. These two measure theoretic results are shown to be equivalent in [36].

Theorem 2.11 (The Subharmonic Addition Theorem: Quasiconvex Version). *Suppose that the subequations \mathcal{F}, \mathcal{G} and \mathcal{H} satisfy*

$$\text{(Jet addition)} \quad \mathcal{F}_x + \mathcal{G}_x \subset \mathcal{H}_x, \text{ for each } x \in X.$$

If $u \in \mathcal{F}(X)$ and $v \in \mathcal{G}(X)$ are locally quasiconvex, then

$$\text{(Subharmonic addition)} \quad u + v \in \mathcal{H}(X).$$

⁵With respect to the Lebesgue measure on \mathbb{R}^n .

This result appears in [37] and follows easily from the almost everywhere theorem. Subharmonic addition extends to $u, v \in \text{USC}(X)$ by quasiconvex approximation in various situations. This extension has been accomplished for constant coefficient subequations \mathcal{F} in [11], for subequations associated to inhomogeneous pure second order equations in [43] and for fiberegular \mathcal{M} -monotone subequations \mathcal{F} in [14].

Subharmonic addition is very important when combined with the following implication between monotonicity and jet addition

$$(2.28) \quad \mathcal{F}_x + \mathcal{M}_x \subset \mathcal{F}_x \implies \mathcal{F}_x + \tilde{\mathcal{F}}_x \subset \tilde{\mathcal{M}}_x, \text{ for each } x \in X.$$

This combination has a very interesting consequence. If \mathcal{M} is a monotonicity cone subequation for \mathcal{F} , then sums of \mathcal{F} -subharmonics and $\tilde{\mathcal{F}}$ -subharmonics are $\tilde{\mathcal{M}}$ -subharmonics. Thus, when \mathcal{M} has constant coefficients, comparison for \mathcal{F} reduces to the validity of the zero maximum principle (ZMP) for $\tilde{\mathcal{M}}$ -subharmonics where $\tilde{\mathcal{M}}$ has constant coefficients since \mathcal{M} does. This is a constant coefficient potential theory and has been analyzed extensively in [11], where the validity of the (ZMP) is well understood for constant coefficient monotonicity cone subequations. This will be briefly reviewed at the end of Section 3 below.

2.5. A “tool kit” for \mathcal{F} -subharmonics (subsolutions)

Some of the “nuts and bolts” of handling \mathcal{F} -subharmonic functions are briefly described here. The first result is both useful for checking whether a given function is \mathcal{F} -subharmonic and also sheds light on the notion of viscosity subsolutions.

Lemma 2.12 (Definitional Comparison). *Let $\mathcal{F} \subset \mathcal{J}^2(X)$ be a subequation and consider $u \in \text{USC}(X)$.*

- (a) *If u is \mathcal{F} subharmonic on X , the following form of comparison holds on any bounded domain $\Omega \subset\subset X$:*

$$u + v \leq 0 \text{ on } \partial\Omega \implies u + v \leq 0 \text{ on } \Omega$$

for each $v \in \text{USC}(\bar{\Omega}) \cap C^2(\Omega)$ which is strictly $\tilde{\mathcal{F}}$ -subharmonic on Ω .

- (b) *Conversely, suppose that for each $x \in X$ there are arbitrarily small neighborhoods $\Omega \subset\subset X$ about x where the above form of comparison holds. Then u is \mathcal{F} -subharmonic on X .*

Part (a) is a well-known principle in the viscosity theory of PDEs and is here recast on the potential theory side with the help of duality. Part (b) is novel (also for its natural formulation on the operator theory side in both constrained and unconstrained cases) and shows that the **fundamental notion** in viscosity theory is the validity of this form of the comparison principle. The proof of definitional comparison can be found in [11] for constant coefficient \mathcal{F} and in [14] for the general case.

The next tool is often used to show that a given function is \mathcal{F} -subharmonic by a contradiction argument.

Lemma 2.13 (Bad Test Jet Lemma). *Let $\mathcal{F} \subset \mathcal{J}^2(X)$ be a subequation. Suppose that $u \in \text{USC}(X)$ is not \mathcal{F} -subharmonic at $x_0 \in X$. Then there exist $\epsilon > 0, \rho > 0$ and a 2-jet $J \notin \mathcal{F}_x$ such that the (unique) quadratic function Q_J with $J_{x_0}^2 Q_J = J$ is an upper test function for u at x in the following ϵ -strict sense:*

$$u(x) - Q_J(x) \leq -\epsilon|x - x_0|^2 \quad \forall x \in B_\rho(x_0) \quad \text{with equality at } x_0.$$

This is merely the contrapositive of the definition of being subharmonic in x_0 making use of ϵ -strict upper test jets yield an equivalent definition see [26].

In addition to the C^2 -coherence property and the uniform translation property (for continuous \mathcal{M} -monotone \mathcal{F}) discussed above, one has many additional properties which are useful in various constructions.

Proposition 2.14 (Elementary properties of $\mathcal{F}(X)$). *Let $\mathcal{F} \subset \mathcal{J}^2(X)$ be a subequation. Then the following properties hold:*

- (i) **local:** $u \in \text{USC}(X)$ is locally \mathcal{F} -subharmonic $\Leftrightarrow u \in \mathcal{F}(X)$;
- (ii) **maximum:** $u, v \in \mathcal{F}(X) \implies \max\{u, v\} \in \mathcal{F}(X)$;
- (iii) **sliding:** $u \in \mathcal{F}(X) \implies u - m \in \mathcal{F}(X)$ for any $m > 0$;
- (iv) **decreasing limits:** if $\{u_k\}_{k \in \mathbb{N}} \subset \mathcal{F}(X)$ is a decreasing sequence then $u := \lim_{k \rightarrow \infty} u_k \in \mathcal{F}(X)$;
- (v) **uniform limits:** if $\{u_k\}_{k \in \mathbb{N}} \subset \mathcal{F}(X)$ locally uniformly converges to u then $u \in \mathcal{F}(X)$;
- (vi) **families locally bounded above:** if $\mathfrak{F} \subset \mathcal{F}(X)$ is a non-empty family of functions which are locally uniformly bounded from above, then the upper semicontinuous envelope⁶ u^* of the Perron function $u(\cdot) := \sup_{w \in \mathfrak{F}} w(\cdot)$ belongs to $\mathcal{F}(X)$.

⁶We recall that $u^*(x) := \limsup_{r \rightarrow 0^+} \{u(y) : y \in X \cap \overline{B}_r(x)\}$ for each $x \in X$ and that u^x is the minimal USC(X) function with $u \leq u^*$ on X .

Furthermore, if \mathcal{F} has constant coefficients, the following also holds:

(vii) **translation:** $u \in \mathcal{F}(X) \iff u(\cdot - y) \in \mathcal{F}(X + y)$, for any $y \in \mathbb{R}^n$.

These are familiar properties for the viscosity theory of nonlinear elliptic PDEs, although stated typically only in the unconstrained case. See [26] for the proofs of this general potential theoretic version.

3. Comparison by monotonicity, duality and fiberegularity

In this section, we present a clean, elegant and flexible method for proving *comparison* (the comparison principle) in nonlinear potential theory. It makes use of the two ingredients monotonicity and duality, along with some form of regularity of the subequation. There are three incarnations of the needed regularity: constant coefficients, tameness for subequations defined by inhomogeneous equations, and fiberegularity. The method works when a given subequation admits a suitable constant coefficient monotonicity cone subequation \mathcal{M} . When the comparison principle is combined with a correspondence principle, comparison can be transferred to nonlinear elliptic PDEs.

3.1. Statement and history of the general result

The method has evolved from the constant coefficient pure second order case [24]. The general theorem in Euclidian space is the following result [14, Theorem 4.3].

Theorem 3.1 (A general comparison theorem). *Let $\Omega \subset \mathbb{R}^n$ be a bounded domain. Suppose that a subequation $\mathcal{F} \subset \mathcal{J}^2(\Omega)$ is fiberegular and \mathcal{M} -monotone on Ω for some monotonicity cone subequation \mathcal{M} . If \mathcal{M} admits a C^2 -strict subharmonic ψ on Ω , then comparison holds for \mathcal{F} on $\overline{\Omega}$; that is,*

$$(CP) \quad u \leq w \text{ on } \partial\Omega \implies u \leq w \text{ on } \Omega$$

for all $u \in \text{USC}(\overline{\Omega})$ which are \mathcal{F} -subharmonic on Ω , and for all $w \in \text{LSC}(\overline{\Omega})$ which are \mathcal{F} -superharmonic on Ω .

The evolution of this result can be summarized in the following way.

- (1) For $\mathcal{F}_x \equiv \mathcal{F} \subset \mathcal{S}(n)$ (constant coefficient pure second order) see [24], where $\mathcal{M} = \mathcal{P}$ and $\psi(x) := \frac{1}{2}|x|^2$. Here one can say

(CP) holds for all subequations $\mathcal{F} \subset \mathcal{S}(n)$ and for all domains $\Omega \subset \subset \mathbb{R}^n$.

- (2) For $\mathcal{F} \subset \Omega \times \mathcal{S}(n)$ (fiberegular variable coefficient pure second order) see [12], where $\mathcal{M} = \mathcal{P}$ and $\psi(x) := \frac{1}{2}|x|^2$. Here one can say

(CP) holds for all fiberegular subequations $\mathcal{F} \subset \Omega \times \mathcal{S}(n)$ and for all domains $\Omega \subset \subset \mathbb{R}^n$.

Of course an interesting case here is an inhomogeneous subequation \mathcal{F} defined by $F(D^2u) - f(x) \geq 0$. In [43], assuming a condition called *tame* on the operator F , (CP) was established. One can show that F tame implies that \mathcal{F} is fiberegular, so this result is a special case of (2).

- (3) For $\mathcal{F}_x \equiv \mathcal{F} \subset \mathbb{R} \times \mathcal{S}(n)$ (constant coefficient gradient free) see [26, Theorem 13.4] where $\mathcal{M} = \mathcal{Q} = \mathcal{N} \times \mathcal{P}$ and $\psi(x) := \frac{1}{2}(|x|^2 - R^2)$, $R \gg 0$. Here one can say

(CP) holds for all subequations $\mathcal{F} \subset \mathbb{R} \times \mathcal{S}(n)$ and for all domains $\Omega \subset \subset \mathbb{R}^n$.

- (4) For $\mathcal{F} \subset \Omega \times (\mathbb{R} \times \mathcal{S}(n))$ (fiberegular variable coefficient gradient-free) see [13], where $\mathcal{M} = \mathcal{Q} = \mathcal{N} \times \mathcal{P}$ and $\psi(x) := \frac{1}{2}(|x|^2 - R^2)$, $R \gg 0$. Here one can say

(CP) holds for all fiberegular subequations $\mathcal{F} \subset \Omega \times \mathbb{R} \times \mathcal{S}(n)$ and for all domains $\Omega \subset \subset \mathbb{R}^n$.

- (5) For $\mathcal{F}_x \equiv \mathcal{F} \subset \mathcal{J}^2(\Omega)$ (general constant coefficients) see [11], where there is also a complete study of which cones \mathcal{M} admit ψ .
 (6) For the general case $\mathcal{F} \subset \mathcal{J}^2(\Omega)$ (fiberegular) see [14], where one imports the class of admissible cones \mathcal{M} from the constant coefficient case.

3.2. Outline of the proof

The main steps in the proof are the following.

Step 1: First, use **duality** to reformulate (CP) as:

$$(CP') \quad u + v \leq 0 \text{ on } \partial\Omega \quad \implies \quad u + v \leq 0 \text{ on } \Omega$$

for all $u \in USC(\bar{\Omega}) \cap \mathcal{F}(\Omega)$ and $v \in USC(\bar{\Omega}) \cap \tilde{\mathcal{F}}(\Omega)$ (both subharmonic). Define $v := -w$ and then the equivalence in (2.10) translates to the equivalence of (CP) and (CP'). Next, note that (CP') is just the zero maximum principle (ZMP) for the sum of \mathcal{F} and $\tilde{\mathcal{F}}$ subharmonics:

$$(3.1) \quad (\text{ZMP}) \quad z \leq 0 \text{ on } \partial\Omega \quad \implies \quad z \leq 0 \text{ on } \partial\Omega,$$

$\forall z \in USC(\bar{\Omega}) \cap (\mathcal{F}(\Omega) + \tilde{\mathcal{F}}(\Omega))$. Thus it remains to prove (ZMP) in (3.1).

Step 2 (Jet Addition): Establish the fundamental *jet addition formula* ([30, Lemma 4.1.2])

$$(3.2) \quad \mathcal{F}_x + \mathcal{M}_x \subset \mathcal{F}_x \implies \mathcal{F}_x + \tilde{\mathcal{F}}_x \subset \tilde{\mathcal{M}}_x, \text{ for each } x \in X.$$

This formula follows from elementary properties of duality and monotonicity.

Step 3: Establish the *Almost Everywhere Theorem* and the quasiconvex version of the *Subharmonic Addition Theorem* (see Theorems 2.10 and 2.11 to conclude

$$z = u + v \in \tilde{\mathcal{M}}(\Omega)$$

if $u \in \mathcal{F}(\Omega)$ and $v \in \tilde{\mathcal{F}}(\Omega)$ are locally quasiconvex. This difficult step relies on the Jensen or Slodkowski or Federer Lemmas.

Step 4: Use **fiberegularity** to prove the full *Subharmonic Addition Theorem*

$$\mathcal{F}(\Omega) + \tilde{\mathcal{F}}(\Omega) \subset \tilde{\mathcal{M}}(\Omega).$$

Step 5: Apply the following constant coefficient result [11, Theorem 6.2].

Theorem 3.2 (The Zero Maximum Principle for Dual Monotonicity Cones). *Suppose that \mathcal{M} is a constant coefficient monotonicity cone subequation that admits a C^2 -strict subsolution ψ on a domain $\Omega \subset\subset \mathbb{R}^n$. Then the zero maximum principle holds for $\tilde{\mathcal{M}}$ on $\bar{\Omega}$; that is,*

$$(ZMP) \quad z \leq 0 \text{ on } \partial\Omega \implies z \leq 0 \text{ on } \Omega$$

for all $z \in \text{USC}(\bar{\Omega}) \cap \tilde{\mathcal{M}}(\Omega)$.

Proof. $\tilde{\mathcal{M}}$ is a (constant coefficient) subequation and hence satisfies the *sliding property*

$$z - m \in \tilde{\mathcal{M}}(\Omega) \text{ for each } m \in [0, +\infty).$$

Since $z - m < 0$ on $\partial\Omega$ compact

$$z - m + \varepsilon\psi \leq 0 \text{ on } \partial\Omega \text{ for each } \varepsilon \text{ sufficiently small.}$$

Since $z - m \in \tilde{\mathcal{M}}(\Omega)$ and since $\varepsilon\psi \in C(\bar{\Omega}) \cap C^2(\Omega)$ is strictly \mathcal{M} -subharmonic, by the *definitional comparison* Lemma 2.12 (with $\mathcal{F} = \tilde{\mathcal{M}}$ and $\tilde{\mathcal{F}} = \tilde{\mathcal{M}} = \mathcal{M}$) one has

$$z - m + \varepsilon\psi \leq 0 \text{ on } \Omega \text{ for each } \varepsilon \text{ sufficiently small.}$$

Passing to the limit for $\varepsilon \rightarrow 0^+$, and then $m \rightarrow 0^+$ yields $z \leq 0$ on Ω . □

The utility of the General Comparison Theorem 3.1 is greatly facilitated by the detailed study of monotonicity cone subequations in [11], which we briefly review. There is a three parameter *fundamental family* of monotonicity cone subequations consisting of

$$\mathcal{M}(\gamma, \mathcal{D}, R) := \left\{ (r, p, A) \in \mathcal{J}^2 : r \leq -\gamma|p|, p \in \mathcal{D}, A \geq \frac{|p|}{R} I \right\}$$

where

$$\gamma \in [0, +\infty), R \in (0, +\infty] \text{ and } \mathcal{D} \subseteq \mathbb{R}^n,$$

where \mathcal{D} is a *directional cone*; that is, a closed convex cone with vertex at the origin and non-empty interior (see Definition 5.2 and Remark 5.9 of [11]). The family is fundamental in the sense that for any monotonicity cone subequation, there exists an element $\mathcal{M}(\gamma, \mathcal{D}, R)$ of the family with $\mathcal{M}(\gamma, \mathcal{D}, R) \subset \mathcal{M}$ (see Theorem 5.10 of [11]). Hence if \mathcal{F} is an \mathcal{M} -monotone subequation, then it is $\mathcal{M}(\gamma, \mathcal{D}, R)$ -monotone for some triple (γ, \mathcal{D}, R) . Moreover, from Theorem 6.3 of [11], given any element $\mathcal{M} = \mathcal{M}(\gamma, \mathcal{D}, R)$ of the fundamental family, one knows for which domains $\Omega \subset\subset \mathbb{R}^n$ there is a C^2 -strict \mathcal{M} -subharmonic and hence for which domains Ω one has the (ZMP) for $\widetilde{\mathcal{M}}$ -subharmonics according to Theorem 3.2. There is a simple dichotomy. If $R = +\infty$, then arbitrary bounded domains Ω may be used, while in the case of R finite, any Ω which is contained in a translate of the truncated cone $\mathcal{D}_R := \mathcal{D} \cap B_R(0)$.

4. The correspondence principle

In this section, we discuss structural conditions on a given proper elliptic operator F with domain $\mathcal{G} \subset \mathcal{J}^2(X)$ which ensure that the constraint set \mathcal{F} defined by the compatibility relation (1.14)

$$(4.1) \quad \mathcal{F} := \{(x, J) \in \mathcal{G} : F(x, J) \geq 0\}$$

satisfies the two conditions needed for the the Correspondence Principle of Theorem 1.3. We recall that these two conditions are:

$$(4.2) \quad \mathcal{F} \text{ defined by (4.1) is a subequation (in the sense of Definition 2.1)}$$

and compatibility (1.16) between \mathcal{F} and F :

$$(4.3) \quad \text{Int } \mathcal{F} = \{(x, J) \in \mathcal{G} : F(x, J) > 0\},$$

or equivalently

$$(4.4) \quad \partial\mathcal{F} = \{(x, J) \in \mathcal{F} : F(x, J) = 0\}.$$

\mathcal{F} defined by (4.1) will be a subequation if it satisfies the three properties of positivity (P), negativity (N) and topological stability (T). The first two (P) and (N) are equivalent to the (fiberwise) monotonicity property that for each $x \in X$

$$(r, p, A) \in \mathcal{F}_x \Rightarrow (r + s, p, A + P) \in \mathcal{F}_x, \quad \forall s \leq 0 \text{ in } \mathbb{R}, P \geq 0 \text{ in } \mathcal{S}(n),$$

which clearly follows from the same monotonicity property for the domain \mathcal{G} and the proper ellipticity of F on \mathcal{G} (see (1.10)):

$$F(x, r+s, p, A+P) \geq F(x, r, p, A), \quad \forall (r, p, A) \in \mathcal{G}_x, s \leq 0 \text{ in } \mathbb{R}, P \geq 0 \text{ in } \mathcal{S}(n).$$

This leaves the topological property (T). Recall that this property requires the three conditions

$$\begin{aligned} \text{(T1)} \quad & \mathcal{F} = \overline{\text{Int } \mathcal{F}}; \\ \text{(T2)} \quad & \mathcal{F}_x = \overline{\text{Int } (\mathcal{F}_x)}, \quad \forall x \in X; \\ \text{(T3)} \quad & (\text{Int } \mathcal{F})_x = \text{Int } (\mathcal{F}_x), \quad \forall x \in X. \end{aligned}$$

In the constant coefficient case, property (T) reduces to property (T1). In the gradient free case, one can show that property (T1) follows from properties (P) and (N) since \mathcal{F} is closed. In the general constant coefficient case, a sufficient condition for (T1) is that \mathcal{F} is closed and is \mathcal{M} -monotone for some monotonicity cone subequation (see Proposition 4.7 of [11]). \mathcal{F} defined by (4.1) is closed by the continuity of F . Hence, if (F, \mathcal{G}) is a constant coefficient \mathcal{M} -monotone pair, then \mathcal{F} defined by (4.1) is indeed a subequation.

In the variable coefficient case, assuming that (F, \mathcal{G}) is an \mathcal{M} -monotone pair, then the argument above (fiberwise) yields the property (T2). This leaves properties (T1) and (T3). It is not hard to see that if \mathcal{F} is closed, then properties (T2) plus (T3) imply (T1) (see Proposition A.2 of [14]). Hence for a \mathcal{M} -monotone pair (F, \mathcal{G}) , the constraint set \mathcal{F} defined by (4.1) will be a subequation if \mathcal{F} is closed and satisfies (T3). Moreover, since the inclusion $(\text{Int } \mathcal{F})_x \subset \text{Int } (\mathcal{F}_x)$ is automatic for each $x \in X$, (T3) reduces to the reverse inclusion, which holds provided that \mathcal{F} is \mathcal{M} -monotone and fiberegular in the sense of Definition 2.7. This fact is proved in Proposition A.5 of [14]. Moreover, as shown in Theorem 6.1 of [14], \mathcal{F} will be fiberegular if \mathcal{G} is fiberegular

provided that F satisfies a mild *regularity condition* (see (4.5) below). In addition, fiberegularity of \mathcal{F} ensures that \mathcal{F} is closed (see Proposition A.6 of [14]). We collect some of these observations in the following Lemma.

Lemma 4.1. *Suppose that (F, \mathcal{G}) is an \mathcal{M} -monotone operator-subequation pair for some monotonicity cone subequation, with $\mathcal{G} = \mathcal{J}^2(X)$ or $\mathcal{G} \subsetneq \mathcal{J}^2(X)$ a fiberegular subequation. Suppose that (F, \mathcal{G}) satisfies the regularity condition: for some fixed $J_0 \in \text{Int } \mathcal{M}$, given $\Omega \subset\subset X$ and $\eta > 0$, there exists $\delta = \delta(\eta, \Omega) > 0$ such that*

$$(4.5) \quad F(y, J + \eta J_0) \geq F(x, J), \quad \forall x, y \in \Omega \text{ with } |x - y| < \delta.$$

Then the constraint set \mathcal{F} defined by (4.1) is a (fiberegular \mathcal{M} -monotone) subequation.

Finally, we discuss structural conditions on a proper elliptic operator F with domain $\mathcal{G} \subset \mathcal{J}^2(X)$ for which the constraint set \mathcal{F} defined by (4.1) satisfies compatibility (4.3) (or equivalently (4.4)). In the situation of Lemma 4.1, which ensures that \mathcal{F} defined by (4.1) is a subequation, by the topological property (T3) it suffices to have (4.3) fiberwise; that is,

$$(4.6) \quad \text{Int } \mathcal{F}_x = \{J \in \mathcal{G}_x : F(x, J) > 0\}, \quad \forall x \in X.$$

This condition is often easily checked for a given pair (F, \mathcal{G}) which determines \mathcal{F} by checking that $F(x, J) = 0$ for $J \in \partial \mathcal{F}_x$ and using some strict monotonicity such as: for each $x \in X$ with some fixed $J_0 \in \text{Int } \mathcal{M}$ there exists $t_0 > 0$ such that

$$(4.7) \quad F(x, J + tJ_0) > F(x, J), \quad \forall t \in (0, t_0), \forall J \in \partial \mathcal{F}_x.$$

Compatibility in this situation of a *homogeneous equation* $F(J^2u) = 0$ is relatively simple because one need only pay attention to F in a neighborhood of the zero locus of F (with domain \mathcal{G}).

More structure is required if one would like to treat the *inhomogeneous equation*

$$(4.8) \quad F(J^2u) = \psi, \quad \psi \in C(X)$$

for a given constant coefficient operator F . This is true even for constant sources $\psi = c$. This case has been studied extensively in [11], which we now review. There the domain \mathcal{G} was denoted instead by \mathcal{F} , which we will also

do below. In the constrained case, where $\mathcal{F} \subset \mathcal{J}^2$ is a (constant coefficient) subequation, **compatibility** is defined by the two conditions

$$(4.9) \quad \inf_{\mathcal{F}} F \text{ is finite (and denoted by } c_0)$$

and

$$(4.10) \quad \partial\mathcal{F} = \{J \in \mathcal{F} : F(J) = c_0\}.$$

Given an operator-subequation pair (F, \mathcal{F}) , the values $c \in F(\mathcal{F})$ are called *admissible levels of F* , since otherwise the level set $\{F = c\}$ is empty.

More is needed in order to treat the inhomogeneous equation $F(J^2u) = c$ for all of the admissible levels. In order to avoid some obvious pathologies, one must assume that the operator $F \in C(\mathcal{F})$ is **topologically tame**; that is, for each admissible level $c \in F(\mathcal{F})$,

$$(4.11) \quad \text{the level set } \mathcal{F}(c) := \{J \in \mathcal{F} : F(J) = c\} \text{ has empty interior.}$$

This condition serves an additional purpose. Namely, if (F, \mathcal{F}) is a proper elliptic operator-subequation pair with F topologically tame, then for every admissible level $c \in F(\mathcal{F})$ the upper level set

$$(4.12) \quad \mathcal{F}_c := \{J \in \mathcal{F} : F(J) \geq c\}$$

satisfies the topological property (T). Hence each \mathcal{F}_c is a subequation since properties (P) and (N) are encoded by the proper ellipticity. The obvious pathologies eliminated by topological tameness of F are explained in [11, section 11.1]. For example, if some admissible level set $\mathcal{F}(c)$ has non-empty interior, then one has many counterexamples for comparison by considering perturbations $v + \varphi$ of a local C^2 solution to $F(J^2v) = c$ with φ smooth, compactly supported and with small C^2 -norm.

Some strict monotonicity for the operator F provides a convenient structural condition on the operator which eliminates such pathologies. More precisely, for constant coefficient compatible pairs (F, \mathcal{F}) which are \mathcal{M} -monotone for some monotonicity cone subequation \mathcal{M} , topological tameness (4.11) is equivalent the following structural condition of **strict \mathcal{M} -monotonicity** on F :

$$(4.13) \quad \exists J_0 \in \text{Int } \mathcal{M} \text{ such that } F(J + tJ_0) > F(J), \quad \forall J \in \mathcal{F}, \forall t > 0.$$

In the gradient-free case this monotonicity is the weakest possible notion of being strictly proper elliptic. Moreover, these equivalent notions (4.11)

and (4.13) are also equivalent to any one of the following three conditions (see Theorem 11.10 of [11]):

- 1) $F(J + J_0) > F(J)$ for each $J \in \mathcal{F}$ and each $J_0 \in \text{Int } \mathcal{M}$;
- 2) $\{J \in \mathcal{F} : F(J) > c\} = \text{Int } \mathcal{F}_c$ for each admissible level $c \in F(\mathcal{F})$;
- 3) $\mathcal{F}(c) = \mathcal{F}_c \cap (-\tilde{\mathcal{F}}_c)$ for each admissible level $c \in F(\mathcal{F})$.

Combining compatibility with strict \mathcal{M} -monotonicity, one has a correspondence principle for the solutions of the inhomogeneous equation (4.8), which are precisely the \mathcal{F}_ψ -harmonics for the subequation with fibers

$$(4.14) \quad \mathcal{F}_{\psi(x)} = \{F \in \mathcal{F} : F(J) \geq \psi(x)\}, \quad x \in X.$$

More precisely, one has the following result whose proof follows directly from the proof of the constant source case $\psi = c$ given in Theorem 11.13 of [11].

Theorem 4.2 (Correspondence principle). *Suppose that (F, \mathcal{F}) is a compatible \mathcal{M} -monotone (operator-subequation) pair for some monotonicity cone subequation \mathcal{M} with F strictly \mathcal{M} -monotone in the sense (4.13). Then, for any $\psi \in C(X)$ taking values in $F(\mathcal{F})$, a function $u \in C(X)$ is an \mathcal{F} -admissible solution of the equation*

$$(4.15) \quad F(J^2u(x)) = \psi(x), \quad \forall x \in X$$

if and only if u is \mathcal{F}_ψ -harmonic in X . In particular, for $u \in \text{USC}(X)$ and $w \in \text{LSC}(X)$ one has

u is an \mathcal{F} -admissible subsolution of (4.15) $\Leftrightarrow u$ is \mathcal{F}_ψ -subharmonic in X

and

w is an \mathcal{F} -admissible supersolution of (4.15) $\Leftrightarrow -w$ is $\tilde{\mathcal{F}}_\psi$ -subharmonic in X .

References

- [1] M. BARDI and P. MANNUCCI, *On the Dirichlet problem for non-totally degenerate fully nonlinear elliptic equations*, Commun. Pure Appl. Anal. **73** (2006), 709–731. [MR2246004](#)
- [2] G. BARLES and J. BUSCA, *Existence and comparison results for fully nonlinear degenerate elliptic equations without zeroth-order term*, Comm. Partial Differential Equations **26** (2001), 2323–2337. [MR1876420](#)

- [3] H. BERESTYCKI, L. NIRENBERG and S. VARADHAN, *The principle eigenvalue and maximum principle for second order elliptic operators in general domains*, Comm. Pure Appl. Math. **47** (1994), 47–92. [MR1258192](#)
- [4] I. BIRINDELLI, G. GALISE and H. ISHII, *A family of degenerate elliptic operators: maximum principle and its consequences*, Ann. Inst. H. Poincaré Anal. Non Linéaire **35** (2018), 417–441. [MR3765548](#)
- [5] I. BIRINDELLI and K.R. PAYNE, *Principal eigenvalues for k -Hessian operators by maximum principle methods*, Math. Eng. **3** (2021), Paper No. 021, 37 pp. [MR4146717](#)
- [6] K.K. BRUSTAD, *Counterexamples to the comparison principle in the special Lagrangian potential equation*, [arXiv:2206.09373v2](#), 7 pages, published online 18 Oct 2022.
- [7] L. CAFFARELLI, L. NIRENBERG and J. SPRUCK, *The Dirichlet problem for nonlinear second-order elliptic equations III. Functions of the eigenvalues of the Hessian*, Acta Math. **155** (1985), 261–301. [MR0806416](#)
- [8] L. CAFFARELLI and X. CABRÉ, *Fully Nonlinear Elliptic Equations*, American Mathematical Society Colloquium Publications, Vol. 43, American Mathematical Society, Providence, RI, 1995. [MR1351007](#)
- [9] S.Y. CHENG and S.T. YAU, *Complete affine hypersurfaces. Part I. The completeness of affine metrics*, Comm. Pure Appl. Math. **39** (1986), 839–866. [MR0859275](#)
- [10] J. CHU, *Quantitative stratification of F -subharmonic functions*, Comm. Geom. Anal. Math. **29** (2021), 1335–1389. [MR4367428](#)
- [11] M. CIRANT, F.R. HARVEY, H.B. LAWSON JR., and K.R. PAYNE, *Comparison Principles by Monotonicity and Duality for Constant Coefficient Nonlinear Potential Theory and PDEs*, Annals of Mathematics Studies, Vol. 218, Princeton University Press, Princeton, NJ, 2023. [MR4671835](#)
- [12] M. CIRANT and K.R. PAYNE, *On viscosity solutions to the Dirichlet problem for elliptic branches of nonhomogeneous fully nonlinear equation*, Publ. Mat. **61** (2017), 529–575. [MR3677871](#)
- [13] M. CIRANT and K.R. PAYNE, *Comparison principles for viscosity solutions of elliptic branches of fully nonlinear equations independent of the gradient*, Math. Eng. **3** (2021), Paper No. 045, 45 pp. [MR4147574](#)

- [14] M. CIRANT, K.R. PAYNE and D.F. RADAELLI, *Comparison principles for nonlinear potential theory and PDEs with continuous coefficients and sufficient monotonicity*, *Nonlinear Anal.* **35** (2023), Paper No. 113343, 52 pp. [MR4612711](#)
- [15] M.G. CRANDALL, H. ISHII and P.-L. LIONS, *User's guide to viscosity solutions of second order partial differential equations*, *Bull. Amer. Math. Soc.* **27** (1992), 1–67. [MR1118699](#)
- [16] G. DE PHILIPPIS and A. FIGALLI, *The Monge-Ampère equation and its link to optimal transportation*, *Bull. Amer. Math. Soc. (N.S.)* **51** (2014), 527–580. [MR3237759](#)
- [17] S. DINEW, H.-S. DO and T.D. TÔ, *A viscosity approach to the Dirichlet problem for degenerate complex Hessian-type equations*, *Anal. PDE* **12** (2019), 505–535. [MR3861899](#)
- [18] F. FORSTERNIČ, *Every smoothly bounded p -convex domain in \mathbb{R}^n admits a p plurisubharmonic defining function*, [arXiv:2111.08113v2](#), 8 pages, published online 12 Jan 2022. [MR4370488](#)
- [19] L. GÅRDING, *An inequality for hyperbolic polynomials*, *J. Math. Mech.* **8** (1959), 957–965. [MR0113978](#)
- [20] D. GILBARG and N.S. TRUDINGER, *Elliptic Partial Differential Equations of Second Order*, Second Edition, Grundlehren der Mathematischen Wissenschaften, Vol. 224, Springer-Verlag, Berlin, 1983. [MR0737190](#)
- [21] F.R. HARVEY and H.B. LAWSON, JR., *Calibrated geometries*, *Acta Math.* **148** (1982), 47–157. [MR0666108](#)
- [22] F.R. HARVEY and H.B. LAWSON, JR., *An introduction to potential theory in calibrated geometry*, *Amer. J. Math.* **131** (2009), 893–944. [MR2543918](#)
- [23] F.R. HARVEY and H.B. LAWSON, JR., *Duality of positive currents and plurisubharmonic functions in calibrated geometry*, *Amer. J. Math.* **131** (2009), 1211–1239. [MR2555839](#)
- [24] F.R. HARVEY and H.B. LAWSON, JR., *Dirichlet duality and the nonlinear Dirichlet problem*, *Comm. Pure Appl. Math.* **62** (2009), 396–443. [MR2487853](#)
- [25] F.R. HARVEY and H.B. LAWSON, JR., *Hyperbolic polynomials and the Dirichlet problem*, [arXiv:0912.5220v2](#), 19 March 2010, 1–48.

- [26] F.R. HARVEY and H.B. LAWSON, JR., *Dirichlet duality and the nonlinear Dirichlet problem on Riemannian manifolds*, J. Differential Geom. **88** (2011), 395–482. [MR2844439](#)
- [27] F.R. HARVEY and H.B. LAWSON, JR., *Plurisubharmonicity in a general geometric context*, in: *Geometry and Analysis*, Adv. Lect. Math. (ALM), Vol. 17, No. 1, 363–402, Int. Press, Somerville, MA, 2011. [MR2882430](#)
- [28] F.R. HARVEY and H.B. LAWSON, JR., *Geometric plurisubharmonicity and convexity: an introduction*, Adv. Math. **230** (2012), 2428–2456. [MR2927376](#)
- [29] F.R. HARVEY and H.B. LAWSON, JR., *Gårding’s theory of hyperbolic polynomials*, Comm. Pure Appl. Math. **66** (2013), 1102–1128. [MR3055586](#)
- [30] F.R. HARVEY and H.B. LAWSON, JR., *Existence, uniqueness and removable singularities for nonlinear partial differential equations in geometry*, in: *Surveys in Differential Geometry* **18**, pp. 103–156, International Press of Boston, Inc., Somerville, MA, 2013. [MR3087918](#)
- [31] F.R. HARVEY and H.B. LAWSON, JR., *p-convexity, p-plurisubharmonicity and the Levi problem*, Indiana Univ. Math. J. **62** (2013), 149–169. [MR3158505](#)
- [32] F.R. HARVEY and H.B. LAWSON, JR., *Removable singularities for nonlinear subequations*, Indiana Univ. Math. J. **63** (2014), 1525–1552. [MR3283560](#)
- [33] F.R. HARVEY and H.B. LAWSON, JR., *The restriction theorem for fully nonlinear subequations*, Ann. Inst. Fourier (Grenoble) **64** (2014), 217–265. [MR3330548](#)
- [34] F.R. HARVEY and H.B. LAWSON, JR., *The equivalence of viscosity and distributional solutions for convex subequations – a strong Bellman principle*, Bull. Braz. Math. Soc. (N.S) **44** (2014), 621–652. [MR3167126](#)
- [35] F.R. HARVEY and H.B. LAWSON, JR., *Potential theory on almost complex manifolds*, Ann. Inst. Fourier (Grenoble) **65** (2015), 171–210. [MR3449151](#)
- [36] F.R. HARVEY and H.B. LAWSON, JR., *Notes on the differentiation of quasi-convex functions*, [arXiv:1309.1772v3](#), 30 July 2016, 1–17.
- [37] F.R. HARVEY and H.B. LAWSON, JR., *The ae theorem and addition theorems for quasi-convex functions*, [arXiv:1309.1770v3](#), 30 July 2016, 1–12.

- [38] F.R. HARVEY and H.B. LAWSON, JR., *Characterizing the strong maximum principle for constant coefficient subequations*, Rend. Mat. Appl. (7) **37** (2016), 63–104. [MR3622305](#)
- [39] F.R. HARVEY and H.B. LAWSON, JR., *The Dirichlet problem with prescribed interior singularities*, Adv. Math. **303** (2016), 1319–1357. [MR3552552](#)
- [40] F.R. HARVEY and H.B. LAWSON, JR., *Tangents to subsolutions: existence and uniqueness, Part II*, J. Geom. Anal. **27** (2017), 2190–2223. [MR3667427](#)
- [41] F.R. HARVEY and H.B. LAWSON, JR., *Lagrangian potential theory and a Lagrangian equation of Monge-Ampère type*, in: *Surveys in Differential Geometry*, Vol. 22, 217–258, International Press of Boston, Inc., Somerville, 2017. [MR3838119](#)
- [42] F.R. HARVEY and H.B. LAWSON, JR., *Tangents to subsolutions: existence and uniqueness, Part I*, Ann. Fac. Sci. Toulouse Math. (6) **27** (2018), 777–848. [MR3884610](#)
- [43] F.R. HARVEY and H.B. LAWSON, JR., *The inhomogeneous Dirichlet Problem for natural operators on manifolds*, Ann. Inst. Fourier (Grenoble) **69** (2019), 3017–3064. [MR4286829](#)
- [44] F.R. HARVEY and H.B. LAWSON, JR., *Plurisubharmonics in general potential theories*, in: *Advances in Complex Geometry*, Contemp. Math., Vol. 735, 145–168, Amer. Math. Soc., Providence, RI, 2019. [MR3996487](#)
- [45] F.R. HARVEY and H.B. LAWSON, JR., *A generalization of PDEs from a Krylov point of view*, Adv. Math. **372** (2020), 107298, 32 pp. [MR4127162](#)
- [46] F.R. HARVEY and H.B. LAWSON, JR., *Pseudoconvexity for the special Lagrangian potential equation*, Calc. Var. Partial Differential Equations, **60** (2021), Paper No. 6, 37 pp. [MR4179860](#)
- [47] F.R. HARVEY, H.B. LAWSON, JR. and S. PLIŚ, *Smooth approximation of plurisubharmonic functions on almost complex manifolds*, Math. Ann. **366** (2016), 929–940. [MR3563228](#)
- [48] H. ISHII and P.-L. LIONS, *Viscosity solutions of fully nonlinear second-order elliptic partial differential equations*, J. Differential Equations **83** (1990), 26–78. [MR1031377](#)
- [49] R. JENSEN, *The maximum principle for viscosity solutions of fully nonlinear second order partial differential equations*, Arch. Rat. Mech. Anal. **101** (1988), 1–27. [MR0920674](#)

- [50] N.V. KRYLOV, *Sequences of convex functions, and estimates of the maximum of the solution of a parabolic equation*, Siberian Math. J. **17** (1976), 226–236. [MR0420016](#)
- [51] N.V. KRYLOV, *Nonlinear Elliptic and Parabolic Equations of Second Order*, Mathematics and its Applications (Soviet Series), Vol. 7, D. Reidel Publishing Co., Dordrecht, 1987. [MR0901759](#)
- [52] N.V. KRYLOV, *On the general notion of fully nonlinear second-order elliptic equations*, Trans. Amer. Math. Soc. **347** (1995), 857–895. [MR1284912](#)
- [53] K.R. PAYNE and D.F. REDAELLI, *Primer on quasiconvex functions in nonlinear potential theory*, preprint 2022.
- [54] Z. SŁODKOWSKI, *The Bremermann-Dirichlet problem for q -plurisubharmonic functions*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4), **11** (1984), 303–326. [MR0764948](#)
- [55] N.S. TRUDINGER, *The Dirichlet problem for the prescribed curvature equations*, Arch. Rational Mech. Anal. **111** (1990), 153–179. [MR1057653](#)
- [56] C. VILLANI, *Optimal Transport. Old and New*, Grundlehren der Mathematischen Wissenschaften, Vol. 338, Springer-Verlag, Berlin, 2009. [MR2459454](#)
- [57] J.B. WALSH, *Continuity of envelopes of plurisubharmonic functions*, J. Math. Mech. **18** (1968/1969), 143–148. [MR0227465](#)

F. Reese Harvey
Department of Mathematics
Rice University
P.O. Box 1892
Houston, TX 77005-1892
USA
E-mail: harvey@rice.edu

Kevin R. Payne
Dipartimento di Matematica “F. Enriques”
Università di Milano
Via C. Saldini 50
20133–Milano
Italy
E-mail: kevin.payne@unimi.it