

Surveys of Modern Mathematics  
Volume I

# Analytic Methods in Algebraic Geometry

by Jean-Pierre Demailly

 International Press  
[www.intlpress.com](http://www.intlpress.com)



高等教育出版社  
HIGHER EDUCATION PRESS

Surveys of Modern Mathematics, Volume I  
Analytic Methods in Algebraic Geometry

Jean-Pierre Demailly  
Institut Fourier, Laboratoire de Mathématiques, Saint-Martin d'Hères, France

*2010 Mathematics Subject Classification.* 14-XX.

Copyright © 2010, 2012 by International Press, Somerville, Massachusetts, U.S.A., and  
by Higher Education Press, Beijing, China.

This work is published and sold in China exclusively by Higher Education Press  
of China.

All rights reserved. Individual readers of this publication, and non-profit libraries acting  
for them, are permitted to make fair use of the material, such as to copy a chapter for use  
in teaching or research. Permission is granted to quote brief passages from this  
publication in reviews, provided the customary acknowledgement of the source is given.  
Republication, systematic copying, or mass reproduction of any material in this  
publication is permitted only under license from International Press. Excluded from these  
provisions is material in articles to which the author holds the copyright. (If the author  
holds copyright, notice of this will be given with the article.) In such cases, requests for  
permission to use or reprint should be addressed directly to the author.

ISBN: 978-1-57146-234-3

Printed in the United States of America.

16 15 14 13 12    1 2 3 4 5 6 7 8 9

## SURVEYS OF MODERN MATHEMATICS

### Series Editors

Shing-Tung Yau  
Harvard University  
Cambridge, Massachusetts  
U.S.A.

Lizhen Ji  
University of Michigan, Ann Arbor  
U.S.A.

Yat-Sun Poon  
University of California at Riverside  
U.S.A.

Jie Xiao  
Tsinghua University  
Beijing, China

Jean-Pierre Demailly  
Institut Fourier  
Laboratoire de Mathématiques  
Saint-Martin d'Hères, France

Eduard J.N. Looijenga  
Universiteit Utrecht  
The Netherlands

Neil Trudinger  
Mathematical Sciences Institute  
Australian National University  
Canberra, Australia



# Preface

The main purpose of this book is to describe analytic techniques which are useful to study questions such as linear series, multiplier ideals and vanishing theorems for algebraic vector bundles. One century after the ground-breaking work of Riemann on geometric aspects of function theory, the general progress achieved in differential geometry and global analysis on manifolds resulted into major advances in the theory of algebraic and analytic varieties of arbitrary dimension. One central unifying concept is positivity, which can be viewed either in algebraic terms (positivity of divisors and algebraic cycles), or in more analytic terms (plurisubharmonicity, Hermitian connections with positive curvature). In this direction, one of the most basic results is Kodaira's vanishing theorem for positive vector bundles (1953—1954), which is a deep consequence of the Bochner technique and the theory of harmonic forms initiated by Hodge during the 1940's. This method quickly led Kodaira to the well-known embedding theorem for projective varieties, a far reaching extension of Riemann's characterization of abelian varieties. Further refinements of the Bochner technique led ten years later to the theory of  $L^2$  estimates for the Cauchy-Riemann operator, in the hands of Kohn, Andreotti-Vesentini and Hörmander among others. Not only can vanishing theorems be proved or reproved in that manner, but perhaps more importantly, extremely precise information of a quantitative nature can be obtained about solutions of  $\bar{\partial}$ -equations, their zeroes, poles and growth at infinity.

We try to present here a condensed exposition of these techniques, assuming that the reader is already somewhat acquainted with the basic concepts pertaining to sheaf theory, cohomology and complex differential geometry. In the final chapter, we address very recent questions and open problems, e.g. results related to the finiteness of the canonical ring and the abundance conjecture, as well as results describing the geometric structure of Kähler varieties and their positive cones.

This book is an expansion of lectures given by the author at the Park City Mathematics Institute in 2008 and was published partly in *Analytic and Algebraic Geometry*, edited by Jeff McNeal and Mircea Mustață. It is a volume in the Park City Mathematics Series, a co-publication of the Park City Mathematics Institute and the American Mathematical Society.

Jean-Pierre Demailly  
Université de Grenoble I, Institut Fourier  
Saint-Martin d'Hères  
April 21, 2010



# Contents

<b>Introduction</b> .....	<b>1</b>
<b>Chapter 1. Preliminary Material: Cohomology, Currents</b> .....	<b>5</b>
1.A. Dolbeault Cohomology and Sheaf Cohomology .....	5
1.B. Plurisubharmonic Functions .....	6
1.C. Positive Currents .....	9
<b>Chapter 2. Lelong numbers and Intersection Theory</b> .....	<b>15</b>
2.A. Multiplication of Currents and Monge-Ampère Operators .....	15
2.B. Lelong Numbers .....	18
<b>Chapter 3. Hermitian Vector Bundles, Connections and Curvature</b> ..	<b>25</b>
<b>Chapter 4. Bochner Technique and Vanishing Theorems</b> .....	<b>31</b>
4.A. Laplace-Beltrami Operators and Hodge Theory .....	31
4.B. Serre Duality Theorem .....	32
4.C. Bochner-Kodaira-Nakano Identity on Kähler Manifolds .....	33
4.D. Vanishing Theorems .....	34
<b>Chapter 5. <math>L^2</math> Estimates and Existence Theorems</b> .....	<b>37</b>
5.A. Basic $L^2$ Existence Theorems .....	37
5.B. Multiplier Ideal Sheaves and Nadel Vanishing Theorem .....	39
<b>Chapter 6. Numerically Effective and Pseudo-effective Line Bundles</b> ..	<b>47</b>
6.A. Pseudo-effective Line Bundles and Metrics with Minimal Singularities .....	47
6.B. Nef Line Bundles .....	49
6.C. Description of the Positive Cones .....	51
6.D. The Kawamata-Viehweg Vanishing Theorem .....	56
6.E. A Uniform Global Generation Property due to Y.T. Siu .....	58
<b>Chapter 7. A Simple Algebraic Approach to Fujita's Conjecture</b> .....	<b>61</b>
<b>Chapter 8. Holomorphic Morse Inequalities</b> .....	<b>71</b>
8.A. General Analytic Statement on Compact Complex Manifolds .....	71
8.B. Algebraic Counterparts of the Holomorphic Morse Inequalities .....	72
8.C. Asymptotic Cohomology Groups .....	74
8.D. Transcendental Asymptotic Cohomology Functions .....	78
<b>Chapter 9. Effective Version of Matsusaka's Big Theorem</b> .....	<b>83</b>
<b>Chapter 10. Positivity Concepts for Vector Bundles</b> .....	<b>89</b>
<b>Chapter 11. Skoda's <math>L^2</math> Estimates for Surjective Bundle Morphisms</b> ..	<b>99</b>
11.A. Surjectivity and Division Theorems .....	99
11.B. Applications to Local Algebra: the Briançon-Skoda Theorem .....	105

<b>Chapter 12. The Ohsawa-Takegoshi <math>L^2</math> Extension Theorem</b> .....	<b>111</b>
12.A. The Basic a Priori Inequality .....	111
12.B. Abstract $L^2$ Existence Theorem for Solutions of $\bar{\partial}$ -Equations .....	112
12.C. The $L^2$ Extension Theorem .....	114
12.D. Skoda's Division Theorem for Ideals of Holomorphic Functions .....	122
<b>Chapter 13. Approximation of Closed Positive Currents by Analytic Cycles</b> .....	<b>127</b>
13.A. Approximation of Plurisubharmonic Functions Via Bergman kernels .....	127
13.B. Global Approximation of Closed (1,1)-Currents on a Compact Complex Manifold ...	129
13.C. Global Approximation by Divisors .....	136
13.D. Singularity Exponents and log Canonical Thresholds .....	143
13.E. Hodge Conjecture and approximation of $(p, p)$ - currents .....	148
<b>Chapter 14. Subadditivity of Multiplier Ideals and Fujita's Approximate Zariski Decomposition</b> .....	<b>153</b>
<b>Chapter 15. Hard Lefschetz Theorem with Multiplier Ideal Sheaves</b> .....	<b>159</b>
15.A. A Bundle Valued Hard Lefschetz Theorem .....	159
15.B. Equisingular Approximations of Quasi Plurisubharmonic Functions .....	160
15.C. A Bochner Type Inequality .....	166
15.D. Proof of Theorem 15.1 .....	168
15.E. A Counterexample .....	170
<b>Chapter 16. Invariance of Plurigenera of Projective Varieties</b> .....	<b>173</b>
<b>Chapter 17. Numerical Characterization of the Kähler Cone</b> .....	<b>177</b>
17.A. Positive Classes in Intermediate $(p, p)$ -bidegrees .....	177
17.B. Numerically Positive Classes of Type (1,1) .....	178
17.C. Deformations of Compact Kähler Manifolds .....	184
<b>Chapter 18. Structure of the Pseudo-effective Cone and Mobile Intersection Theory</b> .....	<b>189</b>
18.A. Classes of Mobile Curves and of Mobile $(n - 1, n - 1)$ -currents .....	189
18.B. Zariski Decomposition and Mobile Intersections .....	192
18.C. The Orthogonality Estimate .....	199
18.D. Dual of the Pseudo-effective Cone .....	202
18.E. A Volume Formula for Algebraic (1,1)-Classes on Projective Surfaces .....	205
<b>Chapter 19. Super-canonical Metrics and Abundance</b> .....	<b>209</b>
19.A. Construction of Super-canonical Metrics .....	209
19.B. Invariance of Plurigenera and Positivity of Curvature of Super-canonical Metrics ...	216
19.C. Tsuji's Strategy for Studying Abundance .....	217
<b>Chapter 20. Siu's Analytic Approach and Păun's Non Vanishing Theorem</b> .....	<b>219</b>
<b>References</b> .....	<b>223</b>



# Introduction

This introduction will serve as a general guide for reading the various parts of this text. The first three chapters briefly introduce basic materials concerning complex differential geometry, Dolbeault cohomology, plurisubharmonic functions, positive currents and holomorphic vector bundles. They are mainly intended to fix notation. Although the most important concepts are redefined, readers will probably need to already possess some related background in complex analysis and complex differential geometry — whereas the expert readers should be able to quickly proceed further.

The heart of the subject starts with the Bochner technique in Chapter 4, leading to fundamental  $L^2$  existence theorems for solutions of  $\bar{\partial}$ -equations in Chapter 5. What makes the theory extremely flexible is the possibility to formulate existence theorems with a wide assortment of different  $L^2$  norms, namely norms of the form  $\int_X |f|^2 e^{-2\varphi}$  where  $\varphi$  is a plurisubharmonic or strictly plurisubharmonic function on the given manifold or variety  $X$ . Here, the weight  $\varphi$  need not be smooth, and on the contrary, it is extremely important to allow weights which have logarithmic poles of the form  $\varphi(z) = c \log \sum |g_j|^2$ , where  $c > 0$  and  $(g_j)$  is a collection of holomorphic functions possessing a common zero set  $Z \subset X$ . Following Nadel [Nad89], one defines the *multiplier ideal sheaf*  $\mathcal{I}(\varphi)$  to be the sheaf of germs of holomorphic functions  $f$  such that  $|f|^2 e^{-2\varphi}$  is locally summable. Then  $\mathcal{I}(\varphi)$  is a coherent algebraic sheaf over  $X$  and  $H^q(X, K_X \otimes L \otimes \mathcal{I}(\varphi)) = 0$  for all  $q \geq 1$  if the curvature of  $L$  is positive as a current. This important result can be seen as a generalization of the Kawamata-Viehweg vanishing theorem [Kaw82, Vie82], which is one of the cornerstones of higher dimensional algebraic geometry, especially in relation with Mori's minimal model program.

In the dictionary between analytic geometry and algebraic geometry, the ideal  $\mathcal{I}(\varphi)$  plays a very important role, since it directly converts an analytic object into an algebraic one, and, simultaneously, takes care of the singularities in a very efficient way. Another analytic tool used to deal with singularities is the theory of positive currents introduced by Lelong [Lel57]. Currents can be seen as generalizations of algebraic cycles, and many classical results of intersection theory still apply to currents. The concept of Lelong number of a current is the analytic analogue of the concept of multiplicity of a germ of algebraic variety. Intersections of cycles correspond to wedge products of currents (whenever these products are defined).

Besides the Kodaira-Nakano vanishing theorem, one of the most basic “effective result” expected to hold in algebraic geometry is expressed in the following conjecture of Fujita [Fuj87]: if  $L$  is an ample (i.e. positive) line bundle on a projective  $n$ -dimensional algebraic variety  $X$ , then  $K_X + (n + 1)L$  is generated by sections and  $K_X + (n + 2)L$  is very ample. In the last two decades, a lot of efforts have been brought for the solution of this conjecture — but reaching the expected optimal bounds will probably require new ideas. The first major results are the proof of the Fujita conjecture in the case of surfaces by Reider [Rei88] (the case of curves is easy and has been known since a very long time), and the numerical criterion for the very ampleness of  $2K_X + L$  given in [Dem93b], obtained by means of analytic techniques and Monge-Ampère equations with isolated singularities. Alternative algebraic techniques were developed slightly later by Kollár [Kol92], Ein-Lazarsfeld [EL93], Fujita [Fuj93], Siu [Siu95, 96], Kawamata [Kaw97] and Helmke [Hel97]. We will explain here Siu’s method because it is technically the simplest method; one of the results obtained by this method is the following effective result:  $2K_X + mL$  is very ample for  $m \geq 2 + \binom{3n+1}{n}$ . The basic idea is to apply the Kawamata-Viehweg vanishing theorem, and to combine this with the Riemann-Roch formula in order to produce sections through a clever induction procedure on the dimension of the base loci of the linear systems involved.

Although Siu’s result is certainly not optimal, it is sufficient to obtain a nice constructive proof of *Matsusaka’s big theorem* [Siu93, Dem96]. The result states that there is an effective value  $m_0$  depending only on the intersection numbers  $L^n$  and  $L^{n-1} \cdot K_X$ , such that  $mL$  is very ample for  $m \geq m_0$ . The basic idea is to combine results on the very ampleness of  $2K_X + mL$  together with the theory of holomorphic Morse inequalities [Dem85b]. The Morse inequalities are used to construct sections of  $m'L - K_X$  for  $m'$  large. Again this step can be made algebraic (following suggestions by F. Catanese and R. Lazarsfeld), but the analytic formulation apparently has a wider range of applicability.

In the subsequent chapters, we pursue the study of  $L^2$  estimates, in relation with the Nullstellenatz and with the extension problem. Skoda [Sko72b, 78] showed that the division problem  $f = \sum g_j h_j$  can be solved holomorphically with very precise  $L^2$  estimates, provided that the  $L^2$  norm of  $|f| |g|^{-p}$  is finite for some sufficiently large exponent  $p$  ( $p > n = \dim X$  is enough). Skoda’s estimates have a nice interpretation in terms of local algebra, and they lead to precise qualitative and quantitative estimates in connection with the Bézout problem. Another very important result is the  $L^2$  extension theorem by Ohsawa-Takegoshi [OT87, Ohs88], which has also been generalized later by Manivel [Man93]. The main statement is that every  $L^2$  section  $f$  of a suitably positive line bundle defined on a subvariety  $Y \subset X$  can be extended to a  $L^2$  section  $\tilde{f}$  defined over the whole of  $X$ . The positivity condition can be understood in terms of the canonical sheaf and normal bundle to the subvariety. The extension theorem turns out to have an incredible amount of important consequences: among them, let us mention for instance Siu’s theorem [Siu74] on the analyticity of Lelong numbers, the basic approximation theorem of closed positive  $(1, 1)$ -currents by divisors, the subadditivity property  $\mathcal{I}(\varphi + \psi) \subset \mathcal{I}(\varphi)\mathcal{I}(\psi)$  of multiplier ideals [DEL00], the restriction formula  $\mathcal{I}(\varphi|_Y) \subset \mathcal{I}(\varphi)|_Y, \dots$ . A suitable combination of these results yields an-

other important result of Fujita [Fuj94] on approximate Zariski decomposition, as we show in Chapter 14.

In Chapter 15, we show how subadditivity can be used to derive an “equi-singular” approximation theorem for (almost) plurisubharmonic functions: any such function can be approximated by a sequence of (almost) plurisubharmonic functions which are smooth outside an analytic set, and which define the same multiplier ideal sheaves. From this, we derive a generalized version of the *hard Lefschetz theorem* for cohomology with values in a pseudo-effective line bundle; namely, the Lefschetz map is surjective when the cohomology groups are twisted by the relevant multiplier ideal sheaves.

Chapter 16 explains the proof of Siu’s theorem on the invariance of plurigenera, according to a beautiful approach developed by Mihai Păun [Pău07]. The proof consists of an iterative process based on the Ohsawa-Takegoshi theorem, and a very clever limiting argument for currents.

Chapters 17 and 18 are devoted to the study of positive cones in Kähler or projective geometry. Recent “algebraic-analytic” characterizations of the Kähler cone [DP04] and the pseudo-effective cone of divisors [BDPP04] are explained in detail. This leads to a discussion of the important concepts of volume and mobile intersections, following S. Boucksom’s PhD work [Bou02]. As a consequence, we show that a projective algebraic manifold has a pseudo-effective canonical line bundle if and only if it is not uniruled.

Chapter 19 presents further important ideas of H. Tsuji, later refined by Berndtsson and Păun, concerning the so-called “super-canonical metrics”, and their interpretation in terms of the invariance of plurigenera and of the abundance conjecture. In the concluding Chapter 20, we state Păun’s version of the Shokurov-Hacon-McKernan-Siu non vanishing theorem and give an account of the very recent approach of the proof of the finiteness of the canonical ring by Birkar-Păun [BiP09], based on the ideas of Hacon-McKernan and Siu.